

SUPPLEMENT TO:

A CATALOGUE/CHECKLIST

OF THE

BUTTERFLIES OF AMERICA
NORTH OF MEXICO

Clifford D. Ferris, Editor

The Lepidopterists' Society

Memoir No. 3

1989

**SUPPLEMENT TO:
A CATALOGUE / CHECKLIST OF THE
BUTTERFLIES OF AMERICA
NORTH OF MEXICO**

Clifford D. Ferris, Editor

Director, Bioengineering Program, University of Wyoming

Research Associate, Allyn Museum of Entomology / Florida Museum of Natural History

Research Associate, Florida Collection of Arthropods

Research Associate in Entomology, Natural History Museum of Los Angeles County

The Lepidopterists' Society
Memoir No. 3
1989

© 1989 by the Lepidopterists' Society
Library of Congress Catalog Card Number: 89:080153
ISBN: 0-930282-04-3
Printed in the United States of America

Typesetting by C. D. Ferris on Macintosh™ Plus computer using Times type font and Apple®
LaserWriter™.

MEMOIRS EDITOR'S NOTE

During the eight years since the publication of *Memoir No. 2*, classification and nomenclature of the North American butterflies has undergone a period of instability. This was predicted at the end of my Editor's Notes in that volume, and is reflected in the varying usages employed by the authors of several major books on North American butterflies that have been published during the interim, plus a number of talks at Annual Meetings and shorter published works.

The present volume reflects nomenclatural changes published since *Memoir No. 2* came out, plus some corrections of errors in that work. A few emendations dictated by the 1985 revision of the *International Code of Zoological Nomenclature* appear here, and further valuable notes are also included. The Lepidopterists' Society owes a debt of gratitude to Dr. Clifford D. Ferris and the other members of the Review Committee (listed on p. v), as well as to the many of our colleagues who have contributed to the substance of this Supplement.

As I stated in *Memoir No. 2*, the search for a more complete and perfect classification system of butterflies continues, and the employment of modern approaches will certainly result in further and, hopefully, well-supported changes in years to come. Increasingly systematists employ such techniques as cladistic analysis, electron microscopy, physiological experimentation, and new life history information to base decisions on more sound footing than before. Further, increased international communication among scholars and the resulting association of our butterfly fauna with that beyond our borders should bring Lepidoptera classification and nomenclature to a new level of worldwide uniformity and stability in years ahead.

In 1989 Dr. William E. Miller will assume the duties of Editor of the *Memoirs*. Bill has already distinguished himself as a tireless and exacting Editor of our *Journal*, and has exercised his duties with skill, timeliness, and courage. I wish to thank the members of the *Memoirs* Advisory Board and the Executive Council for the help they gave me during my tenure as Editor. Intended for long works of broad interest to lepidopterists, the *Memoirs* should be expanded in future to include works other than checklists, and I feel it will be. I want to wish Bill Miller great success in making the *Memoirs* a more frequent and versatile component of our Society's publishing program.

I urge you to support your Society in every way, including suggestions to the Editor for types of works you would like to see appear as *Memoirs*.

Charles V. Covell, Jr.
Editor of the *Memoirs*

PREFACE

A Catalogue/Checklist of the Butterflies of America North of Mexico (*Memoir No. 2* of the Lepidopterists' Society) by Lee D. Miller and F. Martin Brown was published in 1981. At the 34th Annual Meeting of the Lepidopterists' Society, held in Columbus, Ohio in July of 1983, the Executive Council appointed a five-member committee charged with the review of *Memoir No. 2* for errors and omissions. This committee was also instructed to bring the publication up to date regarding new taxa and taxonomic changes in existing nomenclature. It is anticipated that this latter function will be on a continuing basis with periodic notes published in either the *News* or the *Journal of the Lepidopterists' Society*. This initial revision, however, is being published as a separate Supplement to *Memoir No. 2*.

The Committee as finally constituted (one initial appointee declined to serve) is composed of the following:

- Dr. Clifford D. Ferris, Chairman, University of Wyoming.
Pieridae; nymphaloid butterflies, general editing.
- Dr. F. Martin Brown, Colorado Springs, Colorado.
Heliconiidae; Melitaeinae; Danaidae.
- Dr. David L. Hancock, Brisbane, Australia.
Papilionidae.
- Dr. Lee D. Miller, Allyn Museum of Entomology/Florida Museum of Natural History.
Hesperioidae; Satyridae.
- Col. Stanley S. Nicolay, Virginia Beach, Virginia.
Lycaenidae; Riodinidae.

The material presented herein is a synthesis of the efforts of the individual committee members and contributions from numerous individuals representing the lepidopterological community at large, many of whom are recognized in the "Acknowledgments." Although not formally members of the Committee, three individuals have made major contributions to this project and deserve special recognition. They are Dr. Jacqueline Y. Miller, Jonathan P. Pelham, and Steven R. Steinhauser.

It must be emphasized that the geographic region to which *Memoir No. 2* pertains is North America north of Mexico, and not the political entity represented by the United States of America and its protectorates. For this reason, the fauna of Hawaii, Puerto Rico, etc. are not included. Some readers of *Memoir No. 2* have misunderstood its geographic coverage.

An attempt has been made to include all species for which there are any reasonable records of occurrence in the region to which *Memoir No. 2* applies. Some of these records date to the 19th Century and may be erroneous, but museum specimens exist with appropriate locality labels. One must recognize that many environmental changes have occurred in North America during the past century. Distribution patterns have shifted accordingly, and several species are either known to be or are presumed to be extinct; yet we have museum material. Early lists, especially those of Boisduval, erroneously included names of Old World and Asiatic species. These names have generally been omitted from *Memoir No. 2* and this Supplement.

ACKNOWLEDGMENTS

The Committee wishes to thank the following individuals who supplied information used in this revision: P. H. Arnaud, Jr. (San Francisco, CA), G. T. Austin (Las Vegas, NV), K. Bagdonas (Laramie, WY), M. D. Bowers (Cambridge, MA), C. A. Bridges, (Urbana, IL), J. M. Burns (Washington, DC), C. F. Cowan (U.K.), J. P. Donahue (Los Angeles, CA), J. F. Emmel (Hemet, CA), Hans J. Epstein (Switzerland), H. A. Flaschka (Decatur, GA), T. P. Friedlander (Baton Rouge, LA), L. P. Grey (Lincoln, ME), D. W. Jenkins (Sarasota, FL), K. Johnson (New York, NY), O. Kudrna (Germany), J. D. Lafontaine (Canada), B. Mather (Clinton, MS), J. S. Miller (New York, NY), D. J. Murphy (Stanford, CA), N. F. Nash (Urbana, IL), Y. P. Nekrutenko (U.S.S.R.), C. G. Oliver (Scottdale, PA), K. W. Philip (Fairbanks, AK), J. E. Rawlins (Pittsburgh, PA), F. H. Rindge (New York, NY), R. K. Robbins (Washington, DC), C. W. Sabrosky (Washington, DC), J. A. Scott (Lakewood, CO), A. M. Shapiro (Davis, CA), S. Shaw (Cambridge, MA), R. E. Stanford (Denver, CO), J. W. Tilden (San Jose, CA), M. E. Toliver (Eureka, CA), T. W. Turner (Safety Harbor, FL), M. Williams (Urbana, IL). Several anonymous reviewers of the completed draft made valuable suggestions.

In addition to members of the Committee, C. V. Covell, Jr and colleagues (Louisville, KY), J. C. Downey (Sarasota, FL), T. P. Friedlander (Baton Rouge, LA), D. W. Jenkins (Sarasota, FL), and S. R. Steinhauser (Sarasota, FL) graciously assisted in checking sections of the galley proof.

INTRODUCTION

This Supplement to Memoir No. 2 is divided into three parts. The first part consists of some general notes relative to the family-group names and certain literature citations. Much of this information was kindly supplied by the late C. F. Cowan, with additional notes by C. A. Bridges.

The second part contains corrections, emendations, general notes, and additions to the original text of *Memoir No. 2*. Relative to the format, the number code appearing in the left margin refers to page number and the corresponding genus/species entry number. Thus: 84:367 is interpreted as page 84, species entry 367 (*Zerene cesonia*); 85:CXI is interpreted as page 85, genus entry CXI *Phoebeis*.

Because of the numerous changes that have been made in the original text of *Memoir No. 2*, as shown in this Supplement, a rudimentary checklist of the butterflies of North America north of Mexico has been included as Part III. All primary taxa are included, but synonomies, form, and aberrational names have been omitted. This list serves in lieu of an index, and it may also be used in a manner similar to the "life lists" used by birders.

The use of the term CODE refers to the *International Code of Zoological Nomenclature* (February, 1985) adopted by the XX General Assembly of the International Union of Biological Sciences. The permanent body which is the authority on zoological nomenclature is the International Commission on Zoological Nomenclature, herein abbreviated as I.C.Z.N. In this regard, it should be noted that the CODE is revised periodically. Changes in the rules, however, are for the most part not retroactive (an exception is the requirement for genus-species-name gender agreement and certain matters of spelling). Thus actions taken by authors working under the guidelines of previous editions of the CODE are not countermanded when a new edition of the CODE is issued. This condition must prevail so that nomenclatural stability is preserved, which is one of the goals of the CODE.

Editor's Note

Since 1986, several regional books and a general book on North American butterflies have been published. These include: Tilden and Smith, 1986, *A Field Guide to the Western Butterflies*, Houghton Mifflin Co., Boston; Garth and Tilden, 1986, *California Butterflies*, University of California, Berkeley; Heitzman and Heitzman, 1987, *Butterflies and Moths of Missouri*, Missouri Department of Conservation, Jefferson City; Shull, 1987, *The Butterflies of Indiana*, Indiana University Press, Bloomington; Royer, 1988, *Butterflies of North Dakota*, Minot State University; Scott, 1986, *The Butterflies of North America*, Stanford University Press. To varying degrees, the nomenclature used in these books differs from that used in *Memoir No. 2*, and from one another. In some cases, taxonomic changes have occurred since 1981 when *Memoir No. 2* was published, and these are reflected in these books and in this Supplement. Several of the authors mentioned have elected to use the "super genus" approach, for example the use of the single generic name *Papilio* rather than the generic names adopted in *Memoir No. 2* and this Supplement.

The nomenclature used in this Supplement reflects the opinions of the Committee members as to what is appropriate usage at this time. We recognize that nomenclature must be flexible to the extent that it can adapt as our knowledge of the butterflies expands.

A brief comment is in order concerning the nomenclature used in the Scott book. With regard to the boldface type font used therein, species, subspecies and form names are undifferentiated. Many taxonomic changes and new combinations of names appear without any supporting discussion, and for this reason are not adopted herein.

PART I

GENERAL NOTES

1. The method of citing dates in *Memoir No. 2* does not conform to the recommendations given in Art. 22 of the CODE. Recommended format is:

Papilio sennae Linnaeus, 1758. [Literature citation]. This form is used when the date of publication is clearly specified, and the name as originally proposed has not been revised.

Phoebeis sennae (Linnaeus, 1758). [Literature citation]. This form is used when a reviser has altered the generic name, but the date of publication is clearly specified; () used in this case.

Papilio argyrogynomon Bergsträsser, [1779]. [Literature citation]. In this example, the date was not clearly specified and has been determined only from external evidence. [] used in this case.

Lycaeides argyrogynomon (Bergsträsser, [1779]). [Literature citation]. In this instance, both () and [] are used, since the date of publication is clouded and the generic name has been changed.

The nature of this note is purely informative; no attempt has been made to change the existing text, although this format has been followed in new and revised entries, except when split as in: *Phoebeis sennae* (Linnaeus) *PAPILIO*, 1758 [to indicate the genus in which the species name was originally described]. In some instances a date is set in " " to indicate imprinted date of publication when the actual date of publication is different.

2. Dating of the early literature is somewhat uncertain. The late C. F. Cowan has suggested the following revisions to references cited in the text of *Memoir No. 2*, and which represent the current status of this material. The numbers set apart in { } in this and the sections that follow indicate respectively the page and species entry numbers of the associated reference. Only the page numbers are indicated for family-group names. Available names appear in the *Official Lists* (and subsequent Opinions) published by the I.C.Z.N. See ¶ 15 on page 4.

PAPILIONIDAE {55}; **PAPILIONINAE** {59}; **Papilionini** {61}: Latreille, [1802]. Hist. Nat. Crust. Ins. 3: 387, "Ann. X", but [Nov. 1802].

Parnassiinae; **Parnassiini** {55}: Duponchel, [1835]. In Godart, Hist. nat. Lépid. France. Suppl. 1 (Livraison 22): 380; 1832-[1835].

PIERIDAE; **Pierinae**; **Pierini** {68}: Duponchel, [1835]. In Godart, Hist. nat. Lépid. France. Suppl. 1 (Livraison 22): 381; 1832-[1835].

DISMORPHIINAE {91}: This same name was used earlier by Schatz, 1886. In Staudinger and Schatz, Exot. Schmett. 2 (Die Familien und Gattungen. . .): 56, 57. An even yet earlier name is **LEPTALINAE**: Bar, 1878 (as **LEPTALIDES**). Ann. Soc. ent. France (5) 8: 12, 23.

MLETINAE Corbet, "1939" {91}: Since this name is valid under Art. 40(a) of the CODE, it should be cited as: Corbet, 1939 (1886) in accordance with Art. 40(b) 1886 being the date of the superseded **GERYDINAE** Doherty.

RIODINIDAE {127}; **Riodininae** {128}: Grote, 1895 (1827). Mitt. Roemerius. Hildesheim, (1): 1; 1895 (1827), by I.C.Z.N. Opinion 1073.

LIBYTHEIDAE {132}: Boisduval, 1833. Nouv. Ann. Mus. H. N. Paris (2) 2: 200 [reprinted as Faune Entom. Madag., Bourbon et Maurice (Lépid.): 52].

Argynninae {135}: Duponchel, [1835]. In Godart, Hist. nat. Lépid. France. Suppl. 1

(Livraison 23): 394; 1832-[1835].

MELITAEINI; *Melitaeini* {152}: An earlier usage is: Tutt, 1896. Ent. Rec. J. Var. 7: 301.

Nymphalini {170}: An earlier usage of Vanessini was: Duponchel, [1835]. In Godart, Hist. nat. Lépid. France. Suppl. 1 (Livraison 23): 397; 1832-[1835].

Hypolimnini {175}: This name is invalid under Art. 60(a) of the CODE. A suggested replacement is: *Junoniini* Reuter, 1836. Acta Soc. Sci. Fenn. 22: 554.

Epicaliini {180}: Guenée, 1865. In Vinson, Voy. Madagascar, Annexe F: 28.

Eurytelini {182}: An earlier usage is: *Biblidini* Boisduval, 1833. Nouv. Ann. Mus. H. N. Paris (2) 2: 201. [Reprinted as Faune Entom. Madag., Bourbon et Maurice (Lépid.): 53].

Ageroniini {183}: Doubleday, 1847. Gen. diurn. Lepid. 1(8): 81.

Marpsiinae {185}: An earlier usage is: *Cyrestinae* Guenée, 1865. In Vinson, Voy. Madagascar, Annexe F: 180.

Charaxinae {186}: Guenée, 1865. In Vinson, Voy. Madagascar, Annexe F: 28.

DANAIDAE; *Danainae* {203}: Boisduval, 1833. Icones Hist. Lépid. Europe 1 (Livraison 9): 84; 1832-[1834].

3. The name Guérin-Méneville should be spelled with only two acute accent marks rather than the three shown throughout the text. Regarding his publications, the *Iconog. Règne Anim. Cuvier* was issued in 45 livraisons (parts) of 450 plates of all orders from 1829-1838. The complete text followed in 1844. Consequently all page references must be [1844], and all plate references must be earlier:

{94:398} *thoe*: pl. 81, 1831; text [1844].

{128:528} *virginiensis*: pl. 81, 1831; text [1844].

{202:757} *balder*: pl. 80, 1831; text [1844].

Griffith's *Anim. Kingdom Cuvier* commenced publication in 1824, but the insect portion began in 1831. Griffith translated Cuvier's work from the French, and guest authors added supplementary chapters in which new species were introduced. Thus volumes 14 and 15 of the Insects (in 3 and 4 parts, respectively) were direct translations from Latreille in Cuvier's edition 2, vols. 4 and 5 (1829-30), illustrated by plates exactly copied as mirror images from Guérin-Méneville's *Iconographie*, which had just begun to appear. A supplementary chapter by G. R. Gray, with its own original plates, followed each insect order. Gray's concept of *Erycina* can be evaluated by his own plate 102, with three figures of *Erycina* species, one of which is *Erycina (Lamproptera) curius* (Papilionidae). Griffith copied six of Guérin-Méneville's butterfly plates. Griffith's plates 3, 21, 2, 43, 47, 58 are exact copies of Guérin-Méneville's plates 76, 77, 78, 79, 80, 81 respectively. Guérin-Méneville's plate 80 introduced the name *balder*, and plate 81 the names *thoe* and *virginiensis*, all three being "Boisduval" manuscript names. The full legends of Guérin-Méneville's plates were faithfully copied in those of Griffith. Gray had no part in this, although these names have at times been attributed to him. Gray's only involvement in this was in the index to the last (seventh) part of the insects. He gave a short account of the colors of each insect: "for the sake of those without coloured plates." This was issued well after the plates appeared. Even if Griffith had inadvertently issued a copied plate before Guérin-Méneville's original had appeared, the name should be attributed to Guérin-Méneville and not to Gray.

4. The name Le Conte is variously misspelled Leconte, etc. in the text.
 5. Several Boisduval citations should be corrected as follows:

{188}: Satyridae; {190}: Satyrinae; {199}: Satyrini; {200}: Oeneis; 202:758a, b. Icones Hist. Lepid. Europe 1: [page numbers as shown].
 {17}: Achlyodes; 61:301; 65:322; 66:322; 67:324; 68:328; 71:336d; 73:339a; 82:360, a; 87:347, a; {91}: Nathalis; {128}: Euselasia; {132}: LIBYTHEIDAE (see note above about prior usage); 233:500. [1834]. (*In Roret*). Hist. Nat. Ins., Sp. Gén. Lépid. [page or plate citation as shown].

Regarding C. F. Cowan's research into the Boisduval publication dates, the title page of the second book shows the following: "Icones Historique des Lépidoptères Nouveaux ou peu Connus. Collection, avec Figures colorées, des Papillons d'Europe nouvellement découverts; Ouvrage formant le Complément de tous les autres Iconographies; par Le Docteur Boisduval. Paris, A La Librairie Encyclopédique de Roret, 1832." The situation seems to be that signatures of the book were issued successively over a period of several years, and then eventually bound as a whole.

6. Regarding Godart's Enc. Meth.: this publication was issued in two parts; Part 1 to page 328 in 1819; Part 2, pp. 329-828 [1824]. The Supplement was not a separate issue, but rather a portion of Part 2 and began with p. 804. Godart's "Mem. Soc. Linn. Paris 2, 1822" was never published and one copy (at least) exists in proof form at the MNHN in Paris. Mem. Soc. Linn. Paris 1 for the year 1821 was published in 1822; Mem. Soc. Linn. Paris 3 for 1823 was published in 1825. Thus for 60:299, the citation should read: [1824]. Enc. Meth. 9(2): 810.
7. "Johansson", 1763: present day botanists and zoologists are in general agreement that in the time of Linnaeus it was customary for the *Praeses* (Linnaeus) to write the thesis, and for the pupil (Johansson) to propose and defend it. Consequently all new names are credited to the *Praeses*. Thus the references to "Johansson" should be emended to read: Linnaeus (*in Johansson*), 1763; 86:372, for example. Further discussion of this matter is found in Marshall [1983]. Zool. J. Linn. Soc. 78: 375-396].
8. Art. 31 of the CODE clearly states that a species-group name must agree in gender with the generic name with which it is at any time combined, hence such changes as *Eurema mexicana* to *Eurema mexicanum* [89:380 and elsewhere].
9. Members of the Committee have received inquiries as to why the nomenclature in the *Check List of the Lepidoptera of America North of Mexico* edited by R. W. Hodges differs from the nomenclature in *Memoir No. 2*, when Miller and Brown also authored the butterfly section in the *Check List*. Although *Memoir No. 2* was published in 1981 and the *Check List* in 1983, because of publication delays, the latter is actually an earlier edition of the text of *Memoir No. 2*. Nomenclatural changes were made in *Memoir No. 2* that do not appear in the *Check List*. Other changes relate to Note 8 immediately above.
10. C. F. Cowan bibliography (papers related to dates of publication of butterfly names and related subjects).

Cowan, C. F., 1969a. Boisduval's *Species Général des Lépidoptères*. J. Soc. Biblphy nat. Hist. 5(2): 121-122.

_____, 1969b. Boisduval and Le Conte: *Histoire Générale et Iconographie des Lépidoptères et des Chenilles de l'Amérique Septentrionale*. J. Soc. Biblphy nat. Hist. 5(2): 125-134.

_____, 1969c. Notes on Griffith's *Animal Kingdom of Cuvier* (1824-1835). J. Soc. Biblphy nat. Hist. 5(2): 137-140.

_____, 1969d. On the *Mémoires de la Société Linnéenne de Paris*, 1822-27. J. Soc.

- Biblphy nat. Hist. 5(2): 152-154.
- _____, 1970a. Boisduval's *Icones des Lépidoptères d'Europe "1832"* [-1841]. J. Soc. Biblphy nat. Hist. 5(4): 291-302.
- _____, 1970b. Boisduval *et al.*, *Collection . . . des Chenilles d'Europe "1832"* [-1837]. J. Soc. Biblphy nat. Hist. 5(4): 303-323.
- _____, 1971. On Guérin's *Iconographie*: particularly the insects. J. Soc. Biblphy nat. Hist. 6(1): 18-29.
- _____, 1976. On the Disciple's Edition of Cuvier's *Règne Animal*. J. Soc. Biblphy nat. Hist. 8(1): 32-64.
11. Regarding the Gunder names contained in *Memoir No. 2*, frequently the statement appears: "HT should be in AMNH." The Gunder types are in fact at AMNH and a list of those type specimens was published by dos Passos [1938. American Mus. Novitates (999): 1-16].
12. Several publications by Ahmet Ö. Koçak which appeared in the journal Priamus have come to the attention of the Committee [1981. 1(2): 93-96; 1983. 3(1): 3-5; 1984. 3(3): 93-97]. In these articles, Koçak has proposed corrections in nomenclature at the family and species level regarding several entries in *Memoir No. 2*. Where appropriate, changes have been made in this Supplement. In other instances, Koçak has apparently disregarded several I.C.Z.N. Opinions, and his emendations are not followed.
13. The abbreviations used to designate several museums have changed in the past few years (USNM to NMNH, for example). To avoid confusion, the same abbreviations as were used in *Memoir No. 2* are retained herein.
14. Methods of bibliographic citation vary. The journal and serial abbreviations used herein are consistent with those in general usage, with the exception that national names are spelled out (Canadian Ent. as opposed to Can. Ent.) as was done in *Memoir No. 2*.
15. A useful reference for persons seriously interested in matters of nomenclature is:
Melville, R. A. and J. D. D. Smith, eds. 1987. *Official Lists and Indexes of Names and Works in Zoology*. Published by the International Trust for Zoological Nomenclature, c/o British Museum (Natural History), London, UK.
16. Charles A. Bridges of Urbana, IL has published (1984) a useful concordance to *Memoir No. 2* in which appear indices to original descriptions, journals and serials, and a general bibliography. The title of this work is: *Notes on Species-group names mentioned in the Miller & Brown Catalogue/Checklist of the Nearctic Rhopalocera (Lepidoptera)*. This publication may be purchased from the author, who is a member of the Lepidopterists' Society.

PART II

REVISIONS AND EMENDATIONS TO MEMOIR NO. 2

Editorial Note: The section on the HESPERIOIDEA was prepared by the Editor with substantial input from Jonathan P. Pelham and Ray E. Stanford. It was reviewed and emended by L. D. and J. Y. Miller, R. E. Stanford, and S. R. Steinhauser. In many of the entries in *Memoir No. 2* where the taxon was described by W. H. Edwards, the comment is found with respect to type locality and/or type specimens: "designated by F. M. Brown and L. Miller, in press". The related publications are: *Trans. American ent. Soc.*, 106: 43-88, 1980; 113: 29-71, 1987.

Superfamily HESPERIOIDEA Latreille

- 1:1 Add at end of prefatory material: = *Apyrrhothrix* Evans, 1951. Cat. American Hesp. Brit. Mus. (N.H.): 8. *Lapsus calami*.
- 1:1a Correct museum citation to: BM.
- 2:3 Correct third line of entry: = *polybius* (Fabricius), etc.
- 2:3a Delete [] from [1866] entry in 2nd line.
- 2:4 Correct museum citation for *socius* and *decolor* to: BM.
- 2:4 Correct museum citation in 2nd line to: BM.
- 4:7 Add as follows:
 c. c. *californicus* MacNeill [*nec* J. B. Smith, 1891], 1975. In Howe, 7c.
 Butts. N. America: 572. TL - China Flat, El Dorado Co, California. LT
 in CAS. Although this name has been attributed to J. B. Smith, apparent-
 ly he never published it.
- 4:V Revise 3rd line of entry to read: (= *Papilio leo* Gmelin, 1790. Syst. Nat. ed. 13, 1: 2363; subspecies).
 Lines 12 & 13, delete entry: (= *Papilio leo* ... 1790).
 Revise entry to read:
 *1. *leo* (Gmelin), etc. Delete 9 at end of line.
 a. l. *histrio* Röber, 1925. Ent. Mitteil., 14: 161. TL - "Panama". Type 9a.
 probably was in Dresden.
 = f. "arizonensis" (Skinner) *ERYCIDES*, 1911. Etc.
 = f. "pallida" Röber, 1925. Etc.
- Editorial Note: Evans [1955. Cat. American Hesp. in BM (N.H.), part IV, Appendix II, p. 476] has stated: "Comstock 1944 assigns the locality of *lividus* Hübner to Hispaniola: the name should therefore be removed from the synonyms of *leo* and used as a subspecies in place of *ishmael*, which becomes a synonym of *lividus*."
- 6:17 To conform to the CODE, emend spelling to: *mexicana*.
- 6:18 Add to entry:
 = *maculata*, Seitz, 1925. Grossschmett. Erde, 5: pl. 161d. *Lapsus calami*.
- 6:19 Correct end of 1st line of *procerus* entry to read: ... 55: 8 19.
 To conform to the CODE, emend spelling to: *procera*.
- 7:XII Add as follows in *Goniurus* entry:
 = *Goniurus* Westwood, 1852. Gen. diurn. Lepid., (2): 510. *Lapsus calami*.

- = *Goniarus* Mabille 1891. Bull. C. R. Soc. ent. Belgique, 35: 9.
Lapsus calami.
- 7:24 Add to entry: Neotype male designated by S. R. Steinhauser, 1981 [Bull. Allyn Mus., (62): 13] as follows: MCAS, Cherrypoint, Craven Co., N[orth] C[arolina]. NT in AME.
- 7:25 Replace *pronta* Evans with *pronus* Evans as follows:
 2. *pronus* Evans, 1952. Cat. American Hesp. Brit. Mus. (N.H.), 2: 89.
 TL - Ambato, Ecuador. HT in BM.
 = *plinius* Bell, 1956. American Mus. Novitates (1778): 1-5. TL - Santa Cruz, Bolivia. HT in AMNH.
- 9:33 Delete this entry; species does not occur in our region. This record (by Greg Nielson; March, 1972) was reported in McGuire and Rickard [1974. Annotated Check List of the Butterflies. Bentsen-Rio Grande Val. St. Pk. and Vic. Texas Parks and Wildlife Dept., Mission, TX. Entry 223.1]. The record proved to be a misidentification of *U. doryssus* (*fide* W. W. McGuire, *in litt.*).
- 9:XIII Modify end of 2nd line as follows: (1875 *Papilio aulestes recte aulestis* Cramer, [1780]. Util. Kapellen, 3: 161, etc. See: Steinhauser, 1986 [Bull. Allyn Mus., (104): 1-2].
- 10:XIV Add species as follows:
 3. *cinctus* (Plötz) *CECROPTERUS*, 1882. Berl. ent. Zeit., 26: 261. TL - 41.1.
 "Oaxaca", Mexico. Loc. of type unknown.
 = *cinctus* (Godman and Salvin) *CECROPTERUS*, 1894. Biol. Centrali-Americanana, Lepid. Rhop., 2: Tab. 80, fig. 3, emendation of *cincta* (Plötz), 1882.
 = *rotundatus* (Mabille) *TELEGONUS*, 1883. Bull. Soc. ent. Belgique, 27: 52. TL - "S. America". ST's may be in BM.
- 11:46 Correct date entry: 1881[1882].
- 11:48 Scott [1986. Butts. N. America] has elevated f. "albosuffusa" to subspecific status. Since this phenotype occurs to some extent in many *pylades* populations in western North America, we retain its form status.
- 11:50 To conform to the CODE, emend spelling to: *mexicanus*.
- 11:50a It is doubtful that nominate *mexicanus* occurs in our region.
- 12:50 Add as follows:
 d. m. *blanco* Scott, 1981. Papilio (new series), 1: 7. TL - Crooked Creek 50d. Lab., 10,150', 3 airline mi. N. Inyo Co. line, Mono Co., California. HT in California Insect Survey, Berkeley, California.
- 12:51 Add to entry:
 = *confusus* Auct. *Lapsus calami*.
- 12:53 To conform to the CODE, emend spelling to: *valerianus*.
 There is considerable uncertainty about this taxon, and apparently *valerianus* does not occur in our region. It may or may not be a synonym of *mysie*. Tilden [1949. Bull. S. Calif. Acad. Sci., 48: 4-6; 1975. J. Lepid. Soc., 60-61], however, reviewed the taxonomy of *mysie* Dyar and concluded that it is a valid species in the genus *Phoedinus* Godman and Salvin. Several recent publications have followed this arrangement. This genus was placed by Evans in his Catalogue (1953) as a synonym of *Cogia* Butler. The type-species of *Phoedinus* is *Eudamus caicus* Herrich-Schäffer, 1869 which is now placed as *Cogia caicus*. On this basis then, the taxon *mysie* should be placed under *Cogia*. See entry 14:XXI below.

- 12:XVIII Correct spelling to: *Ancistrocampta*.
 13:XXI The comments in Note 36 are superseded by Cowan, 1970. Ann. Rhop.: 53.
 14:62a It is doubtful that nominate *caicus* occurs in our area.
 14:XXI Add new species as follows:
 4. *mysie* (Dyar) *THORYBES*, 1904. J. New York ent. Soc., 12: 40. TL - 62.1.
 Patagonia Mtns., Arizona. HT in USNM. This species has been collected
 in southern Arizona. See comment above under entry 12:53.
 14 New entry as follows:

Genus XXIA: *Arteurotia* Butler and Druce

1872. Cist. Ent., 1: 112. Type-species by original designation: *Arteurotia tractipennis* Butler and Druce, 1872. Cist. Ent., 1: 112.
 1. *tractipennis* Butler and Druce, 1872. Cist. Ent., 1: 112. TL - Costa Rica. HT in BM.
 a. t. *tractipennis* Butler and Druce, 1872. Cist. Ent., 1: 112. TL - Costa Rica. HT in BM.
 = *ribbei* (Staudinger) *HELIAS*, 1875. Verh. zool.-bot. Ges. Wien, 25: 117. TL - Chiriquí, Panama. LT may be in ZMHU.

- 15:XXIII Add as follows:
 3. *dimidiata* Herrich-Schäffer, 1870. CorrespBl. zool.-min. Ver. Regensburg, 24: 160. TL - "Mexico". Loc. of type unknown.
 a. d. *dimidiata* Herrich-Schäffer, 1820. CorrespBl. zool.-min. Ver. Reg. ensburg, 160. TL - "Mexico". Loc. of type unknown.
 = *didia* Möschler, 1876. Verh. zool.-bot. Ges. Wien., 26: 340. TL - "Surinam". Type may be in Vienna.
 = *corinna* Plötz, 1882. Berl. ent. Zeit., 26: 254. TL - "Mexico". Type may be in Munich.
 = *nivonicus* (Plötz) *ACHLYODES*, 1884. Jahrb. nass. Ver. Nat., 37: 14. TL - "Mexico". Type may be in Munich.
 = *bilinea* Mabille, 1889. Le Naturaliste 1889: 216. TL - "Chiriquí", Panama. Type should be in MNHN.
 = *bobae* (Weeks), *PAPIAS*, 1906. Canadian Ent., 38: 203. TL - Suapure, Venezuela. HT may be in MCZ.
 = *hypsiptyle* Hayward, 1947. Acta zool. lilloana, 3: 121. TL - "Columbia". HT in Museo Argentino de Ciencias Naturales, Buenos Aires.

- 16:XXV Add as follows:
 4. *azteca* (Scudder) *PHOLISORA*, 1872. 4th Ann. Rept. Peabody Acad. Sci. [1871]: 72. TL - Tehuantepec, Oaxaca, Mexico. HT should be in MCZ.

- 16:74 Add to entry:
 = *trixus* Auct. *Lapsus calami*.

Editorial Note: In the text of Uitlandsche Kapellen 4:87, Stoll referred to "Tryxus" and stated that it belongs "aux Plebeiens nobles" or "onder de Dikkop-Kapellen". dos Passos [1960. J. Lepid. Soc., 14(1): 24-36] considered similar use in the text by Cramer to be a *nomen nudum* and used the index spelling. If one accepts this reasoning, then the index spelling *trixus*

would apply with the date 1784 rather than 1780 [see Brown, 1941. Ann. ent. Soc. America, 34(1): 128].

17 Insert genus entry as follows:

Genus XXIXA *Antigonus* Hübner

[1891]. Verz. bekannt. Schmett. (7): 108. Type-species by monotypy: *Urbanus erosus* Hübner, [1812]. Samml. exot. Schmett., 1: pl. [153].

1. *emorsus* (R. Felder) *LEUCOCHITONEA*, 1869. Verh. zool.-bot. Ges. 74.5. Wien, 19: 479. TL - "Mexico". HT in BM.

= *albimedia* (Draudt) *SYSTASEA*, [1922]. In Seitz, Grossschmett. Erde 5: 904. TL - "Mexico". Loc. type not known.

18:78 To conform to the CODE, emend spelling to: *stigmatica*.

18:81a The synonym *llano* was treated as a separate subspecies by Durden [1982. J. Lepid. Soc., 36(1): 8].

19:82 To conform to the CODE, emend spelling to: *brunneus*.

20:85a Add:

= *juvenis* (Hübner) *NISONIADES*, [1819]. Verz. bekannt. Schmett., (7): 108.

21:93c To conform to the CODE, emend spelling to: *perniger*.

21:99a Add:

= *cervantes* (Boisduval) *THANAOS*, 1852. Ann. Soc. ent. France (2)10: 310. TL - "California". HT may be in USNM or BM.

22:101 Change entry number to 101a and emend entry as follows:

TL of *ruralis* restricted by Scott, 1981 [Papilio (new series), 1: 7] to Tuolumne Co., California.

23:101 Add:

TL of *caespitatis* restricted by Scott, 1981 [Papilio (new series), 1: 7] to Marin Co., California.

Add entry 101b as follows:

b. r. *lagunae* Scott, 1981. Papilio (new series), 1: 7. TL - N. end, E. La- 101b. guna Mtns., San Diego Co., California. HT in LACM.

24:110 To conform to the CODE, emend spelling to: *lavianus*.

25:116 To conform to the CODE, emend spelling to: *mejicana*.

25:XLII Delete reference to *Hesperopsis* and split it out as a separate genus with species as follows:

Genus XLIA *Hesperopsis* Dyar

1905. J. New York ent. Soc., 13: 118. Type-species by original designation: *Thanaos alpheus* W. H. Edwards, 1876. Trans. American ent. Soc., 5: 206. See Stanford, 1980 [*in Ferris and Brown, Butts. Rocky Mtn. States: 87*].

1. *libya* (Scudder) *HETEROPTERUS*, 1878. Etc.

a. l. *libya* (Scudder) *HETEROPTERUS*, 1878. Etc. 117a.

b. l. *lena* (W. H. Edwards), *ANCYLOXYPHA*, 1882. Etc. 117b.

2. *alpheus* (W. H. Edwards) *THANAOS*, 1876. Etc.

= *alphaeas* Dyar, 1905. *Lapsus calami*.

a. a. *alpheus* (W. H. Edwards) *THANAOS*, 1876. Etc. 118a.

b. a. *oricus* (W. H. Edwards) *PHOLISORA*, 1879. Etc. 118b.

- c. a. *texanus* Scott, 1981. *Papilio* (new series), 1: 8. TL - Boca Chica, 118c.
20-22 mi. E. Brownsville, Cameron Co., Texas. HT in LACM.
3. *gracielae* (MacNeill) *PHOLISORA*, 1970. Etc. 119.
- 26:XLII Add at end of genus entry:
= *Pamphila* W. J. Holland [*nec* Fabricius], 1898. *Butterfly Book:*
342.
- 26:120a Add:
= *mackenziei* Wyatt, 1965. *Zeit. wien ent. Ges.*, 50: 70. TL - ???
HT presumed to be in Wyatt Collection in Karlsruhe Museum, West
Germany.
- 27:123 To conform to the CODE, emend spelling to: *microstictus*.
Editorial Note: There is considerable confusion about this taxon. No re-
vision is presented herein, but the reader is referred to: Freeman, 1979 [Bull.
Allyn Mus., (52): 5-6] for the discussion of *Piruna mexicana* Freeman;
Scott, 1986 [Butts. N. America: 426] for his discussion of *Piruna cingo*
Evans.
- 27:XLIV Add species as follows:
3. *syraces* (Godman) *CYMAENES*, [1901]. In Godman and Salvin, Biol. 126.1.
Centrali-American, Lepid. Rhop., 2: 595, pl. 103, figs. 1-3. TL -
"Mexico". HT in BM.
- 28:131 Add to entry:
= *lherminieri* Auct. *Lapsus calami*.
- 28:134 Correct spelling to: *tripunctus*.
- 29 New entry as follows:

Genus LIA Vettius Godman

- In* Godman and Salvin, 1901. Biol. Centrali-American, Lepid. Rhop., 2:
589. Type-species by original designation: *Papilio phyllus* Cramer,
[1777]. Uitl. Kapellen 2(15): 122, pl. 176, figs. B, C.
1. *fantasos* (Stoll) *PAPILIO*, 1780. Pap. Exot., 4: T. 300, figs. E, F. TL - 137.5.
Surinam. Loc. of type unknown.
= *abebalus* (Stoll) *PAPILIO*, [1781]. Uitl. Kapellen 4(31): 145, pl.
365, figs. G, H. TL - "Guinea and Cape of Good Hope". Loc. of
type unknown.
= *eucherus* (Plötz) *HESPERIA*, 1882. Stett. ent. Zeit., 43: 452.
TL - "Surinam". Loc. of type unknown, perhaps in Munich.
- 30:139 Correct reference to Note 87 to: Note 84.
- 30:LIV Add species as follows:
2. *huasteca* (H. A. Freeman), *TIRYNTHIA*, 1966. J. Lepid. Soc., 23 140.1.
(Suppl. 2): 44, pl. 14, figs. 3-6; pl. 10, fig. 5. TL - 7 mi. S. Valles, San
Luis Potosi, Mexico. HT in USNM.
- 31:144 Correct spelling to: *powsheik*. Also misspelled in MONA Check List
(the species name differs from the spelling of the county name).
- 31:147 To conform to the CODE, emend spelling to: *aurantiacus*.
- 32:148 To conform to the CODE, emend spelling to: *minimus*.
- 32:148 Add to entry:
= *singularis* (Plötz [*nec* Herrich-Schäffer, 1865]) *THYMELICUS*,

- 32:LX 1884. Stett. ent. Zeit., 45: 284. TL - ??. Loc. of type unknown.
 Correct spelling in 4th line of entry to: *Adopoea*. Scudder [1875. Proc. Amer. Acad. Arts. Sci., 10: 103] misspelled this name as *Adopaea*.

32:150 Correct Note 73 to read: Note 75.

33:151 Add at end of entry: Many other palearctic synonyms omitted.
 Add to entry:
 a. p. *phyleus* (Drury) *PAPILIO*, [1773]. Etc. 151a.
 b. p. *muertovalle* Scott, 1981. *Papilio* (new series), 1: 10. TL - Furnace Creek, date grove, Death Valley, Inyo Co., California. HT in LACM.

34:154a Add to entry:
 TL restricted by Scott, 1981 [*Papilio* (new series), 1: 12] to "Nr. Victorville, San Bernardino Co., California".

34:154 Add entry as follows:
 c. e. *alinea* Scott, 1981. *Papilio* (new series), 1: 12. TL - Afton, San Bernardino Co., California. HT in LACM. 154c.

34:LXV Correct spelling in 7th line of prefatory material to: Sodovsky.
 Add at end of entry:
 = *Herpina* W. H. Edwards, 1872. In Hayden, Geol. Surv., Montana: 466-467. *Lapsus calami*.

35:158 Add at end of entry: Many palearctic synonyms omitted.

36:158 Add to entry:
 o. c. *oroplata* Scott, 1981. *Papilio* (new series), 1: 8. TL - Spring Creek, Fremont Co., Colorado. HT in LACM. 158o.

36:161, Scott and Stanford ["1981(82)". J. Res. Lepid., 20(1): 18-35] demonstrated that *leonardus* and *pawnee* are conspecific. Thus these entries should be as 161 (delete existing 162) and arranged, starting at end of *leonardus* entry, as follows:
 a. l. *leonardus* Harris, 1862. Etc. 161a.
 b. l. *pawnee* Dodge, 1874. Etc. 161b.
 c. l. *mONTANA* (Skinner) *PAMPHILA*, 1911. Etc. 161c.

36:164 Add following *california* entry:
 = *californica* Skinner, 1920. Ent. News, 31: 175. *Lapsus calami*.

36:165a Correct entry to read: HT in AMNH.

36:165b Durden, 1982 [J. Lepid. Soc., 36(1): 9] has treated this taxon as a separate species.

37:174 Delete *coras*, and replace with *peckius* (W. Kirby). The taxon *coras* refers to a questionable species from South America. See Stanford, 1980 [in Ferris and Brown, Butts. Rocky Mtn. States: 118].

38:175 Add entry as follows:
 d. s. *ministigma* Scott, 1981. *Papilio* (new series), 1: 9. TL - 8 mi. W. Crestone, Saguache Co., Colorado. HT in LACM. 175d.

38:175 The following changes should be made based upon a recent revisionary paper [Austin, 1987. Bull. Allyn Mus., (109): 1-24]:
 The name *genoa* listed under 175a should be placed as a separate subspecies 175e. Austin (*op. cit.*, p. 3) has fixed the TL as Carson River Valley, Douglas Co., Nevada.

38:175 Add five new subspecies as follows:
 f. s. *alkaliensis* Austin, 1987. Bull. Allyn Mus., (109): 4-5. TL - Nev. 8A, 1.3 mi. W. Nev. 34, Granite Mtns., Washoe Co., Nevada. HT in Nevada State Museum. 175f.

- g. s. albamontana* Austin, 1987. Bull. Allyn Mus., (109): 6-7. TL - 175g. Trail Canyon, White Mtns., Esmeralda Co., Nevada. HT in Nevada State Museum.
- h. s. sinemaculata* Austin, 1987. Bull. Allyn Mus., (109): 7-9. TL - 175h. Nev. 140, 5.0 mi. W. Denio Jct., Humboldt Co., Nevada. HT in Nevada State Museum.
- i. s. basinensis* Austin, 1988. Bull. Allyn Mus., (120): 1. Replacement 175i. name for *P. s. pallida* Austin, 1987. Bull. Allyn Mus., (109): 9-10. TL - Nev. 722, 4.0 mi. E. of Reese River, Lander Co., Nevada. HT in Nevada State Museum. Preoccupied by *Pamphila pallida* Skinner, 1911. Ent. News, 22: 412.
- j. s. nigrescens* Austin, 1987. Bull. Allyn Mus., (109): 10-12. TL - 175j. Steptoe Valley, Warm Springs, T21N, R63E, S25 on USGS Monte Neva Hot Springs, Nevada, 7.5° quadrangle, White Pine Co., Nevada. HT in Nevada State Museum.
- 38:179 Add: = *turneri* Freeman, 1944. Ent. News, 55(2): 47. TL - Jesmond, British Columbia. HT should be in Freeman Collection in AMNH.
- 40:184 Freeman, 1950 [Field and Laboratory, 18(2): 78] reported from Texas W. otho *curassavica* (Snellen) *HESPERIA*, 1886. Tijdschr. Ent., 30: 28, pl. 2, figs. 3, 3a. TL - Curaçao, Venezuela. Loc. of type unknown. Although this record has been questioned by some specialists, C. D. Ferris took a male of what appears to be this subspecies in Hidalgo Co., TX on 8.vii.72. J. Y. Miller is revising this group regarding subspeciation.
- 40:186 Correct author font in last entry to: (H. A. Freeman).
- 41:187a Add to note references: [see Notes 22, 134].
Add: = *flaveola* (Mabille) *PAMPHILA*, 1891. Bull. C. R. Soc. ent. Belgique, 35: cixxiv. TL - ??? HT in ???
- 41:189 Add to entry:
3. *mazai* H. A. Freeman, 1969. J. Lepid. Soc., 23 (Suppl. 2): 39, pl. 13, 189.1. figs. 1-2, pl. 15, fig. 8. TL - Mexcala, Guerrero, Mexico. HT in USNM. One paratype from Laredo, Webb Co., Texas.
4. *potosiensis* H. A. Freeman, 1969. J. Lepid. Soc., 23 (Suppl. 2): 40, 189.2. pl. 13, figs. 3-4; pl. 15, fig. 9. TL - 7 mi. S. Valles, San Luis Potosi, Mexico. HT in USNM.
- 42:LXXII The taxonomy in genus *Ochlodes* is currently in a state of flux. Several recent publications have placed *pratincola* (192b) as a separate species, and *nemorum* (193c) has been omitted entirely. Until the taxonomy in this genus is resolved, we make no changes here except to add some recently described names.
- 42:192a Correct reference to: [see Note 137].
TL restricted by Scott, 1981 [Papilio (new series), 1: 11] to Tuolumne gold fields, Tuolumne Co., California.
- 42:192b TL restricted by Scott, 1981 [Papilio (new series), 1: 11] to Broderick, Yolo Co., California.
- 42:192c Add: = *amanda* (Plötz) *HESPERIA*, 1883. Stett. ent. Zeit., 44: 197. TL - ??? HT in ???
- 42:192 Add as follows:
d. *s. santacruzus* Scott, 1981. Papilio (new series), 1: 11. TL - Central 192d. Valley, Santa Cruz I., Santa Cruz Co., California. HT in California Insect Survey, Berkeley, California.

- e. s. orecoastus Scott, 1981. *Papilio* (new series), 1: 11. TL - Cullaby 192e.
Lake, Clatsop Co., Oregon. HT in LACM.
- f. s. bonnevillus Scott, 1981 *Papilio* (new series), 1: 10. TL - Thomas 192f.
Can., Ruby Mtns., Elko Co., Nevada. HT in LACM.
- 42:193a The TL for *milo* (based upon H. K. Morrison's records) must be in error.
O. agricola does not occur in extreme NW Oregon (*fide* J. P. Pelham).
- 42:193c TL fixed by Scott, 1981 [*Papilio* (new series), 1: 11] as Broderick, Yolo
Co., California.
- 42:LXXII Add at end of entry:
W. J. Holland recorded a specimen of *O. venatus faunus* (Turati) from
the United States. This was either a misidentification or an import of a
European specimen.
- 43:197 Rearrange and add as follows:
2. *hobomok* (Harris) *HESPERIA*, 1862. Rept. Ins. Inj. Veg., 3rd ed.:
313. TL - "Massachusetts". Loc. of type not known [see Note 144].
a. h. *hobomok* (Harris) *HESPERIA*, 1862. Etc. 197a.
b. h. *wetona* Scott, 1981. *Papilio* (new series), 1: 9. TL - Sand Gulch, S. 197b.
Greenwood, 7200', Custer Co., Colorado. HT in LACM.
- 44:LXXV Add species as follows:
3. *vitellius* (Fabricius) *HESPERIA*, 1793. Ent. syst., 3(1): 327. TL - 205.1.
"Central American Islands". HT - Zoological Museum of Copenhagen.
Evans [1955. Cat. American Hesp. Brit. Mus. (N.H.), 4: 353] lists a
female of this species from Florida in the British Museum.
= *huebneri* (Plötz) *HESPERIA*, 1883. Stett. ent. Zeit., 44: 199.
TL - "West Indies". HT in ???
= *insularis* (Mabille) *TAGIADES*, 1876. Annales Soc. ent. France,
5(6): 272. TL - St. Thomas, Virgin Islands. HT in BM.
= *portensis* (Mabille) *PAMPHILA*, 1891. C. R. Soc. ent. Belgique., 35: cixxv. TL - Puerto Rico. HT in ???
= *commodus* Kirby, 1903. In Wytsman, Fac. ed., Hübner,
Samml. exot. Schmett.: 110. A replacement name for *vitellius*
Hübner, erroneously considered to be a different species from
vitellius Fabricius.
- 44:206 Delete extra (in (Plötz).
- 45:213 To conform to the CODE, emend spelling to: *conspicuus*.
- 45:215 This taxon almost certainly represents a hybrid.
- 46:216 Stanford [in Ferris and Brown, 1981. Butts Rocky Mtn. States: 101]
elevated *illinois* to subspecific status. Thus entry should show:
a. b. *bimacula* (Grote and Robinson) *HESPERIA*, 1867. Etc. 216a.
b. b. *illinois* (Dodge) *HESPERIA*, 1872. Etc. 216b.
- 46:217 Because of the confusion surrounding the use of the name *ruricola*, one
cannot consider that the Principle of Priority applies here. Therefore
vestris Boisduval should be restored as the specific epithet. For further
discussion, see Brown and McGuire, 1983. [Trans. San Diego Soc. nat.
Hist., 20:69-79]. [c.f. Note 153].
- 46:217b The name *kiowah* applies to the Rocky Mtns. populations as used by
Stanford, 1980 [in Ferris and Brown, Butts. Rocky Mtn. States]. See
also Brown and McGuire, 1983 [Trans. San Diego Soc. nat. Hist., 20: 69-
79]. Durden [1982, J. Lepid., 36(1): 9] treated *osyka* as a valid sub-
species. We suggest that *vestris* be arranged as follows:

- a. v. *vestris* (Boisduval) *HESPERIA*, 1852. Etc. 217a.
 b. v. *metacomet* (Harris) *HESPERIA*, 1862. Etc. 217b.
 c. v. *kiowah* (Reakirt) *HESPERIA*, 1866. Etc. 217c.
 d. v. *harbisoni* J. W. Brown and McGuire, 1983. Trans. San Diego Soc. 217d.
 nat. Hist., 20(3): 58-59. TL - 13.3 km E. Dulzura, N. slope Tecate Peak,
 500 m, San Diego Co., California. HT in San Diego Nat. Hist Museum.
- 47:226b The taxon *margarita* should be placed as f. "margarita" of *A. python* (225), and the *cestus* entry renumbered as 226. See Tilden and Smith [1986. Field Guide Western Butts.: 223].
- 47:227 Delete all references to *ovinia* and elevate *edwardsi* to a full species 247. See Burns, 1983 [Proc. ent. Soc. Washington, 85(2): 335-358].
- 48:234 Based upon rearing studies, J. A. Scott [1976. J. Res. Lepid., 15(2): 92] has determined that *erna* is simply a form of *aenus* (231). Delete this entry and place *erna* as a junior synonym of *aenus*. In its place, insert as follows:
 7. *elissa* Godman, [1900]. In Godman and Salvin, Biol. Centrali-American, Lepid. Rhop., 2: 505, pl. 95, figs. 40-41. TL - Mexico. HT in BM. This species is closely related to *eos* (244), and has been reported from southeastern Arizona. 234.
- 48:238 This taxon may prove to be a subspecies of *tolteca* Scudder, 1872 [4th Ann. Rept. Peabody Acad. Sci. 1871: 76].
- 48:240, 241 Editorial note: These taxa have been considered to be synonymous by some specialists.
- 51:260 To conform to the CODE, emend spelling to: *hecebola*.
- 52 In subfamily *Megathyminae*, delete reference to Note 170.
- 52:LXXXVII The taxonomy in this genus is confused. K. Roever (*in litt.*) has conducted chromosome studies on this genus (unpublished as yet). We feel that the best arrangement of species at this time is as presented by Roever, 1975 [*in Howe, Butts. N. America*]. We will not duplicate this information here. One item should be noted, however, Freeman (*in litt.*), as reported by Ferris [1977. J. Lepid. Soc., 31(1): 74-75] has pointed out that *chisosensis* (271) differs in chromosome number (18) from *neumoegeni* (265) (10). Roever placed *chisosensis* as a subspecies of *neumoegeni*. The CHECK-LIST (PART III of this Memoir) has been rearranged to conform to Roever's treatment with the one exception noted above.
- 52:267 Correct spelling to: *florenciae*.
- 53 In Tribe *Megathymini*, delete reference to Note 170.
- 54:286j To conform to the CODE, emend spelling to: *albasuffusus*.
- 54:289, 290 As suggested by Ferris, 1980 [*in Ferris and Brown, Butts. Rocky Mtn. States: 143*], these taxa should be arranged after the initial *streckeri* entry as follows:
 a. s. *streckeri* (Skinner) *AEGIALE*, 1876. Etc. 289a.
 b. s. *texanus* Barnes and McDunnough, 1912. Etc. 289b.
 c. s. *leussleri* Holland, 1931. Etc. 289c.
- 54:292 To conform to the CODE, emend spelling to: *maculosa*.

Superfamily PAPILIONOIDEA Latreille

- 55: Revise headings as follows:

Superfamily PAPILIONOIDEA Latreille

Hist. Nat. Crust. Ins., 3: 387, "Ann. X", but [Nov., 1802].

Family PAPILIONIDAE Latreille

Hist. Nat. Crust. Ins., 3: 387, "Ann. X", but [Nov., 1802]. See I.C.Z.N. direction 116, 1985. Bull. zool. Nom., 42(pt. 1): 41-42.

Subfamily Parnassiinae Duponchel

Duponchel, 1832-[1835]. In Godart, Hist. nat. Lépid. France, Suppl. 1, (Livraison 22): 380.

- 55:XC Correct lit. cit. 4th line of text to read: VetenskAkad.; 18th line: Das Tierreich.
- 55:239a Correct to read:
 *1. *eversmanni* [Ménétriés] in Siemaschko, [1851]. Russkaya Fauna, fasc. 17, Lepidoptera, tab. 4 as explanation to fig. 5. TL - Kansk (Krasnoyarsk Dist., U.S.S.R.). Male HT in Zool. Inst. Leningrad.
 a. e. thor Hy. Edwards, 1881. Papilio, 1: 4. TL - Yukon River, Alaska 293a. about 100 mi. from mouth. HT in AMNH [see note 178].
 = *pinkensis* Gauthier, 1984. Ent. Zeitschr., 94: 319. Replacement name for *meridionalis* Eisner, 1978. Zool. Meded., 53: 109, pl. 2, figs. 9, 10. TL - Pink Mt., British Columbia. HT in Rijksmus. Nat. Hist. (Leiden); preoccupied by *P. apollo meridionalis* Pagenstecher, 1909. [see Note 179].
 = ab. "kohlsatti" Gunder, 1932. Etc.
- 56:294b This subspecies is now presumed to be extinct.
- 56:294h Delete synonym *shepardi*, and add this taxon as subspecies k.
 K. c. *shepardi* Eisner, 1966. Zool. Meded., 41: 145-146. TL - Wawa-wai Snake R., Washington. HT in Rijksmus. Nat. Hist. (Leiden). This subspecies, which is now extinct at the TL owing to construction of a dam and flooding of the site, bears no resemblance to *altaurus*. See discussion in Ferris, 1976. J. Res. Lepid., 15(2): 65-74. The *shepardi* phenotype does still occur in isolated pockets in areas close to the TL.
- 57:295d Add as follows:
 = *olympianus* Auct. *Lapsus calami*.
- 57:295e Shepard, 1984 (Quaestiones Entomologicae, 20: 35-44) has restricted type localities and designated lectotypes for several Doubleday taxa. Based upon Shepard's research, the type of *smintheus* was collected by Joseph Burke in 1844 [not 1845 as stated by Doubleday] near Jasper, Alberta, and restricted by Shepard to vic. of Rock Lake (53°27'N, 118°16'W). Shepard has selected as the LT of *smintheus* a male specimen presented to the British Museum in 1845 by Doubleday and labeled: "Syntype, Rocky Mts. Pres. by Earl of Derby, 45-136, 33.6, spec. exam C. Eisner". A label has been attached to this specimen with the data: "Lectotype of *Parnassius smintheus*, designated by Jon H. Shepard, 1983". This LT is in the BM.
 Correct the 4th line as follows:
 = *sedakovii* [Ménétriés] in Siemaschko, [1851]. Russkaya Fauna,

fasc. 17, Lepidoptera, tab. 4 as explanation to fig. 2. TL - Irkutsk.
Female LT in Zool. Inst., Leningrad.

- 58:295g Correct the literature citation in the 8th line of entry to read: Lepidopterist.
 57:295 Although stated on p. vi of *Memoir No. 2* that the authors followed "the best available revision" within genera, the published revision by Ferris, 1976 [J. Res. Lepid., 15(1): 1-22] was not followed for the entries under *Parnassius phoebus*. The following reassessments should be made: 295c *magnus* should be placed as a junior synonym of 295e *smintheus*, which now becomes 295d. 295f *xanthus* should be placed as a junior synonym of 295d (new) *smintheus*. 295e (new) should now be shown as subspecies *montanus* Bryk and Eisner, with *maximus* Bryk and Eisner as a junior synonym. The remaining subspecies are renumbered as shown in the Checklist.
- 59: Heading should read:
- Subfamily PAPILIONINAE Latreille
- Hist. Nat. Crust. Ins., 3: 387, "Ann. X", but [Nov., 1802]. General Note: The generic names used in this section are in general agreement with the family revision on a World basis published by Hancock [1983. *Smithersia*, 2: 1-48].
- 59:XCI Correct 2nd line of type species entry to read: *Princeps echelus* Hübner, [1815] (= *Princeps echemon* Hübner, [1813]. Samml. exot. Schmett., 1: pl. [121]).
- 59:296 Correct date of publication of *eurimedes* to: [1782].
- 59:296a Correct entry to read:
 = *#arcas* (Stoll) *PAPILIO*, [1872]. In Cramer, Uitl. Kapellen, 4: 174. TL . . .
- 60:297 For [see Note 000], read: [see Note 196].
- 60:297a At end of entry, insert additional synonym and explanatory note as follows:
 = *septentriae* Fabricius, 1938 [in Bryk, J. Chr. Fabricius Systema Glossatorum. Im Anhang: K. Illiger: Die neuste Gattungs-Einteilung der Schmetterlinge aus den Linneischen Gattungen Papilio und Sphinx. J. Chr. Fabricius: Rechenschaft an das Publicum über seine Classification der Glossaten. Verlag Gustav Feller, Neubrandenburg, Germany, pp. 23-24]. This reference is subsequently abbreviated as: J. C. Fabr. Syst. Gloss. This is a replacement name for *philenor* based upon a foodplant name.
- 60:297b The HT of ab. "inghami" is in AMNH.
- 60:299 Correct lit. cit. for Tribe Leptocircini to read: 1896. Allen's Nats. Libr.; Lepid. (Butterflies) 2: xviii, 307.
- Correct 3rd line of entry to read:
 = *Teinopalpidae* Grote, 1899. Proc. American phil. Soc., 38: 16.
- 60:300 (Cramer) should be shown as: (Cramer)
 Insert additional synonym as follows:
 = *annoneae* Fabricius, 1938. In Bryk, J. C. Fabr. Syst. Gloss., p. 49. A replacement name for *ajax* based upon an erroneous food-plant name.
- 61:302 Correct page ref. for Lucas in 1st line to read 130, not 180.
 Correct *sinon* (Cramer) to: *sinon* (Stoll).

61: Heading should read (below middle of page):

Tribe Papilionini Latreille

Hist. Nat. Crust. Ins., 3: 387, "Ann. X", but [Nov., 1802].

61:XCIV Correct 4th line of text to read:

= *Amaryssus* Dalman, 1816. K. Svenska VetensAkad. Handl., Stockholm, 37: 60. Type species by original designation *Papilio machaon* . . .

Correct spelling in 6th line to: *Aernauta*.

62:303a Insert additional synonym as follows:

= *gracehus* Fabricius, 1938. In Bryk, J. C. Fabr. Syst. Gloss., p. 23. This name was proposed for a female specimen of *asterius* in the collection of Smith Barton [in Barton Collection from Smith?] from Philadelphia, PA.

62:303a 12th line (f. "pseudoamericanus"). Correct page reference to: 291.

At end of 303a entry, add as follows:

b. p. *coloro* W. G. Wright, 1905. Butts. W. Coast, 86: pl. 3, fig. 25. 303b.
TL - Colorado Desert, SE California; fixed by Ferris & J. Emmel [Bull. Allyn Mus., (76): 1-13] as Whitewater Hill, west end of Coachella Valley, Colorado Desert, Riverside Co., California. HT in CAS.

= *rudkini* F. & R. Chermock, 1937. Bull. S. California Acad. Sci., 36: 8. TL - Ibanpah Mtns., California. HT in LACM. See Ferris & J. Emmel (*Op. cit.*) and [see Note 211].

= f. "clarki" F. & R. Chermock, 1937. Bull. S. California Acad. Sci., 36: 8-9. TL - Ibanpah Mtns., California, HT in LACM.

= f. "comstocki" F. & R. Chermock, 1937. Bull. S. California Acad. Sci., 36: 10. TL - Ibanpah Mtns., California, HT in LACM.

62:305 Delete entire entry for *Papilio rudkini*.

62:306 Add the following note at end of entry: Several specialists have suggested that this taxon represents a hybrid between *Papilio machaon* and *P. polyxenes asterius*. Scott [1981. Papilio (new series), 1: 1], however, has asserted that *kahli* is a subspecies of *polyxenes* and has used the combination *P. polyxenes kahli*. While Scott has cited a rearing study, his paper is not a definitive taxonomic revision. Until such a paper is published, the taxonomic placement of *kahli* remains unclear.

62:308 Correct date in 1st line to: 1866.

63:308 Correct spelling in 3rd line to: "hollandii".

63:309 This taxon has been variously treated by authors as a separate species and as a subspecies of *P. bairdii*, the latter most recently by Fisher [in Ferris and Brown, 1981. Butts. of the Rocky Mtn. States, p. 183]. Under the concept that *oregonius* is a yellow expression of *bairdii*, then the entries should be changed as follows:

- | | |
|---|-------|
| a. b. <i>baridii</i> W. H. Edwards, 1869. | 308a. |
| b. b. <i>oregonius</i> W. H. Edwards, 1876. | 308b. |
| c. b. <i>dodi</i> McDunnough, 1939. | 308c. |

Scott [1986. Butts. N.A.] without comment has placed *oregonius* and all

- of the subspecies of *bairdii* as subspecies of *machaon*.
- 63:310a Add at end of entry: Seyer [1977. Mitteilungen ent. Ges. Basel, 27(4): 113] published three names attributed to F. Chermock. These names are also cited in Zoological Record for 1977. Apparently the taxa were intended to be new subspecies, but Chermock never published the names and no designated type specimens can be located in the Chermock material at the AME. Seyer listed only the names and did not provide descriptions. Hence the taxa *kwakwapoochesi*, *prestoni*, and *frechini* are clearly *nomina nuda* with no taxonomic status.
- 63:310b Add at end of entry: The name *avinoffi* may well apply to a yellow expression of *P. kahli* (entry 306).
- 63:311a Delete the entry:
 = *chloro* [sic] W. G. Wright, etc. See new entry 303b under the correct spelling *coloro*.
- 64:312c The TL has been fixed as Devil Canyon, ca. 11 km NNW of San Bernardino Co., California by S. E. Miller [1984. J. Res. Lepid., 23(2): 175].
- 64:312 Add two new subspecies entries as follows:
 h. i. *phyllisae* J. Emmel, 1981. J. Lepid. Soc., 35(4): 297. TL - Butter- 312h. bread Peak and ridge running to the southwest, 4900-5900', S. 30 & 31, T. 29 S., R. 36 E., Kern Co., California. HT in LACM.
 i. i. *panamintensis* J. Emmel, 1981. J. Lepid. Soc., 35(4): 300. TL - 312i. Thorndike Campground, Wildrose Canyon, 7400', Panamint Range, S. 35, T. 19 S., R. 45 E., Inyo Co., California. HT in LACM.
- 65:314 Correct 5th line of entry to read: ST's in AMNH.
 Add at end of entry:
 = *luxuriosus* (Reiff ms.); actually published by Forbes, 1960. Cornell Exp. Sta. Mem. 371: 104. No specific TL indicated. This is a *nomen nudum*.
- 65:316a Correct font to read: (E. M. Sharpe).
- 65:317 Correct lit. cit. to read: Spec. gén. Lépid.
- 66:XCVI Correct lit. cit. for *Euphoeades* to read: (5): 83 [1819].
- 66:320a Correct type font: a. g. *glaucus* . . .
 Insert additional synonyms as follows:
 = *lauri* Fabricius, 1938. In Bryk, J. C. Fabr. Syst. Gloss., p. 22
 A replacement name proposed for *glaucus* based upon a foodplant name.
- Add at end of entry:
 = ab. "ehrmanni" (Ehrmann), 1925. Bull. Brooklyn ent. Soc., 20(2): 84. TL - Hammett Place, Alleghany Co., PA. HT in CM.
- Correct punctuation: = ab. "gerhardi" . . .
- 66:320b Remove synonym *arcticus* (Skinner) and place as a separate subspecies 320d. This butterfly is phenotypically distinct from *P. g. canadensis* and occurs widely in Alaska and the Yukon Territory.
- 66:320c Replace entry as follows:
 c. g. *maynardi* (Gauthier) PAPILIO, 1984. Revista Lepid., 12: 210. Re- 320c.
 placement name for *australis* Maynard, 1891 (*nec* Esper, 1781). Man.
 N. American Butt.: 215. TL - "Florida". HT in MCZ.
- 66:321 Correct page ref. from 158 to 138 here and also in 321a.

- 66:321a Change [see Note 000] to: [see Note 231].
- 66:321b This is a color variant and should be placed as f. "ammoni" under 321a.
- 66:322 To conform to the CODE, emend spelling to: multicaudatus.
- Correct lit. cit. to read: Spec. gén. Lépid.
- 67:323 Add note at end of entry: Upton [1984(85). J. Lepid. Soc., 38(3): 165-170] has asserted that *eurymedon* (Lucas) is predated by an earlier name proposed by Donovan [1805. Nat. Hist. Ins. New Holland, pl. 16]. Donovan's type specimen, which appears to be *eurymedon*, bears a locality label "New Holland", and the specimen is in the C.S.I.R.O., Canberra, Australia. Because Donovan's name has not been used since 1891, a petition is being filed with the I.C.Z.N. (under Art. 79 of the CODE) to suppress the taxon in favor of the well-established *eurymedon* (Lucas).
- 67:324 Correct lit. cit. to read: Spec. gén. Lépid.
- 67:325a Prior to the aberration entries, insert the following synonym:
 = *anethi* Fabricius, 1938. In Bryk, J. C. Fabr. Syst. Gloss., p. 21.
 A replacement name proposed for *troilus* based upon a foodplant name.
- In 10th line, correct author cit. to read: (Dufrane).
- 67:326 Add note: H. A. Freeman [1951. Field and Laboratory, 19(1): 32] reported the occurrence in Texas of the Mexican subspecies *p. leontis* (Rothschild and Jordan) PAPILIO, 1906. Novit. zool., 13: 599. TL - Monterrey, Mexico. HT in BM.
- 67:326.1 Add new entry as follows:
 8. *victorinus* (Doubleday) PAPILIO, 1844. Ann. Mag. nat. Hist., 14: 326.1.
 418. TL - "West coast of Amer." Female HT in BM. This species has been reported from Texas by Adams [1984. J. Lepid. Soc., 37: 318]; Kendall and McGuire [1984. Bull. Allyn Mus., (86): 11].
- 67:XCVII According to Hancock's revision [1982. Smithersia, 2: 1-48], *anchisiades* should be associated with the genus HERACLIDES (XCV) and placed as 319.1a.
- In 2nd line of text, correct spelling to read: *Priamides hippoionus*.
- 68:327a Add at end of entry:
 = *pandonius* (Staudinger) PAPILIO, 1894. Deut. ent. Zeits. [Iris], 7: 104. Unjustified replacement name for *pandion* (C. and R. Felder).

Family PIERIDAE Duponchel

- 68:329 Beginning with 4th line, revise as follows:
- a. m. *menapia* (C. and R. Felder) PIERIS, 1859. Wiener ent. Monats., 3: 271. TL - "Utah" [see Note 240], here restricted to west of Pyramid Lake, Nevada. HT in BM [see Note 241].
 = *ninonia* (Boisduval) PIERIS, 1869. Ann. Soc. ent. Belgique, 12: 38. TL - "le plus orientale de la Californie". HT apparently in USNM [see Note 22].
 = f. "suffusa" Stretch, 1882. Papilio, 2: 109-110. TL - vic. Spokane Falls, Washington. HT in MCZ.
 = ab. "nigricosta" J. A. Comstock, 1918. The Lepid., 2: 13. TL - "high Sierras of Tulare Co., California, 9-10,000 ft. (near Olancha Peak)". HT in LACM.

- b. m. tau (Scudder) *PIERIS*, 1861. Proc. Boston Soc. nat. Hist., 8: 183. 329b.
 TL - Gulf of Georgia, [Washington]. HT in MCZ. [See: Dornfeld, 1980. Butts. of Oregon].
- c. m. melanica Scott, 1981. *Papilio* (new series), 1: 2. TL - 6 mi. W. 329c.
 Willits, Mendocino Co., Calif. HT in LACM. Note: The relation between this taxon and *ninonia* needs to be elucidated.
- 69:C Add at end of genus citations:
 = *APPIUS* F. M. Brown, 1942. Ent. News, 53: 82-83. *Lapsus calami*.
- 70:CI Add at end of genus citations:
 = *PIERIS* Auct., part, not Schrank, 1801. Fauna boica. 2(1): 152, 161.
- 70:332 Correct line 3 and add as noted:
 = *pseudochloridice* (McDunnough) ...
 = *pseudochlorodice* Ferris and Brown, 1980. Butts. Rocky Mtn. States. *Lapsus calami*.
- 70:334 Add at end on entry:
 = *protodin* (W. H. Edwards) *PINIS* [*sic* = *PIERIS*], 1872. In Hayden, F. V., Geol. Surv. Montana, p. 466. *Lapsus calami*.
- 71:CII Add at end of genus citations:
 = *PIERIS* Auct., part, not Schrank, 1801. Fauna boica. 2(1): 152, 161.

71:CII Discussion of generic status of Artogeia

The use of the generic name *Artogeia* in *Memoir No. 2* was based upon a revisionary paper "Artogeia Verity 1947, gen. rev. for *Papilio napi* Linnaeus" published by Otakar Kudrna in 1974 in *Entomologist's Gaz.*, 25: 9-12. Many specialists have questioned the use of this generic name. A year after *Memoir No. 2* was issued, Kudrna published another paper in which he rescinded his earlier action elevating *ARTOGEIA* to full generic status [*in Blab, J. & O. Kudrna, 1982. Hilfsprogramm für Schmetterlinge. Naturschutz aktuell*, 6: 1-135]. He further recommended that *ARTOGEIA* might be used in the subgeneric sense as defined by Klots [1933]. A generic revision of the Pieridae. *Entomologica am. (N.S.)*, 12: 139-242]. Robbins and Henson [1986. J. Lepid. Soc., 40(2): 79-92] have also presented data that suggest the use of *PIERIS* to replace *ARTOGEIA*. On the basis of these actions, it appears preferable to use *PIERIS* as the appropriate generic name. The generic entry should be changed as follows to reflect this action, and the () removed from author names when the associated species was described under the genus *PIERIS*.

Genus CII: *Pieris* Schrank

1801. *Fauna boica. Burchgedrachte Geschichte der in Baiern cinheimischen und zahmen Thiere*, 2(1): 152, 161. Type-species by selection by Latreille [1810. *Consid. gén. Anim. Crust. Arach. Ins.*; 440, 351] as *Papilio brassicae* Linneaus, 1758, *Syst. Nat.* (ed. 10), 1: 467.
 = *Artogeia* Verity, 1947. *Farfalle diurn. d'Italia*, 3: 192. Type-species by original designation *Papilio napi* Linneaus, 1758, *Syst.*

Nat. (ed. 10), 1: 468. [Note 252 no longer applies].

71:336

Discussion of *Pieris napi* in North America

In 1981, Ulf Eitschberger published a paper suggesting that *Pieris napi* does not occur in North America and he proposed several new names [Atalanta, 11: 366-371]. In 1983, he published a major work (in two volumes) devoted to the "Pieris napi-bryoniae complex" in which many new taxa were proposed, including many new North American taxa. This opus: Systematische Untersuchungen am *Pieris napi-bryoniae-Komplex* (s.l.) (Lepidoptera, Pieridae), Herbipoliana 1(1): 1-504; 1(2): 1-601, published by U. Eitschberger & H. Steiniger, has been the subject of two very critical reviews. The first, by A. M. Shapiro [1984. J. Lepid. Soc., 38(4): 324-327] is in the format of a conventional book review. Shapiro raises a number of questions about the validity of the treatment by Eitschberger of the North American taxa. The second review by O. Kudrna and H. Geiger [1985. A critical review of "Systematische Untersuchungen am *Pieris napi-bryoniae-Komplex* (s.l.)" (Lepidoptera: Pieridae) by Ulf Eitschberger, J. Res. Lepid., 24(1): 47-60] is in the form of a technical paper. Numerous errors in Eitschberger's work are detailed and improprieties regarding the I.C.Z.N. CODE are cited. The authors conclude that Eitschberger's study should be placed in the I.C.Z.N. Official Index of Rejected and Invalid Specific Names in Zoology.

On the basis of these two reviews, no changes to reflect Eitschberger's work are being made in this revision to *Memoir No. 2*. The new taxa and their status are included here for the sake of completeness. The numbers after each entry refer to the page in which the name appears in Volume 1 of Eitschberger's study. Only North American taxa are included here.

Nomina Nuda

<i>Pieris marginalis tremblayi</i>	327.
<i>Pieris marginalis shapiroi</i>	330.
<i>Pieris marginalis browni</i>	332.
<i>Pieris virginensis hyatti</i>	358.

Available Names

<i>Pieris angelika</i> [nec Eitschberger, 1981]	340.
<i>Pieris oleracea ekisi</i>	272.
<i>Pieris marginalis reicheli</i>	301.
<i>Pieris marginalis meckyae</i>	322.
<i>Pieris marginalis guppyi</i>	324.

71:336d Correct date in line 1 of entry to read: 1829.

72:336h Insert as 8th line of entry:

= macdunnoughi dos Passos, 1964. Lepid. Soc. Mem. No. 1: 40,
emendation.

73:338 Add after "(1889)" in 4th line: , preoccupied by "immaculata" de Selys-Longchamps, 1857. Ann. Soc. ent. Belge, 1: 5.

- 73:339 Add to text related to *m. monuste* as follows: Based upon the Linnaean description and the figures of Kleemann [1761. *Der Beyträge zur Natur-oder Insecten-Geschichte, etc.*, in *Der monathlich-herausgegebenen Insecten-Belustigung, von August Johann Rosenhof*, IV: 9-68; Pls. I-VIII] and Cramer [1775-1791. *Papillons exotiques de trois parties du monde l'Asie, l'Afrique et l'Amérique etc.*, Amsterdam, I-IV], W. P. Comstock fixed the type locality for *monuste* as Surinam [1943. *American Mus. Novitates*, (1229): 1-2].
- 73:339c Add: The name *cleomes* may apply to a pale migratory form of *phileta*. See: Pease, R. W. Jr., 1962. *Science*, 137: 987-988. Scott [1981. *Papilio* (new Series), 1: 3] has applied the form name "nigra" to the dark form of *phileta*. No type specimen was specified and this is an infrasubspecific name with no standing under the CODE.
- 74:341a Add at end of entry:
- = *ansonoides* (W. H. Edwards) *ANTHOCARIS* [sic], 1987. *In* Hayden, F. V., *Geol. Surv. Montana*, p. 466. *Lapsus calami*. Incorrectly cited in MONA Check List, 1983: 51 as *ausonoides* [sic].
 - = *ausoniedes* (Hy. Edwards) *ANTHOCARIS* [sic], 1877. *Proc. Calif. Acad. Sci.*, 7(12): 168. *Lapsus calami*.
 - = *aussonides* Elrod & Maley, 1096. *Butts. of Montana*, p. 32. *Lapsus calami*.
- 74:342 Correct TL data as follows: TL - Vic. of Rock Lake, nr. Jasper, Alberta (53°27'N, 118°16'W) as restricted by Shepard, 1984 [*Quaectiones Entomologicae*, 20: 35-44]. LT (as selected by Shepard from Lord Derby material) in BM. [see Notes 260 and 269].
- Add at end of entry:
- = *elsa* Beutenmüller, 1898. *Bull. American Mus. nat. Hist.*, 10: 25.
 - TL - Laggan, "British Columbia" [Lake Louise, Alberta]. ST in AMNH. This species was originally described as a variety of *creusa*. Examination of a syntype in the AMNH collection proved it to be a slightly aberrant *creusa*. The additional specimen that Beutenmüller stated was in the Strecker Collection cannot be found in that collection.
 - = *crensa* Hy. Edwards, 1881. *Papilio*, 1: 15. *Lapsus calami*.
- 75:343d Delete entire entry. See: 74:342 (above).
- 75:344 Add at end of entry: Form "rosa" does not apply to the color of the insect, but rather to a form in which the forewing apical dark markings are much reduced.
- 75:CVI In the 1st line of text, correct citation as follows: [Feb. 1833], Coll. Icon. Hist. Chenilles Europ., 12: pl. Papilionides 5; and [June, 1833], 13: p. [35].
- Add at end of genus citations:
- = *Anthocaris* Hemming, 1934. *Gen. Names hol. Butts.*, 1: 132. *Lapsus calami*.
 - = *Anthocaris* dos Passos, 1964. *Lepid. Soc., Mem. No. 1*: 48-49. *Lapsus calami*.
 - = *Anthochris* Cook, C., 1948. *Lepid. Soc. News*, 2(2): 22. *Lapsus calami*.
- 75:345a Add to entry:
- = *angelina* Boisduval, 1869. *Ann. ent. Soc. Belgique*, 12: 40.

- 75:346 Austin [(1986). J. Lepid. Soc., 39(2): 101] has suggested that *pima* and *cethura* are conspecific based upon intermediate populations in the Mojave Desert. Emmel and Emmel [1973. Butts. S. California] used the combination *A. cethura pima*. This situation requires further study.
- 75:347 Further credence is given to the hybrid theory regarding this taxon [see Note 276] by Shields and Mori, "1978(1979)", J. Res. Lepid., 17(1): 53-55.
- 76:348a The HT of ab. "wrighti" is not in the LACM. A note in the extant Riker mount photographed for Plate 11 of Comstock's Butts. California regarding this specimen states: "ret'd to W. S. Wright, San Diego".
- 76:348h Correct spelling of first word in second line to: Alaska.
- 76:CVII The name *Falcapica* Klots is predicated by *Paramidea* Kusnezov. Therefore the genus entry should be replaced as follows:

Genus CVII: *Paramidea* Kusnezov

[1928]. Faune URSS, Ins. Lépid. 1 (livr. 2): 58, footnote. Type-species by original designation *Midea scolymus* (Butler) *ANTHOCHARIS*, [1866]. J. linn. Soc. Lond., Zool., 9: 52. (cf. Hemming, 1967: 340, *fide* Nekrutenko).

= \ddagger *Midea* Herrich-Schäffer, 1867. Corresp.-Bl. zool.-min. Ver. Regensburg, 21: 105, 143. Type-species by monotypy *Papilio genutia* Fabricius, 1793. Ent. Syst., 3: 193; preoccupied (= *Mancipium midea* Hübner, [1809]. Samml. exot. Schmett., 1: pl. [142], figs. 1-4). Preoccupied by *Midea* Bruzelius, [1855]. Beskr. Hydrachn.: 35.
= *Falcapica* Klots, 1930. Bull. Brooklyn ent. Soc., 25: 83. Type-species, as replacement name, *Papilio genutia* Fabricius, 1793. Ent. Syst., 3: 193; preoccupied (= *Mancipium midea* Hübner, [1809]. Samml. exot. Schmett., 1: pl. [142], figs. 1-4).

- 77:349a Add at end of entry:
= *medea* (Leussler) *ANTHOCARIS* [sic], 1938. Ent. News, 49: 77. *Lapsus calami*.

- 77:349b Change author citation as follows:
(dos Passos and Klots) *ANTHOCHARIS (FALCAPICA)*, 1969.

- 78:CVIII Add at end of entry:
= *Eucolias* Berger, 1986. Lambillionea, Suppl. to 86(7-8): 22-23. Type-species by original designation *Papilio palaeno* Linnaeus, 1761, Fauna Suecia: 272 [incorrectly cited by Berger as: "C. *palaeno* L."].

- 78:351a Correct date (3 entries) of Strecker Syn. Cat. from 1876 to: 1878.
79:351a In 3rd line, correct date from 1886 to: 1891.
The HT of ab. "rothkei" is in AMNH.
79:351c The name *eriphyle* may in fact apply to a hybrid form *eurytheme* \times *philodice*. Pending further study, should this be the case, then the name *hageni* W. H. Edwards (1883) is available to denote the western subspecies of *philodice*.
79:352 The page citation for *ariadne* should be: 12-13.
80:353 This entire entry should be reassigned as 355i under *Colias alexandra*. See: Ferris, 1973, J. Lepid. Soc., 27: 57-73; 1988. Bull. Allyn Mus.,

(116): 1-28. The initial portion of the entry entry should now read:

h. a. *harfordii* Hy. Edwards, 1877. Pacific Coast Lepid., (24): 9 [see 355i. Note 291]. TL - Havilah, Kern Co., California. ST's in AMNH [see Notes 292 and 293].

80:354a Delete statement "HT should be in MCZ" and emend as follows:

The types of this taxon have been examined by C. D. Ferris. The male is indeed *occidentalis*. The accompanying female, however, is a specimen of the "alba" form of *Colias eurytheme* Boisduval. A red lectotype label has been affixed to the pin of the male specimen. It reads: "LECTOTYPE/ *Colias philodice/occidentalis* Scudder/Designated by C. D./Ferris 20 Jan., 1986." This label is handwritten in black ink. LT in MCZ.

80:345b Correct the HT statement to: ST's in AMNH.

Add as follows:

= *chryomelas* Hy. Edwards, 1877. Trans. Amer. ent. Soc., 6: 1-68. *Lapsus calami*.

= *chrysomelaena* Auct., 1983. MONA Check List: 52. This is a questionable emendation with regard to the CODE and the original description of the insect.

81:355e Correct date of Cockerell citation to: 1889.

81:355 Add entry 355g as follows:

g. a. *kluanensis* Ferris, 1981. Bull. Allyn Mus., (63): 1-12. TL - Haines 355g. Junction, Yukon Territory. HT in AME.

= *yukonensis* Berger, 1986. Lambillonea, Suppl. to 86(7-8): 45.

This is a *nomen nudum*. The name appears in a key without a description. No type specimen was designated, nor was a type locality designated other than "Yukon (Canada)". It is assumed that Berger's name refers to *kluanensis*, but it may equally refer to *Colias alexandra christina* which occurs at several localities in the Yukon Territory.

h. a. *apache* Ferris, 1988. Bull. Allyn Mus., (116): 8-11. TL - 13 mi. E. 355h. of McNary, 2520 m, Apache Co., Arizona. HT in AME.

Add entry 355i as noted above (80:353).

81:356 Add entry 356c as follows:

c. m. *lemhiensis* Curtis and Ferris, 1985. Bull. Allyn Mus., (91): 1-9. 356c. TL - Near Meadow Creek Lake, 4 mi. W. of Gilmore, 9100 ft., Lemhi Range, Salmon Nat. For., Lemhi Co., Idaho. HT in AME.

82:357c Per the revision of this species by Ferris [1982. Bull. Allyn Mus., (71): 1-19], *glacialis* is a junior synonym of *h. hecla*, and should be so placed under entry 357a.

82:357.1 Add new species as follows:

8. *canadensis* Ferris, 1982. Bull. Allyn Mus., (71): 1-19. TL - Mile 209 357.1. Alaska Highway, British Columbia. HT in AME. This species was initially described as a subspecies of *hecla*, but subsequently elevated to full species status [Ferris, 1988. Bull. Allyn Mus., (116): 1-28].

82:358 This taxon is tentatively retained, but should be renumbered as species 9. As shown by Ferris [1985. Bull. Allyn Mus., (96): 1-51], there is strong evidence that this taxon represents a hybrid *hecla* x *nastes* [see Note 298]. As such, *boothii* would be placed at the end of the revised entry for 357a. Various authors have assumed that the HT was placed in the USNM, but this does not appear to be correct. The specimens illustrated by Curtis seem

to be those in the Hope Museum at Oxford University, England. As also demonstrated by Ferris [1985. Bull. Allyn Mus., (96): 1-51], *thula* is a color form of *boothii*, and the entry for *boothii* should be emended as follows (deleting species entry 359):

= *thula* Hovanitz, 1955. Wasmann J. Biol., 13: 2-4. TL - 70°45'N, 156°30'W, along the Meade R., S. of Point Barrow, Alaska. HT in USNM. This is presumed to be a hybrid *nastes* x *hecla*, although there is limited evidence that *thula* is becoming a stable phenotype in some areas.

= f. "chione" Curtis, 1835. Nat. Hist Append. in Ross., Voy. Search Northwest Pass.: 65. TL - Boothia peninsula, Northwest Territories. Type in Hope Museum, Oxford University, England.

82:360 C. F. Cowan [1970. J. Soc. Biblphy nat. Hist., 5(4): 300] has established the date of publication of the Boisduval name *nastes* as 1834. The title page of the volume in question, however, is clearly imprinted "1832". Based upon a recent extensive revision by Ferris [1985. Bull. Allyn Mus., (96): 1-51], the entry for *C. nastes*, starting at the end of 360a should be rearranged as follows:

= f. "rossii" Guenée, 1864. Ann. Soc. ent. France, 4(4): 199. TL - Boothia peninsula, Northwest Territories. HT apparently in USNM [see Note 22]. This is presumed to be a hybrid *nastes* x *hecla*.

= f. "gueneei" Avinoff, 1935. In Holland and Avinoff, Mem. Carnegie Mus., 12: 13-14, 21-22, 25-27. TL - Southampton I., Northwest Territories. HT in CM. This is presumed to be a hybrid *nastes* x *hecla*.

b. n. *moina* Strecker, 1880. Bull. Brooklyn ent. Soc., 3: 34. TL - "A considerable distance above Fort Churchill" [possibly Manitoba, but probably in the vicinity of Eskimo Point, Northwest Territories]. LT [designated by Ferris, 1988. Bull Allyn Mus., (116): 22-23] in Strecker Collection at FMNH.

= ab. "harperi" (Gunder) EURYMUS, 1932. Canadian Ent., 360c. 64: 278. TL - Fort Churchill, Manitoba. HT in AMNH.

c. n. *alaska* Bang-Haas, 1927. Horae Macrolepid., 1: 41. TL - "Rompert" = Rampart, Alaska. HT in ZMHU [see Note 299].

= *subarctica* (McDunnough) EURYMUS, 1928. Canadian Ent., 60: 270-271. TL - Bernard Harbour, Northwest Territories. HT in CNC.

= ab. "cocandicides" Verity, 1911. Rhop. Palaeearctica: xxxvii; 355. TL - "Territoire de Barren", 67°40'N, 114°30'W [vic. Coppermine R., Northwest Territories]. HT may be in BM.

d. n. *streckeri* Grum-Grschimailo, 1895. Horae Soc. ent. Rossiae, 29: 360d. 290. TL - Laggan, Alberta. LT [designated by Ferris, 1988. Bull Allyn Mus., (116): 1-28] in Strecker Collection at FMNH.

= ab. "obscurata" Verity, 1911. Rhop. Palaeearctica: pl. 61, fig. 6. TL - Lake Louise, Alberta. HT may be in BM.

= ab. "palliflava" (McDunnough) EURYMUS, 1927. Canadian Ent., 59: 154. TL - Mt. McLean nr. Lillooet, British Columbia. HT in CNC.

- 83:361 Emend entry to read as follows:
11. *scudderii* Reakirt, 1865. Proc. ent. Soc. Philadelphia, 4: 217-218. 361.
 "Rocky Mountains, Colorado Territory", fixed in *Memoir No. 2* by Brown as "vic. Empire, Clear Creek Co., Colorado". HT in FMNH. As ascertained by Ferris, the HT is lost. A neotype has been designated by Ferris, [1987. Bull. Allyn Mus., (112): 6-7] as follows: a male specimen collected by E. M. and S. F. Perkins, 19.vii.63 on the east slope of Berthoud Pass along Hwy. 40 at Hoop Creek in the Arapahoe National Forest, 10,800', Clear Creek Co., Colorado. NT in AME.
- = f. "flavotincta" Cockerell, 1901. Psyche, 9: 186. TL not stated, but reference made to specimens from vic. Twin Lakes and the upper Arkansas River Valley, Lake Co., Colorado, figured by W. H. Edwards, Butts. N. America, 1: pl. *Colias* VIII, fig. 5. Ferris [1987. Bull. Allyn Mus., (112): 8-11] has fixed the TL and designated a LT as follows: TL - Lake Co., Colorado. LT in CM (from W. H. Edwards Collection).
- = *ruckesi* Klots, 1937. J. New York. ent. Soc., 45: 324-326. TL - Windsor [sic] Cr. Canyon, W. of Cowles, New Mexico. HT in AMNH. See Ferris (*op. cit.*).
- Note: The original spelling (1865) of this name is *scudderii*. Reakirt apparently emended the name to *scuddereri* [1866. Proc. ent. Soc. Philadelphia, 5: 136].
- 83:362c Delete 362c as a subspecies and enter it as a junior synonym of 362a. See Ferris [1987. Bull. Allyn Mus., (112): 1-25].
- 83:363a Nothing corresponding to the HT can be identified at the USNM and the HT is apparently lost, or perhaps at the BM.
 Regarding *labradorensis*, Ferris [1988. Bull. Allyn Mus., (122): 1-34] has designated a lectotype which is in the MCZ.
- 83:363b Delete this entry as a separate subspecies. Bean never described this insect, nor was a type specimen designated. The journal article cited is not a description. On this basis, *minisni* is a *nomen nudum*. If the name is to be credited, it should be credited to Barnes & Benjamin by virtue of their description of f. "isni". The form name "neri" of *pelidne skinneri* has line priority over f. "isni". As shown by Ferris [1988. Bull. Allyn Mus., (122): 12-19], the trinomial *skinneri* applies to all of the Rocky Mountain populations of *pelidne*. Thus *minisni* may be taken either to be a *nomen nudum*, or credited to Barnes & Benjamin. If the latter, it falls as a junior synonym to *skinneri* Barnes. Verity misspelled the name as *menisme* [1911. Rhop. Palaearctica: xxxiv; 218].
- 83:364a Ferris [1988. Bull. Allyn Mus., (122): 2] has fixed the type locality as "Mouth of the Saskatchewan River, vic Grand Rapids. W. shore of Lake Winnipeg, Manitoba". The HT (type no. 5081) is in the MCZ.
- 83:364b Ferris [1988. Bull. Allyn Mus., (122): 1-34] has recognized only the taxon
 364c *interior* without subspecies. Thus the entries 364b, c should be placed as synonyms of *interior*. The HT of *laurentina* is lost. A neotype has been designated by Ferris [*op. cit.*] as follows: A male from Baddeck, Cape Breton I., Nova Scotia, collected by J. McDunnough on 16 July, 1938. The NT is in the CNC.
- 84:365a Correct the page citation for Synopsis 9 to: 8.
- 84:367 Correct the citation from Ann. Soc. Ent. to: Bull. Soc. ent.

- 84:367 Add the following form names (all from the same reference):
 = (female) f. "marginata" Riddell, 1941. Trans. R. ent. Soc., 91: 453. TL - San Bernardino Mtns., California. HT in CAS.
 = (male) f. "rubrosuffusa" Riddell, 1941. *Op. cit.*: 454. TL - San Bernardino Mtns., California. HT in CAS.
 = (female) ab. "nigrocapitata" Riddell, 1941. *Op. cit.*: 452. TL - Mill Creek Canyon, San Bernardino Mtns., California. HT in CAS.
 = (male) ab. "flavolineata" Riddell, 1941. *Op. cit.*: 454. TL - San Bernardino Mtns., California. HT in CAS.
- 87:CXII Insert additional species under APHRISSA as follows:
 *2. *orbis* (Poey) *CALLIDRYAS*, 1832 Cent. Lepid. Cuba, 1: 5 figs. TL - 376.1. "Cuba". HT in ANSP.
- 88:CXIV In keeping with the change from ARTOGEIA to PIERIS (71:CII - above) and the sense of Klots (*op. cit.* - above), PYRISITIA and ABAEIS are best considered as subgeneric names of EUREMA, thus delete Genus CXV and Genus CXVI. Add at the end of the current entry for Genus CXIV:
 = *Pyrisitia* Butler, 1870. Cist. Ent., 1: 35, 44. Type-species by original designation *Papilio proterpia* Fabricius, 1775. Syst. Ent.: 478. [see Note 330].
 = *Abaeis* Hübner, [1819]. Verz. bekannt. Schmett., (7): 96. Type-species by designation of Butler, 1870, Cist. Ent., 1: 35 *Papilio nicippe* Cramer, [1779], Util. Kapellen, 3: 31.
 = *Xanthidia* Boiduval and Le Conte, 1829. Hist. Lépid. Amerique sept.: 48-49. Type-species by designation of Scudder, 1875, Proc. American Acad. Arts. Sci., 10: 288 *Papilio nicippe* Cramer, [1779], Util. Kapellen, 3: 31.
- Agassiz used the name HEUREMA as an unjustifiable emendation of EUREMA Hübner [Nomencl. zoo. (4to ed.), Index univ.: 181, 1846].
- 88:379 To conform to the CODE, emend spelling to: *boisduvalianum*.
- 88:380 To conform to the CODE, emend spelling to: *mexicanum*.
- 89:CXV Delete heading and the 2 lines immediately beneath.
- 89:381a To conform to the CODE, emend spelling to: *limoneum*.
- 90:387b Correct author entry as follows: (M. Bates) EUREMA, etc.
- 90:CXVI Delete heading and the 6 lines immediately beneath.
- 91:CXVII This now becomes Genus CXV based upon the two deletions above. Add the following to the prefatory material:
 The incorrect spelling NATALIS was used by Doubleday [List. Spec. Ins. Brit. Mus., Appendix: 10, 1836].
- 91: Change the mid-page subfamily entry to read:

Subfamily Dismorphiinae Schatz

1866. In Staudinger and Schatz. Exot. Schmett. 2 (Die Familien und Gattungen der Tagfalter, Systematischen und analytisch bearbeitet): 56, 57.
- 91:CXVIII This now becomes Genus CXVI to accomodate the two earlier deletions.
- 91:390 Change entry to read as follows:
 1. *albania* (Bates) *LEPTALIS*, 1864. Ent. month. Mag., 1: 6. TL - Cen- 390.

tral Guatemala. HT in BM. According to the most recent revision of this genus by Llorente-Bousquets [1983. Folia Entomológica Mexicana 58: 3-206], *albania* is the species that has been recorded from southern Texas.

Family LYCAENIDAE Leach

- 93: Lines 2 and 6 of text at top of page, correct spelling to: Sodovsky.
- 93:393a Correct ab. "neui" and ab. "fulvus" entries to read: HT in AMNH.
- 93:394 To conform to the CODE, emend spelling to: **cuprea**.
Add subspecies as follows:
d. c. *artemisia* Scott, 1981. *Papilio* (new series), 1: 6. TL - The Potholes, 394d. Teton Co., Wyoming. HT in LACM.
- 94:396 Add subspecies as follows:
c. e. *nevadensis* (Austin) *LYCAENA*, 1984. J. Res. Lepid., 23(1): 83- 396c.
85. TL - Jarbidge Canyon between Pine Creek and Gorge Gulch, 6600', T46N, R9E, S33, Elko Co., Nevada. HT in Nevada State Museum.
- 94:398 Based upon the Principle of Priority and the actions of the I.C.Z.N. cited by Brown and Field [1970. J. New York ent. Soc., 78(3): 175-184] and Miller and Brown [1979. Bull. Allyn Mus., (51): 15-17], we feel that *hyllus* is the correct specific epithet. Therefore the actions of Koçak [1983. Priamus, 3(1): 3-5; 1984. 3(3): 95] are deemed invalid.
Correct spelling to: "wormsbacheri".
Add at end of entry:
= ab. "sternitzkyi" (Gunder) *LYCAENA*, 1927. Canadian Ent., 59: 285. TL - Petaluma, California. HT in AMNH.
- 95:399 To conform to CODE, emend spelling to: **rubida**.
- 96:402b To conform to CODE, emend spelling to: **phaedra**.
- 96:403c Delete entry from "TL - . . . (1969)." and replace as follows:
The action taken by F. M. Brown [1969. Trans. American ent. Soc., 95: 172-173] was in error owing to confusion about two similar place names. The Edwards ST's were extant, unbeknown to Brown. Bird and Ferris [1979. Canadian Ent., 111:637-639] corrected this error. The TL was fixed in the publication cited as: "Crow Nest [sic, Crow's Nest] Pass", Alberta, and a LT designated from one of the original W. H. Edwards ST's. LT in CNC.
- 98:411 Kendall and McGuire, [1984. Bull. Allyn Mus., (86): 15-16] have asserted that the species recorded from the United States is *E. toxea* (Godart), and additionally that *toxea* is a species distinct from *minijas* (Hübner) and not a junior synonym thereof as stated in *Memoir No. 2*. Goodson [1947. The Entomologist, 80: 273-276] identified the Texas butterfly as *toxea*. Scott [1986. Butts. North America has retained *minijas*]. This situation needs additional study.
- 100:420 To conform to the CODE, emend spelling to: **academicum**.
- 101:421 To conform to the CODE, emend spelling to: **californicum**.
Austin, 1985[86], J. Lepid. Soc., 39(2): 103 has removed *clygnus* from synonymy and treated it as a valid subspecies. Brown and Opler [1970. Trans. American ent. Soc., 96: 43-46] considered *clygnus* to be a synonym of *californicum*.
- 101:422,a To conform to the CODE, emend spelling to: **sylvinum**.
- 101:422e Add as follows:

- = *putmani* (F. M. Brown et al.) *STRYMON*, 1957. Colorado Butts.: 131. *Lapsus calami*.
- 101:424a Correct spelling in 3rd line of entry to: *wittfeldii*.
- 101:424b Insert after *lorata* entry as follows:
 = *heathii* (Fletcher) *THECLA*, 1903. Trans. Royal Soc. Can., Sect. IV: 211-212; 1904. Canadian Ent., 36(5): 125. TL - Long River Valley nr. Cartwright, Manitoba. HT in USNM.
- 101:424b Correct spelling in last line to: *inorata*.
- 102:424b Add to 2nd line: Lafontaine originally described this taxon as *borealis* (1969), but emended the spelling to *boreale* in 1970 in his redescription of the taxon [J. Lepid. Soc., 24(2): 83-86].
- 102:424 Add subspecies as follows:
 d. *c. albidus* Scott, 1981. *Papilio* (new series), 1: 5. TL - NW Hayden, 424d. Routt Co., Colorado. HT in LACM.
- 102:427b Insert date of publication: 1862.
- 102:430 Add after 4th line of entry:
 = *soepium* dos Passos, 1970 [*nec* Boisduval, 1852]. J. Lepid. Soc., 24(1): 28. *Lapsus calami*.
- 103:430b To conform to the CODE, emend spelling to: *okanaganum*.
- 103:CXXXV Correct Roman numerals to: CXXXIII.
- 103:431 According to R. K. Robbins at the Smithsonian Institution, there are several species that have been lumped under the name *ocrisia*, and it is not clear that the Texas record applies to this species.
- 103:CXXXVI Correct Roman numerals to: CXXXIV.
- 103:CXXXVII Correct Roman numerals to: CXXXV.
- 104:CXXXVIII Correct Roman numerals to: CXXXVI.
- 104:CXXXIX Correct Roman numerals to: CXXXVII.
 R. K Robbins has a paper in press revising *Thereus*, and new assignments will occur at both the genus (to *REKOA* Kaye, 1904) and species levels.
- 104:439 Insert before entry = *juicha* the following:
 = *mytilus* (Godman and Salvin) *THECLA*, 1887. Biol. Centrali-American, Lepid. Rhop., 2: 37. *Lapsus calami*.
- 104:CXL Correct Roman numerals to: CXXXVIII.
- 104:440 According to Lathy [1926. Ann. Mag. nat. Hist., 17: 35-47] *pion* is a junior synonym of *strophius* Godart. This situation bears further study.
- 104:CXLI Correct Roman numerals to: CXXXIX.
- 104:441 Correct spelling in 3rd line of entry to read: *RUSTICUS*.
- 105:CXLII Correct Roman numerals to: CXL.
- 105:443b J. A. Scott [1986. Butts. N. America] treated *perplexa* as a subspecies of *affinis* (447). J. F. Emmel has informed this editor that *perplexa* will be treated as a separate species in the forthcoming book on California butterflies by J. F. Emmel, T. C. Emmel and S. O. Mattoon.
- 105:448 J. A. Scott [1986. Butts. N. America] has treated *viridis* as a junior synonym of *d. dumetorum* (443a). J. F. Emmel has informed this editor that the same treatment will be used in the forthcoming book on California butterflies by J. F. Emmel, T. C. Emmel and S. O. Mattoon.
 Editorial Note: The taxonomy of genus *Callophrys* has been complicated by two publications: Scott and Justice ["1981(82)", J. Res. Lepid., 20(2): 81-85; Scott, 1986, Butts. N. America]. In the first paper, *C. affinis* is treated as a subspecies of *C. dumetorum*. No mention is made of *C. viridis*.

In the second publication, Scott has combined the taxa *affinis*, *apama*, *homoperplexa*, and *perplexa* as subspecies of *C. affinis*. *C. viridis* and *C. dumetorum* are combined as the single species *dumetorum*. The following subspecies are assigned to the taxon *sheridanii*: *sheridanii*, *comstocki*, *lembergi*, *newcomeri*, and a *nomen nudum paradoxa*. This latter is an unpublished manuscript name attributed to R. E. Stanford, and its use in Scott, 1986 (p. 379) does not meet the publication requirements of the CODE.

106:CXLIII Correct Roman numerals to: CXLI.

Correct spelling in 1st line of prefatory material to read: Ehrlich.

106:450 Correct lit. cit. to read: The Entomologist, 79: 186.

106:CXLIV Correct Roman numerals to: CXLII.

106:452 Revise entry as follows:

1. *spinctorum* (Hewitson) *THECLA*, 1867. Ill. diurn. Lepid. Lyc.; 1: 94. TL - "California". HT in BM.

a. s. *spinctorum* (Hewitson) *THECLA*, 1867. Ill. diurn. Lepid. Lyc.; 452a. 1: 94. TL - "California". HT in BM.

= *cuyamaca* W. G. Wright, 1922. Bull. S. California Acad. Sci., 21: 19. TL - Julian, San Diego Co., California. Type in CAS.

b. s. *inus* (W. H. Edwards) *THECLA*, 1871. Trans. American ent. Soc., 3: 270-271. TL - "Colorado", restricted to about 1 mi. E. Kenosha Pass, Park Co., Colorado by F. M. Brown [1970]. Trans. American ent. Soc., 96: 59]. See also: Clench, 1981 [Bull. Allyn Mus., (64): 21].

106: Add entry as follows:

1. *millerorum* (Clench) *CALLOPHRYS*, 1981. Bull. Allyn Mus., (64): 452.1. 21-23. TL - Vic. El Encarnacion, 2400-2450 m, Hidalgo, Mexico. HT in AME. One specimen has been recorded from Otero Co., New Mexico See Johnson, "1985(1986)", J. Lepid. Soc., 39(2): 119-124.

107:457b Correct spelling *TEHCLA* to: *THECLA*.

107:458,a Delete references to *rhodope* (Godman and Salvin). Species is Mexican. Ft. Wingate is located in New Mexico, not Arizona.

107:458 Add two subspecies:

d. s. *chalcosiva* (Clench) *CALLOPHRYS*, 1981. Bull. Allyn Mus., 458d. (64): 11-13. TL - S. Willow Creek, Stansbury Mtns., Tooele Co., Utah. HT in AME.

e. s. *clenchi* K. Johnson, 1988. Insecta Mundi, 2(1): 76-80. Replacement name for *Callophrys (Mitoura) siva rhodope* Clench, 1981. Bull. Allyn Mus., (64): 14-15 [nec Godman and Salvin, 1887. Biol. Centrali-American, Lepid. Rhop., 2: 48], misidentification; preoccupied. TL - Pinery Canyon, 6500', Chiricahua Mtns., Cochise Co., Arizona. HT in CM.

107:460 To conform to the CODE, emend spelling to: *grynea*.

Correct spelling to: *patersonia*.

107:460a Insert after the entry for *damon* (*Stoll*):

= *demon* dos Passos, 1970. J. Lepid. Soc., 24(1): 32. *Lapsus calami*. dos Passos credited this misspelling to "Skinner, 1897", but the name is spelled correctly as *damon* in Skinner's 1897 paper [Canadian Ent., 29: 156]. In this paper, Skinner described *Thecla damon* "n[e]w var. *discoidalis*". See entry 460d.

107

Add new species:

11. *thornei* J. W. Brown, [1983]. J. Res. Lepid., 21(4): 246-249. TL - 461.1.
 Little Cedar Canyon, north slope of San Ysidro Mtns., (Otay Mtn.), east end of Lower Otay Lake, 200 m, 32°37'N, 116°52'W, San Diego Co., California. HT in San Diego Nat. Hist. Museum. [NB: Taxonomically this species is closest to *loki* (459)].

108:CXLV Correct Roman numerals to: CXLIII.

108:462 Delete last two lines of entry and insert as follows:

- a. *x. xami* (Reakirt) THECLA, 1866. Proc. Acad. nat. Sci. Philadelphia, 462a. [1866]: 332. TL - "nr. Vera Cruz, Mexico". Loc. of type unknown.
 b. *x. texami* (Clench) CALLOPHRYS, 1981. Bull. Allyn Mus., (64): 6- 462b.
 7. TL - Corpus Christi [Nueces Co.], Texas. HT in CM.

108:CXLVI Correct Roman numerals to: CXLIV.

108:463 Delete entry number (463) at right margin, and add at end of entry:

- a. *m. mcfarlandi* P. Ehrlich and Clench, 1960. Ent. News, 71: 138. 463a.
 TL - La Cueva Canyon, W. slope Sandia Mtns., Bernalillo Co., New Mexico. HT in AMNH. An additional subspecies from Mexico was recently described.

108:CXLVII Correct Roman numerals to: CXLV.

108:464 The correct name is *Incisalia augustinus* Westwood, 1852; not *I. augustus* Kirby, 1837. Westwood made *Thecla augustus* Kirby a secondary homonym of *Hesperia augustus* Fabricius 1793 when he switched the latter to *Thecla*. Consequently Westwood proposed *augustinus* as a replacement name for *augustus* Kirby. Apparently Comstock and Huntington in their 1959 paper [J. New York ent. Soc., 67: 85] overlooked this situation since *augustus* Kirby was placed under *Incisalia*. The CODE [Art. 59(b)] is specific about the treatment of secondary homonyms: "A junior secondary homonym replaced before 1961 is permanently invalid." Thus [*Thecla*] *augustus* Kirby is permanently invalid, and *augustinus* Westwood applies. The text of entry 464 should be rearranged accordingly so that the primary name is *Incisalia augustinus* (Westwood) with *I. augustus* (Kirby) as the junior secondary homonym. Several recent books have used the incorrect spelling *augusta*.

108:464c Correct spelling to read: *croesioides*.

108:465 The separation of the *Sedum*-feeding *mossii* group from *fotis* was made by Fisher [1980]. In Ferris and Brown, Butts. Rocky Mtn. States]. Subsequent publications have also used this nomenclature, including Tilden and Smith [1986. Field Guide to W. Butts.], Garth and Tilden [1986. California Butts.]. Thus this entry should be rearranged as follows:

2. *fotis* (Strecker) THECLA, 1878. Lepid. Rhop. Het.: 129. TL - "Ari- 465. zona". ST's in FMNH (Strecker Collection).
 2.1. *mossii* (Hy. Edwards) THECLA, 1881. Papilio, 1: 54. TL - Esquimalt, Vancouver Island, British Columbia. HT in AMNH.
 a. *m. mossii* (Hy. Edwards) THECLA, 1881. Papilio, 1: 54. TL - Es- 465.1a. quimalt, Vancouver Island, British Columbia. HT in AMNH.
 b. *m. schryveri* Cross, 1937. Proc. Colorado Mus. nat. Hist., 16: 20. 465.1b. TL - Chimney Gulch, Colorado. HT is not in the Denver Museum of Natural History and is presumed to have been destroyed by dermestids. [see Note 371].
 c. *m. bayensis* (R. M. Brown) CALLOPHRYS, 1969. J. Lepid. Soc., 465.1c.

23(2): 95-96. TL - San Bruno Mtns., San Mateo Co., California. HT in CAS.

d. m. *doudoroffi* dos Passos, 1940. Canadian Ent., 72: 168. TL - Big Sur, Monterey Co., California. HT in AMNH.

e. m. *windi* Clench, 1943. Canadian Ent., 75: 185. TL - Placer Co., California. HT in AMNH.

109:466 To conform to the CODE, emend spelling to: *polia*.

109:466b To conform to the CODE, emend spelling to: *obscura*.

109:467c To conform to the CODE, emend spelling to: *hadra*.

109:468c To conform to the CODE, emend spelling to: *solata*.

109:470a Add at end of entry:

= *plautus* Scudder [*nec* Fabricius], 1987. Bull. Buffalo Soc. nat. Sci., 3: 104. Scudder credited the name, which is a *nomen nudum*, to Abbot [*Papilio plautus*], Draw. Ins. Georgia. Brit. Mus., 6: 55, figs. 173-175; 16, 36, tab. 112. This reference is an unpublished manuscript. Scudder, unfortunately, had the habit of citing unpublished manuscript names.

109:CXLVIII Correct Roman numerals to: CXLVI.

110:CXLIX Correct Roman numerals to: CXLVII.

Clench [1978. J. Lepid. Soc., 32(4): 277-281] rescinded his prior action erecting the genus *EURISTRYMON*. Thus the prefatory material should be changed as follows:

Genus CXLVII: Fixsenia Tutt

1907. Nat. Hist. British Butts., 2: 142. Type-species by original designation *Thecla herzi* Fixsen, 1887. In Romanoff, Mém. Lépid., 3: 279; pl. 13, fig. 4.

= *Leechia* Tutt, 1907, *op. cit.* Type-species by original designation *Thecla thalia* Leech, [1893]. Butts. China Japan Corea (2) (text-part 3): 367; (plate-part 3/4): pl. 30, fig. 15. Invalid junior homonym of *Leechia* South, 1901. Trans. ent. Soc. London, 1901: 400. Replaced by the name *Strymondia* Tutt, [1908]. See below.

= *Strymondia* Tutt, [April, 1908]. Nat. Hist. Brit. Butts., 2: 483. Type-species through Art. 67(i) of CODE (replacement names) *Thecla thalia* Leech, [1893]. Butts. China Japan Corea (2) (text-part 3): 367; (plate part 3/4): pl. 30, fig. 15. New subjective synonym.

= *Euristrymon* Clench, 1961. In P. and A. Ehrlich, How Know Butts.: 212. Type-species by original designation *Papilio favorius* J. E. Smith, in Abbot and Smith, 1797. Nat. Hist. Rarer Lepid. Ins. Georgia, 1: 27.

110:475 Modify entry as follows:

a. p. *polingi* (Barnes and Benjamin) *STRYMON*, 1926. Bull. S. California Acad. Sci., 25: 94. TL - Sunny Glen Ranch, nr. Alpine, Brewster Co., Texas. HT in NMNH.

b. p. *organensis* Ferris, 1980. J. Lepid. Soc., 34(2): 221-222. TL - Finley Canyon, W. Slope Organ Mtns., ca. 6500', Doña Ana Co., New Mexico. HT in AME.

110:CL Correct Roman numerals to: CXLVIII.

- 110:CLI Correct Roman numerals to: CXLIX.
 110:CLII Correct Roman numerals to: CL.
 111:478d To conform to the CODE, emend spelling to: pudicus.
 111:478f To conform to the CODE, emend spelling to: atrofasciatus.
 111:480 To conform to the CODE, emend spelling to: ruofuscus.
 111:482 Correct entry to read: HT in ANSP.
 111:484a To conform to the CODE, emend spelling to: albatus.
 112:487b Occurrence of this subspecies in our area is questionable.
 112:490 According to R. K. Robbins at the Smithsonian Institution, at least two species are presently lumped under *bazochii*, and it is not clear if *bazochii* is the species that has been collected in Texas.
 112:CLIII Correct Roman numerals to: CLI.
 112:CLIV Correct Roman numerals to: CLII.
 113:493a R. K. Robbins at the Smithsonian Institution has suggested that *cypbara* is a separate species rather than a subspecies of *endymion*.
 113 To agree with Eliot's revision [1973. Bull. Brit. Mus. Nat. Hist., 28(6): 443-588], change Tribe entry as follows:

Tribe Polyommatini Swainson

1827. Phil. Mag., (2)1(3): 187 [see Note 1].

- 113:CLV Correct Roman numerals to: CLIII.
 113:495 To conform to the CODE, emend spelling to: exile.
 113:CLVI Correct Roman numerals to: CLIV.
 114:CLVII Correct Roman numerals to: CLV.
 114:CLVIII Correct Roman numerals to: CLVI.
 115 Delete Tribe Everini heading and associated reference.
 115:CLIX Correct Roman numerals to: CLVII.
 115:503a Based upon Wright's illustration of *sissona*, this taxon should be placed as a synonym of *amyntula* (504a).
 115:503b To conform to the CODE, emend spelling to: texanus. Durden [1982. J. Lepid. Soc., 36(1): 13] has suggested that *texanus* is neither a subspecies of *comyntas* nor *amyntula*, but should be associated with *herrii* (504d), which he considers to be a separate species. This situation requires further study.
 115:504a In the entry for ab. "dodgei", delete the superscript 5 immediately after the = sign.
 115:504d Bailowitz [1982. J. Lepid. Soc., 36(4): 308-309] has reaffirmed the placement of *herrii* with *amyntula*.
 116 Delete Tribe Celastrini heading and associated reference.
 116:CLX Correct Roman numerals to: CLVII.
 116:505 In the most recent revision of the genus *CELASTRINA* by Eliot and Kawazoé [1983. Blue Butts. *Lycaenopsis* Group], the taxon *ladon* (Cramer) is placed as a subspecies of the holarctic *argiolus* (Linnaeus). The name *ladon* is used to represent the race of *argiolus* that occurs in the south-eastern portion of North America. On the basis of the Eliot and Kawazoé revision, the following changes should be made in entry 505: The species name is changed to *argiolus*. The second two lines of the prefatory material are replaced as follows:

*1. *argiolus* (Linnaeus) *PAPILIO*, 1758. Syst. Nat. ed. 10, 1: 483. See Eliot and Kawazoé (*op. cit.*), pp. 213-214 for synonymy.

116:505a Studies in progress (unpublished as yet) by several researchers indicate that f. "neglectamajor" may in fact represent a separate species.
Delete references to Note 385 and Note 000.

117:506 The Committee received various letters questioning the validity of the name *ebenina*. This situation was carefully researched by L. D. Miller and C D. Ferris with regard to various W. H. Edwards names and their subsequent treatment by Strecker and Dyar. Miller and Ferris concluded that the name *ebinina* is valid for the edition of the CODE under which Clench worked. The reasons are as follows:

1. W. H. Edwards clearly described "nigra" as a melanic and dimorphic male of the winter form (not variety) of *Lycaena pseudargiolus*; the winter form being "violacea". The arrangement shown by Edwards in Volume 2 of *Butts. N. America* is:

Lycaena pseudargiolus

winter form "violacea"

dimorphic (melanic) male "nigra".

Thus "violacea" in this use by Edwards has only infrasubspecific status, since *pseudargiolus* now has subspecific status, and in addition, Edwards clearly denoted "violacea" as a winter form. This use of "violacea" differs from the use by Edwards of *violacea* [1866. Proc. ent. Soc. Philadelphia, 6: 201]. Consequently "nigra" as used by Edwards has less than infrasubspecific status.

2. In the Strecker Catalogue, entry #136, p. 95 there appears the name "Nig." This has been misinterpreted by some authors as "nig" and not "nig.", which is an abbreviation for "nigra". It was common for 19th Century authors to abbreviate form names. Strecker named "Nig[ra]" based upon the Edwards illustrations of the female of *Lycaena violacea* (fig. 4) in Volume 1 of *Butts. North America*. W. H. Edwards named his male form "nigra" in Volume 2 of *Butts. N. America*. Strecker very clearly designated his "Nig." as an aberration, and thus a status below an infrasubspecific name. The Strecker arrangement is:

Lycaena lucia

ab. [entry] a. "Nig[ra]" (applied to a female aberration)

ab. [entry] b. "Intermedia" (applied to a female aberration)

3. In his 1902 Catalogue, Dyar sunk "nigra" in synonymy and ignored Strecker's "Nig[ra]" entirely.

4. While contemporary authors (on the first reviser basis) have interpreted "var." or "variety" to be synonymous with subspecies, "form" and "aberration" are clearly defined. Edwards stated clearly that his "nigra" applied to a melanic/dimorphic [he used both descriptors] male of a winter form, and Strecker clearly stated that his "Nig[ra]" applied to a female aberration. On this basis, both of these names have no status under the CODE and cannot be applied (nor are they available) to replace *ebenina* Clench.

Delete reference to: Note 000.

Delete Tribe *Scolitantidini* heading and associated reference.

117:CLXI Correct Roman numerals to: CLIX.

118:CLXII Correct Roman numerals to: CLX.

The treatment of the species in this genus generally followed that of Shields

[1975. Bull. Allyn Mus., (28): 1-36] in *Memoir No. 2*. A subsequent revision by Mattoni [1977. J. Res. Lepid., 16(4): 223-242] refuted a portion of Shields's work. Most recent publications on the butterflies of western North America have followed the arrangement proposed by Mattoni. Based upon the comments above, the entries on page 119 should be rearranged as follows:

c. e. enoptes . . .	509c.
d. e. bayensis . . .	509d.
e. e. smithi . . .	509e.
f. e. tildeni . . .	509f.
g. e. langstoni . . .	509g.
h. e. dammersi . . .	509h.
3. mojave . . .	510.
4. rita . . .	
a. r. rita . . .	511a.
b. r. coloradensis . . .	511b.
c. r. emmeli . . .	511c.
d. r. mattonii . . .	511d.
e. r. pallescens . . .	511e.
d. r. elvira . . .	511f.
5. spaldingi (Barnes and McDunnough) <i>PHILOTES</i> , 1917. Contrib. nat. Hist. Lepid. N. America, 3: 216. TL - Provo, Utah. HT in USNM.	
a. s. spaldingi (Barnes and McDunnough) <i>PHILOTES</i> , 1917. Contrib. nat. Hist. Lepid. N. America, 3: 216. TL - Provo, Utah. HT in USNM.	511.1a.
b. s. pinjuna Scott, 1981. <i>Papilio</i> (new series), 1: 6. TL - Shilling's Spring, Conejos Co., Colorado. HT in LACM.	511.1b.

119:CLXIII Correct Roman numerals to: CLXI.

119:CLXIV Correct Roman numerals to: CLXII.

120:514a Add:

= *mildredi* Auct., 1983. MONA Check List: 56. An incorrect emendation that contravenes Art. 31(a) of the CODE.

120:514h In 2nd line of entry, correct to read: HT in AMNH.

120:514i To conform to the CODE, emend spelling to: incognita.

121:514j In 2nd line, correct date of publication to: 1977.

121:515 Correct reference to Note 000 to read: Note 395.

121:CLXV Correct Roman numerals to: CLXIII.

121:516 The species name should be *idas* [see Higgins, 1985. J. Lepid. Soc., 39(2): 145-146]. The initial species entry should be modified as follows:

*1. *idas* (Linnaeus) *PAPILIO*, 1761. Fauna Svecica, (ed. 2): 284, no. 1075. TL - Sweden. ST in Linnaean Collection, London. [See Opinion 269 of the I.C.Z.N., 1954].

122:520 Add the subspecies:

m. i. *helios* (W. H. Edwards) *LYCAENA*, [1871]. Trans. American ent. Soc., 3: 208-209. TL - "From California", restricted to Fawn Lodge, Trinity Co., California by F. M. Brown [1970. Trans. American ent. Soc., 96: 390]. NT in CM, designated by J. C. Downey in Brown 1970 [Trans. American ent. Soc., 96: 392].

122:520i Correct spelling in 2nd line to: Francisco.

123:CLXVI Correct Roman numerals to: CLXIV.

- 123:518b The two entries: f. "caerulescens" and ab. "leussleri" should be removed from this entry and placed under entry 518e.
- 124:CLXVII Correct Roman numerals to: CLXV.
- 124:CLXVIII Correct Roman numerals to: CLXVI.
- 126:521 Add new subspecies at end of entry as follows:
d. s. charlestensis (Austin) *PLEBEJUS*, 1980. J. Lepid. Soc., 521d.
 34(1): 20-22. TL - Lee Canyon, Spring Range, 8250-8800', Clark Co., Nevada. HT in LACM.
- 126:CLXIX Correct Roman numerals to: CLXVII.
- 127:CLXX Correct Roman numerals to: CLXVIII.

Discussion:

In the dos Passos List [1964. Lepid. Soc. Memoir No. 1], only the taxon *aquilo* (Boisduval) was included under the genus *Agriades*. Subsequent publications on both sides of the Atlantic have generally expressed the concept that the taxa *aquilo* and *glandon* (de Prunner) should be used to represent Old World insects. In some instances, *glandon* and *aquilo* have been considered to be conspecific with the former carrying publication priority; in other instances they have been considered to be separate species. In the most recent general publication on European butterflies [Higgins and Riley, 1975. Butts. Britain and Europe: 286-290], they are treated as distinct species. *A. glandon* inhabits interior montane habitats, while *aquilo* is a low-altitude coastal species. Other factors used to separate these two butterflies are differences in adult maculation and larval host plant preference. The North American fauna are perhaps more complex than their European counterparts. There is a leguminosae-feeding, low-altitude butterfly to which the name *franklinii* (Curtis) has been applied. In the Old World, *aquilo* uses *Astragalus alpinus* (Leguminosae) as a larval host. In the Rocky Mountains, the butterfly that W. H. Edwards named *rusticus* is reported to use *Androsace* (Primulaceae) as a larval foodplant, as does also *glandon*. Thus we have two parallel species in the New and Old Worlds. A problem arises, however, with the interior arctic populations of *Agriades*. Reported larval host plants are *Saxifraga* (Saxifragaceae). Possibly a third species is involved, or perhaps out of necessity *rusticus* has adapted to another host. Various legumes do normally exist where these butterflies occur, since *Agriades* is often found flying with *Colias nastes*, which is a legume-feeding species. See Fisher in Ferris and Brown [1980. Butts. Rocky Mtn. States: 208-209] for a discussion of *rusticus*.

Under the assumption that only two species occur in North America, it is suggested that entry 526 be rearranged as follows: Genus description and 526a remain as shown. Entries 526b-f now become 526.1a-e, as noted below:

1. *rusticus* (W. H. Edwards) *LYCAENA*, 1865. Proc. ent. Soc. Philadelphia, 4: 203-204. TL - "Pikes Peak", restricted to vic. Empire, Clear Creek Co., Colorado by F. M. Brown, 1970, Trans. American ent. Soc., 96: 404. LT in FMNH designated by F. M. Brown, *op. cit.* p. 406.
 - a. r. *rusticus* (W. H. Edwards) *LYCAENA*, 1865. Proc. ent. Soc. Philadelphia, 4: 203-204. TL - "Pikes Peak", restricted to vic. Empire, Clear Creek Co., Colorado by F. M. Brown [1970. Trans. American ent. Soc., 96: 404]. LT in FMNH designated by F. M. Brown, *op. cit.* p. 406.
- 526.1a.

b. r. podarce (C. and R. Felder), etc.	526.1b.
c. r. megallo (McDunnough), etc.	526.1c.
d. r. lacustris (T. N. Freeman), etc. [Correct spelling of TL to: Norway House].	526.1d.
e. r. bryanti (Leussler), etc.	526.1e.

Family RIODINIDAE Grote

- 127 Change first line of prefatory material to read:
1895(1827). Mitt. RoemerMus. Hildesheim, (1): 1 [see Note 405].
- 128:CLXXI Correct Roman numerals to: CLXIX.
Correct lit. cit. in 1st line of *Eurygona* entry to read: Spec. gén. Lépid.
- 128 Change 1st line under Subfamily Riodininae entry to read:
1895(1827). Mitt. RoemerMus. Hildesheim, (1): 1 (as Riodinidae) [see Notes 1 and 405].
- 128:CLXXII Correct Roman numerals to: CLXX.
Replace genus description by:
1869. Trans. American ent. Soc., 2: 310. Type-species by designation of I.C.Z.N. [1966. Opinion 755] *Erycinia virginiensis* Guérin-Méneville [1831]. Icon. Ins.: Plate Ins. 81 [see Notes 408 and 409].
= *Calephalis* W. G. Wright, 1908. J. New York ent. Soc., 16(3): 160. *Lapsus calami*.
= *Lephelisca* Barnes and Lindsey, 1922. Ann. ent. Soc. America, 15: 93. Type-species by original designation *Erycinia virginiensis* Guérin-Méneville, [1831]. Icon. Ins.: Plate Ins. 81.
- 128:528 Revise entry to read:
1. *virginensis* (Guérin-Méneville) ERYCINIA, [1831]. Icon. Ins.: Plate 528.
Ins. 81. TL - "Georgia". HT apparently in USNM [see Notes 22, 410, 411 and 412].
= *caenus* Auct. [nec Linnaeus, 1767].
= *pumila*, etc.
- 128:533 To conform to the CODE, emend spelling to: mutica.
Correct footnote reference from 414 to: 415.
- 128:534 Correct footnote reference from 415 to: 414.
- 128:535 In 2nd line of entry, correct "min." to: mi.
- 129:CLXXIII Correct Roman numerals to: CLXXI.
- 129:CLXXIV Correct Roman numerals to: CLXXII.
- 130:CLXXV Correct Roman numerals to: CLXXIII.
- 130:CLXXVI Correct Roman numerals to: CLXXIV.
- 130:541a Add at end of entry:
= *zeia area* F. M. Brown et al., 1957. Colorado Butts.: 120. *Lapsus calami*.
- 131:CLXXVII Correct Roman numerals to: CLXXV.
- 131:544e To conform to the CODE, emend spelling to: mejicana.
- 132:549 Correct spelling in 2nd line to: Arizona.

Family LIBYTHEIDAE Boisduval

There have been two recent revisionary publications devoted to this family. They are included here only for completeness; no changes have been made regarding North

American fauna: Shields, 1983[84]. J. Res. Lepid., 22: 264-266; 1985; Tokurana (Acta Rhopalocerologica), No. 9: 1-58. [In this latter publication, Shields credits the family name Libytheidae to Duponchel (no date)]. See Part I, Note 2 in this Supplement regarding the date and authorship of the family name.

132:CLXXVIII Correct Roman numerals to: CLXXVI.

Family HELICONIIDAE Swainson

133 Delete 3rd and 4th lines at top of page: = Argynitae, etc.

133:CLXXXIX Correct Roman numerals to: CLXXXVII.

133:CLXXX Correct Roman numerals to: CLXXXVIII.

133:CLXXXI Correct Roman numerals to: CLXXXIX.

133:CLXXXII Correct Roman numerals to: CLXXX.

134:CLXXXIII Correct Roman numerals to: CLXXXI.

134:CLXXXIV Correct Roman numerals to: CLXXXII.

Regarding the 2nd line of the genus text, the original spelling was *chari-thonia*. Linnaeus in his Index [1767. Vol. 1, pt. 2, p. 757] corrected the spelling to *charitionia* to conform to its Greek base.

134:559a Correct reference to footnote 443 to read: 442.

135 In the entry for *Blanchardia* (16 lines from top of page), correct the final sentence to read as follows: Preoccupied by *Blanchardia* Castelnau, 1875. Philadelphia Cent. Exhib. of 1876, Off. Rec. Section 7, Essay 2: 47. The generic name as herein used applies to a small Australian fish.

135:560b Correct spelling to: *vazquezae*.

Add:

= *vasquezae*. 1983. MONA Check List: 58. *Lapsus calami*.

135:561a To conform to the CODE, emend spelling to: *petiveranus*.

Family NYMPHALIDAE Swainson

135:CLXXXV Correct Roman numerals to: CLXXXIII.

Add at end of prefatory material:

= *EUPTNETA* dos Passos, 1964. Lepid. Soc. Mem. No. 1: 96.

Lapsus calami. dos Passos credited this misspelling to "[W. H.] Edwards, 1873", but no such form has been found in Edwards's papers.

136:562 Replace the 1st 2 lines at top of page as follows:

= *daunus* Auct.

= *mariamne* Scudder, 1889. Butts. Eastern U.S. and Canada, 1: 519.

= (female) f. "dodgei" Gunder, 1927. Ent. News, 38: 135-136. TL - nr. Scribner, Dodge Co., Nebraska. HT may be in CAS.

136:CLXXXVI Correct Roman numerals to: CLXXXIV.

The name dos Passos is consistently misspelled as dosPassos in this section.

138:568 The taxon *Speyeria nokomis wenona* dos Passos and Grey, 1945 was purposely omitted from *Memoir No. 2* because this insect has been recorded only from Mexico.

138:568a Regarding the type locality comment, specimens have recently been collected

in Ouray Co., Colorado in sight of Mt. Sneffels. Thus the stated TL is most probably correct. An anonymous reviewer has made the following comment: "However, the phenotype of the New Mexico *S. n. nigrocaerulea* is almost identical to these Mt. Sneffels specimens, and is not so similar to the Arizona *S. n. nitocris*." It appears that *nokomis* from New Mexico is variable. Material in the C. D. Ferris collection from Taos and Catron Cos., New Mexico is very close to *nitocris*, and was the basis for Ferris and Fisher [1971. J. Lepid. Soc., 25(1): 44-52] placing *nigrocaerulea* as a synonym of *nitocris*. Thus it appears that *nigrocaerulea* could be equally placed under 568a or 568b.

f. "valesinoidesalba" (Reuss) may apply to 568d (*n. apacheana*); no TL was stated. Note also: = omitted from entry.

ab. "rufescens" (T. Cockerell) should be placed immediately following the entry for *nigrocaerulea*.

138:569 Insert before last line of page:

= *aglaia* (W. H. Edwards) ARGYNNIS, 1864. Proc. ent. Soc. Philadelphia, 2: 505.

139:570,a Correct TL in both entries to: Vic. Agua Fria, Mariposa Co., California. See Masters, 1971 [J. Lepid. Soc., 33(2): 137-138].

139:570d There has been considerable confusion about the correct placement of this taxon. According to L. P. Grey [1989. J. Lepid. Soc., 43(1): 1-10], *gunderi* is a senior synonym of *S. zerene cynna* (571p). Thus entry 570d should be moved to entry 571p. Entries 570e-g now become 570d-f.

140:571d An anonymous reviewer has commented as follows:

"The taxon *S. zerene sordida* (W. G. Wright) does not appear to represent any known population. In fact, Wright himself described *sordida* as an aberration of *bremneri* (Edwards), and never intended to list it at a higher taxonomic level. Moreover, Wright's concept of "*bremneri*" actually represents our modern concept of *S. hydaspe*!"

140:571i Correct date to: 1945.

140:571j See comment under 571d.

140:571o This taxon was described from a mixed series which contained several species and the *zerene* in this series are *platina*. See Grey [1989. J. Lepid. Soc., 43(1): 1-10]. Thus entry 571o as shown should be moved under entry 571n as follows:

= *pfoutsi* (Gunder) ARGYNNIS, 1933. Canadian Ent., 65: 171.
TL - Mt. Loafer, nr. Payson, Utah Co., Utah. HT in AMNH.

140:571p This entry should be renumbered 571o and modified as follows:

o. z. *gunderi* (J. A. Comstock) ARGYNNIS, 1925. Bull. S. California Acad. Sci., 24: 67. TL - N. of Alturas, Modoc Co., California. HT in LACM.

= *cynna* dos Passos and Grey, 1945. American Mus. Novitates, (1927): 4. TL - Humboldt Natl. Forest, Ruby Valley, Elko Co., Nevada. HT in AMNH, [see 139:570d (above)].

140:572 In 1975, Howe [Butts. N. America: 230-235] treated *callippe* and *nevadensis* as separate species with associated subspecies. This arrangement is contrary to current literature and has not been followed in *Memoir No. 2*. A recent revisionary paper by R. A. Arnold [1985. Pan-Pacif. Ent., 61(1): 1-23] in which the number of *callippe* subspecies is reduced to three is not

followed in this revision to *Memoir No. 2*. This paper has proved to be controversial. A cogent rebuttal of Arnold's paper has been published by Hammond, "1985(86)" [J. Res. Lepid., 24(3): 197-208]. See also the treatment by Grey [1989. J. Lepid. Soc., 43(1): 1-10].

141:572g Add at end of entry:

= *sierra* dos Passos and Grey, 1945. American Mus. Novitates, (1927): 5. TL - Gold Lake, Sierra Co., California. HT in AMNH.

141:572h An anonymous reviewer has commented as follows:

"There are two serious problems with the taxon *laura* (W. H. Edwards). First, the designated type locality 'Nevada' is both ambiguous and probably in error. Second, the characteristics of the *laura* lectotype closely match those of *S. c. juba* in the northern Sierra Nevada. Therefore, *laura* is probably best listed as a synonym of *S. c. juba*.

"In conclusion, I would like to suggest that the taxon *inornata* should be raised from synonymy with *juba* to a separate subspecies status, which may require a change in the designated type locality, while the taxa *sierra* and *laura* should be reduced to synonymy with *juba*."

141:572i Delete entire entry. See discussion by Grey [1989. J. Lepid. Soc., 43(1): 1-10].

142:573b-d Emmel and Emmel [1973. Butts. S. California: 29-30] treated these taxa as follows:

Speyeria a. adiaste

a. atossa

a. clemencei

S. a. atossa was last collected in 1959 and it is now thought to be extinct. Garth and Tilden [1986. California Butts.] and Tilden and Smith [1986. Field Guide West. Butts.] have treated *adiaste* as a separate species. It is therefore recommended that 573b-d be separated from *egleis* and made a new entry 573.1; see Checklist.

143:573m Add new subspecies (see Checklist for revised numbering):

j. e. moecki Hammond and Dornfeld, 1983. J. Lepid. Soc., 37(2): 115- 573j.
120. TL - Skookum Meadow, Walker Rim, Klamath Co., Oregon. HT in AMNH.

143:574d,p The problem here is that two putative subspecies have the same type locality. For discussion, see: Ferris, 1983 [J. Res. Lepid., 22: 101-114].

144:574r Insert at end of entry: [see Note 532].

144:574 After 574u, insert additional subspecies:

v. a. *ratonensis* Scott, 1981. Papilio (new series), 1: 4. TL - Raton Me- 574v.
sa, Colfax Co., New Mexico. HT in LACM.

w. a. *elko* Austin, 1983. J. Lepid. Soc., 37(3): 244-248. TL - Owyhee 574w.
Valley, Wildhorse Creek C.G., ca. 10 mi. S. Mountain City, Elko Co.,
Nevada. HT in Nevada State Museum.

x. a. *capitanensis* R. Holland, 1988. Bull. Allyn Mus., (115): 1-9. TL - 574x.
Padilla Point, crest of Capitan Ridge, Capitan Mtns., Lincoln Co., New
Mexico. HT in AMNH.

145:575e See discussion of TL by Grey [1989. J. Lepid. Soc., 43(1): 1-10].

145:575g This subspecies is known from only three specimens collected in 1932.
Despite intensive searching by many collectors, no additional examples have
been found and this subspecies is presumed to be extinct.

145:576b,c The subspecies *jesmondensis* (576c) should be placed as a junior synonym

- of *opus* (576b) as discussed by Grey [1989. J. Lepid. Soc., 43(1): 1-10].
- 146:576f,h The subspecies *arge* (576f) should be placed as a junior synonym of *mormonia* (576h) as discussed by Grey [1989. J. Lepid. Soc., 43(1): 1-10].
- 146:576i The name *clio* has been used generally to describe the unsilvered form of *eurynome*. Silvered and unsilvered forms occur in various subspecies of *mormonia*. Howe [1975. Butts. N. America] treated *clio* as a separate subspecies, despite the fact that mixed pairs are regularly taken *in copulo*.
- 147:CLXXXVII Correct Roman numerals to: CLXXXV.
- 147:577 According to Crosson du Cormier [1977. Alexanor, 10: 31-43], *napaea* does not occur in North America and *alaskensis* should be the species name. This paper has been disputed by many North American specialists, and no change is made herein pending further study.
- 147:577b Add at end of entry:
 = *pales* Lehmann, 1913. In Seitz, Grossschmett. Erde, 5: 423.
- 147:CLXXXVIII Based upon genitalic studies conducted by L. P. Grey [1989. J. Lepid. Soc., 43(1): 1-10]. the genus *PROCLOSSIANA* should be lumped with the renumbered genus *CLOSSIANA* (CLXXXVI) and the heading changed to:
- Genus CLXXXVI: *CLOSSIANA* Reuss
1920. Ent. Mitt., 9: 192 (footnote). Type-species by original designation *PAPILIO selene* [Denis and Schiffermüller] *PAPILIO*, 1775. Ankündigung eines syst. Werkes Schmett. Wienergegend: 321 [see Note 466].
 = *Proclossiana* Reuss, 1926. Deutsche ent. Zeits. [Iris], 51: 168.
 Type-species by original designation *Papilio aphirape* Hübner [1799-1800]. Samml. Europäischer Schmett.: pl. 5, figs. 23-35.
 (= *Papilio eunomia* Esper, [1779]. Die Schmett., Suppl. Band 1: 94).
- [Entry 578 now follows, then entry 579].
- 147:578a Correct date of *ossianus* (Boisduval) to: "1832" [1834].
- 148:578d This butterfly is only a melanic altitudinal form of *dawsoni* (578c) and is therefore omitted from the Checklist. See Ferris and Groothuis "1970(1971)" [J. Res. Lepid., 9(4): 243-248].
- 148: Delete heading CLXXXIX and the first three lines of text immediately beneath.
- 148:579a Regarding the synonym *marilandica*, this name was applied to a presumed univoltine population. Anderson and Simmons, 1979 [J. Lepid. Soc., 33(2): 143-145], have demonstrated that the type locality population is multivoltine.
- 149:580c Correct first portion of entry to read:
 (D. Stallings and Turner) *BOLORIA*, 1946. Canadian Ent., 78: 135-136.
 TL - etc.
- 149:581 Karsholt and Nielsen, 1986 [Ent. scand., 16: 446] have designated a LT for this taxon and fixed the TL as Lappland.
- 149:581b Rearrange the first part of entry as follows:
 b. f. *gibsoni* (Barnes and McDunnough) *BRENTTHIS*, 1926. Bull. S. California Acad. Sci., 25: 92. TL - Barter's I., N. Alaska (presumably

incorrect [see Note 470]). HT stated to be NMNH, but probably in Senckenberg Mus., Frankfurt.

= *alaskensis* (Lehmann) *ARGYNNIS*, 1913. In Seitz, Gross-schmett. Erde, 5: 424. TL - Lat. 69°40'N., Long. 141°W., NE Alaska. HT should be in Senckenberg Mus., Frankfurt [see Note 469].

149:582b Move this entry under 582a as follows:

= *youngi* (Holland) *B[RENTTHIS]*, 1900. Ent. News, 11: 383-384. TL - "mountains between Forty-Mile and Mission Creeks, NE Alaska". HT in CM. See: Ferris, 1984 [Bull. Allyn Mus., (89): 1 7].

149:582b Add new entry as follows:

b. i. *harryi* Ferris, 1984. Bull. Allyn Mus., (89): 2. TL - Near Summit of 582b. Bears Ears Trail, 0.5 mi. NW Mt. Chauvenet, 11, 750', Fremont Co., Wyoming. HT in AME.

149:582 Add new species as follows:

4.1. *acrocema* (Gall and Sperling) *BOLORIA*, 1980. J. Lepid. Soc., 34 582.1. (2): 231. TL - Mt. Uncompahgre, 13 km NW Lake City, Hinsdale Co., Colorado. HT in YPM.

149:583 In entry for *laurenti*, correct page citation to 450 and spelling of TL to Brighton.

150:584c The subspecific name is an invalid homonym, preoccupied by *C. thore borealis* (Staudinger), 1861. The replacement name *uslui* has been proposed by Koçak [1984. Priamus, 3(3):96]. The type specimen and TL remain the same.

150:586 Karsholt and Nielsen, 1986 [Ent. scand., 16: 446] have designated a LT for this taxon and fixed the TL as Lappland, Sweden.

150:586c Mt. Natazhat is in the Yukon Territory. For further discussion of this taxon, see Ferris *et al.*, 1983 [Canadian Ent., 115: 832-833]. Collecting in the Arctic by J. Troubridge in 1987, and Troubridge and K. W. Philip in 1988 has confirmed that *natazhati* is a distinct species. A publication is in preparation. No change is made here, however, pending this publication.

150:586d Delete this entry as a separate subspecies and place *nabokovi* as a junior synonym of *natazhati* (586c). This action is based upon independent examination of the type specimen of *nabokovi* by J. H. Shepard and C. D. Ferris. Correct date of publication for *nabokovi* to 1946.

151:586e Correct the citation to: The Entomologist, 86: 210.

151:588a The TL has been restricted to the vicinity of Rock Lake, Alberta (53°27'N, 118°16'W) by J. H. Shepard [1984. Questiones Entomologicae, 20: 35 44].

151:588 Add additional subspecies:

c. a. *tschukotkensis* Wyatt, 1961. Zeits. Weiner ent. Gesell., 46: 99. 588c. TL - Tschukotka Mtns., NE Siberia. HT in Colin Wyatt Collection, Karlsruhe Museum, West Germany. Editorial Note: This subspecies occurs in the Seward Peninsula and the NW Brooks Range in Alaska. It was originally described as a subspecies of *distincta*. It has yet to be satisfactorily demonstrated that *astarte* and *distincta* are separate species.

151:589f Add at end of entry:

= *sangredecristo* (Scott) *BOLORIA*, 1981. Papilio (new series), 1: 5. TL - Hermit Pass, Sangre de Cristo Mtns., Custer Co., Colorado.

HT in LACM. Based upon the original description, this name applies to an altitudinal female form of *helena*.

152:CXC Correct Roman numerals to: CLXXXVII.

This entry in its entirety does not follow the treatment of the W. H. Edwards names by Brown [1966. Trans. American ent. Soc., 92: 357-468]. In view of present (and perhaps incomplete) knowledge of this genus, Brown's 1966 treatment seems to be a viable approach. Based upon examination of museum specimens from the early 1900s and contemporary material, it appears that *P. minuta* no longer occurs in the United States. The taxon *nymphula* appears to apply to *minuta* (as placed by Brown) and not to *arachne*. While at one time, *m. minuta* occurred in Texas and *m. nymphula* apparently occurred in Arizona, there are no contemporary collections of either butterfly, although a *minuta-arachne* clinal form has been taken by M. E. Toliver in NE New Mexico. See discussion in Ferris and Brown [1980. Butts. Rocky Mtn. States: 329]. A series of contemporaty specimens of *m. nymphula* from northern Mexico is in the AME. Based upon present knowledge, a reasonable arrangement of *Poladryas* is as follows:

1. *minuta* (W. H. Edwards) *MELITAEA*, 1861. Etc.
 - a. *m. minuta* (W. H. Edwards) *MELITAEA*, 1861. Etc. 591a.
 - b. *m. nymphula* (W. H. Edwards) *MELITAEA*, 1884. Etc. 591b.
2. *arachne* (W. H. Edwards) *MELITAEA*, 1869. Etc.
 - a. *a. arachne* (W. H. Edwards) *MELITAEA*, 1869. Etc. 592a.

= *gilensis* (Holland) *MELITAEA*, 1931. Etc. TL - Along the west fork of the Gila River in Catron Co., New Mexico. The HT bears the locality label: "W. F. Gila/7.16 N". The Ft. Gila, S. Arizona locality is incorrect. The HT figured by Holland is a female and not a male as stated. HT in CM. The name *gilensis* actually applies to an aberration in which the VHW dark markings are much reduced.

= ab. "gunderiae" (Holland), etc.
 - b. *a. monache* (J. A. Comstock) *MELITAEA*, 1918. Etc. 592b.

152:CXCI Several changes are necessary in this section based upon a recent revision of the genus by M. J. Smith and J. P. Brock [1988. Bull. Allyn Mus., (118): 1-21].

152:CXCI Correct Roman numerals to: CXXXVIII.

153:595 Break entry at end of 2nd line, delete 595, and insert as follows:

- a. *c. cyneas* (Godman and Salvin) *MELITAEA*, 1878. Proc. zool. Soc. 595a.
London, [1878]: 269. TL - Oaxaca, Mexico. HT in BM.

153:596 Break entry at end of 4th line and insert as follows:

- a. *f. fulvia* (W. H. Edwards) *MELITAEA*, 1879. Canadian Ent., 11: 596a.
117-118. TL - "W. Texas", restricted to Archer Co., Texas by F. M. Brown, 1966, Trans. American ent. Soc., 92: 391.
= ab. "sinefasciata" (R. C. Williams), etc.
- b. *f. coronado* M. Smith and Brock, 1988. Bull. Allyn Mus., (118): 6- 596b.
10. TL - Summit of Redington Pass, 20 km ESE of Tucson, Santa Catalina Mtns., Pima Co., Arizona. HT in LACM.
- c. *f. pariaensis* M. Smith and Brock, 1988. Bull. Allyn Mus., (118): 10- 596c.
12. TL - The Cockscomb Ridge, ca. 61 road km E. of Kanab, Kane Co., Utah. HT in LACM.

153:597a Insert at end of entry for f. "obsoleta" as follows:

Misspelled "obliterata" by Strecker [1878. Syn. Cat., : 125].

153:597c Smith and Brock (*op. cit.*) treat *cerrita* as a blend-zone form.

153:597 Add new subspecies as follows:

e. l. *oregonensis* (Bauer) *CHLOSYNE*, 1975. In Howe, Butts. N. America: 166-167. TL - Not clearly stated; Jackson Co., Oregon implied.

The presumed female type is illustrated in pl. 40, fig. 17 with data: "Mt. Ashland, Loop Road, Jackson Co., Ore." HT in LACM.

154:CXCII Correct Roman numerals to: CXXXIX.

155:CXXXIX Add two new species as follows:

8. *ehrenbergerii* (Geyer) *MORPHEIS*, 1833. Samml. exot. Schmett., 3: 604.1.

pl. 443 (5) (named figure only). TL - Mexico? Loc. type unknown.

= *ehrenbergerii* (Hübner [Kirby]) *ANEMECA*, 1871. Synon. Cat., no. 179.

= *ehrenbergeri* Higgins, 1960. Trans. R. ent. Soc., 12: 404. Higgins (*op. cit.*) placed this species in *CHLOSYNE*.

9. *melitaeoides* (C. and R. Felder) *SYNCHLOE*, 1867. Reise Freg. No- 604.2. vara, 3: 396. TL - "Mexico". HT in BM.

= *melitaeoides* (Godman and Salvin) *SYNCHLOE*, 1882. Biol. Centrali-American, Lepid. Rhop., 1: 181.

= *melitaeoides* Higgins, 1960. Trans. R. ent. Soc., 12: 419. Higgins treated this taxon as a form of *C. marina* (Geyer), 1837. Kendall and McGuire [1984. Bull. Allyn Mus., (86): 28] treated *melitaeoides* as a full species.

155:CXCIII Correct Roman numerals to: CXC.

Charidryas was elevated to generic status by Ferris and Fisher [1977. J. Res. Lepid., 16(3): 133-140]. In his massive revision of the Melitaeinae, Higgins [1960. Trans. R. ent. Soc. London, 112: 381-475], did not recognize this taxon. We feel that the species associated with *Charidryas* in *Memoir No. 2* are sufficiently distinct in maculation to merit recognition. There are excellent biogeographical reasons for maintaining *Charidryas* as a genus. It is north-temperate in origin, whereas *Chlosyne* is tropical. On this basis, *Charidryas* is retained as a distinct genus.

156:606a Add the following:

= *nyctis* (Boisduval) *MELITAEA*, 1869. Ann. Soc. ent. Belgique, 12: 53. *Lapsus calami*.

Correct entry for ab. "milburni" to read: HT in AMNH.

156:608a To be consistent with geographic-race distribution, the entry, ab. "blackmorei" should be placed with 608f.

156:608b The status of *whitneyi* is currently under revision and the reader should check on the revised status as published in the forthcoming book on California butterflies by J. F. Emmel, T. C. Emmel and S. O. Mattoon.

Insert as follows after the entry = *hewesi*:

= ab. "hopfingeri" (Gunder) *MELITAEA*, 1934. Canadian Ent., 66: 129. TL - Aster I. (Columbia R.), Brewster, Okanogan Co., Washington. HT in AMNH.

157:608f The HT of f. "dorothyae" is in LACM.

Editorial Note: There are several problems here in regard to the illustrations in Howe [1975. Butts. N. America] as pointed out by J. P. Pelham and J. P. Donahue (*in litt.*). The ensuing discussion applies to Note 497 in *Memoir No. 2* and plate 40 in Howe. In the latter, figs. 13, 16 are *sterope*

and figs. 9, 12 are "dorothyae". The HT male of "dorothyae" (fig. 12) is from Durkee [Baker Co.], Ore. The AT female (fig. 9) is from Gypsum, Snake R., Baker Co., Ore. Fig. 13 is *sterope* with erroneous data *fide* Bauer; fig. 16 is *sterope* from Fulton Canyon, Sherman Co., Ore. According to the CODE, the taxon "dorothyae" is inappropriately described in Howe. The butterfly itself remains somewhat of an enigma. It occurs in the blend-zone between the ranges of *acastus* (to the south) and *palla sterope* (to the north). It remains to be elucidated whether "dorothyae" is a varietal form of *sterope*, a subspecies of *acastus*, or possibly a hybrid between *acastus* and *sterope*. The emendation to "dorothy" in the 1983 MONA Check List contravenes Art. 31(a) of the CODE.

- 157:611 The original spelling is: *gabbi*.
- 157:612 The status of *damoetas* is currently under revision and the reader should check on the revised status as published in the forthcoming book on California butterflies by J. F. Emmel, T. C. Emmel and S. O. Mattoon.
- 158:613c Correct entry as follows:
c. h. *manchada* (Bauer) *CHLOSYNE*, 1959[1960]. Etc.
- 158:CXCIV Correct Roman numerals to: CXCI.
- 158:614 Add:
= *horni* Rebel, 1906. Verh. zool.-bot. Ges. Wein: 9. TL - ?? Loc.
type not known.
= *draudi* Röber, 1913. In Seitz, Grossschmett. Erde, 5: 453. TL -
Mexico. ST's probably in ZMHU.
- 158:CXCV Correct Roman numerals to: CXCII.
- 159:CXCVI Correct Roman numerals to: CXCIII.
- 159:616a Add at end of entry:
= *ubrica* (Skinner) MELITAEA, 1898. Syn. Cat. N. A. Rhop.: 14.
Lapsus calami.
- 159:CXCVII Correct Roman numerals to: CXCIV.
- 159:617a Add at end of *texana* entry as follows:
Misidentified as *Eresia cincta* Edwards, by Scudder [1875. Bull. Buffalo
Soc. nat. Hist., 2: 239, 268].
- 159:617b Opler and Krizek [1984. Butts. E. Great Plains: 143] have suggested that
seminole may be a distinct species.
- 160:CXCVIII Correct Roman numerals to: CXCV.
- 160:623a To conform to the CODE, emend spelling to: *arcticus*.
Add:
= *artica* dos Passos, 1969. J. Lepid. Soc., 23(2): 120. *Lapsus
calami*.
- 161:623b Insert between *euclea* and *selenis* entries as follows:
= *pulchella* (Boisduval) MELITAEA, 1852. Ann. Soc. entomol.
France, (2)10(2): 306, no. 4. TL - "Californie" [the alleged type
bears no locality label]. This specimen is actually *P. pratensis*, but
Boisduval clearly intended the name *pulchella* as a replacement
name for *Papilio tharos* Drury. For further discussion, see Tilden
1974 [J. Lepid. Soc., 28(4): 353-353].
- 161:623b Insert at end of 2nd line of *selenis* entry: [see Note 507].
- 161:623c To conform to the CODE, emend spelling to: *distinctus*.
The HT is in LACM.
- 161:623d Opler and Krizek [1984. Butts. E. Great Plains: 145-146] elevated this tax-

on to species status based upon the presumption that it is univoltine, while *tharos* represents a multivoltine species. Based upon extensive studies, J. P. Pelham has ascertained that *pascoensis* from the TL region is found in typical *tharos* habitats, and is at least bivoltine. On this basis, it is thus a form or subspecies of *tharos*. Hence it appears incorrect to use the name *pascoensis* to represent the univoltine sibling species of *tharos*.

Add to end of entry as follows:

- = f. "herse" G. C. Hall, 1924. J. New York ent. Soc., 32: 110. TL - Taft, British Columbia. HT in AMNH.
- = f. "nigrescens" G. C. Hall, 1924. J. New York ent. Soc., 32: 110. TL - Taft, British Columbia. HT in AMNH.

- 161:623 Editorial Comment: The *tharos*-complex is still under study by several specialists, and the "correct" nomenclature has yet to be established. Scott [1986. Butts. N. America: 309-311] has used *morpheus* (Fabricius) [previously considered as the spring form of *tharos*] to represent the univoltine species. He has placed *pascoensis* as a synonym, which is a contradiction of Pelham's findings. In south-central Wyoming and along the Arizona-New Mexico border, there are *tharos* populations that will probably prove to be distinct species (C. G. Oliver, *in litt.*). Thus there are perhaps a minimum of four sibling species currently included under the taxon *tharos*. Much additional research is required.

- 161:624 Add:
= ab. "harperi" Gunder, 1932. Canadian Ent., 64: 283-284.
TL - McCreary, Manitoba. HT in AMNH.

- 161:625 Editorial Note: With regard to the use of *pratensis* in preference to *campestris*, it should be noted that in the original Behr publication, the name order was *Melitaea montana* (p. 85), *M. campestris* (p. 86), then *M. pratensis* (p. 86). Strecker in his transfer of these taxa to *Phyciodes*, reversed the order placing *P. pratensis* (no. 233) and *P. montana* (no. 234). If the "First Reviser" principle is adhered to, then the correct specific epithet is *pratensis*. On the other hand, if the Principle of Priority is strictly applied per the current edition of the CODE, then the specific epithet becomes *montanus* (correct spelling to conform to the CODE), since the "50-Year Rule" no longer exists. In neither case does *campestris* have precedence. In the interest of maintaining stability, we retain *pratensis* as the species name.

- 161:625b To conform to the CODE, emend spelling to: *montanus*.
162:626 To conform to the CODE, emend spelling to: *pictus*.
162:626b Based upon the revision of the genus by Higgins [1981. Bull. Brit. Mus. (N.H.), Entomol Ser., 43(3): 91], *canace* is a junior synonym of *pictus pictus*.
162:628 To conform to the CODE, emend spelling to: *pallidus*.
162:629 The name *mylitta* was incorrectly emended to *mylittus* by Higgins [1981. Bull. Brit. Mus. (Nat. Hist.), 43(3)].

- 163:629a Insert after 4th line of entry:
= *mylittae* W. G. Wright, 1883. Papilio, 3: 119. *Lapsus calami*.
163:629b The HT of *arizonensis* is in the Bauer collection.
163:629c Add: [NB: Contemporary occurrence of this subspecies in our region is questionable].
163: Insert new Genus CXCVI: *Tegosa* Higgins as follows:

Genus CXCVI: *Tegosa* Higgins

1981. Bull. Brit. Mus. (Nat. Hist.), 43(3): 121. Type-species by original designation *Acraea claudina* Eschscholtz, 1821 in O. von Kotzebue: Entdeckungs-Reise etc.: 3, Appendix (5): 112, pl. [8], figs. 18a, b.
 *1. *anieta* (Hewitson) *ERESIA*, [1864]. Illus. exot. Butts., 3: [23], pl. [12], figs. 43, 44.

a. a. *luka* Higgins, 1981. Bull. Brit. Mus. (Nat. Hist.), 43(3): 126-127. 629.5a.
 TL - Tamazunchale, San Luis Potose [*sic*, Potosi], Mexico. HT in BM.
 There is a specimen from Texas of this species in the LACM. The sub-specific assignment is based solely upon geography and is tentative.

163:CXCIX, CC Delete and renumber as noted below. Based upon a recent paper that treated genetic distances in the checkerspot butterflies [1985. Brussard *et al.*, J. Kansas ent. Soc., 58(3): 403-412], *Hypodryas* and *Occidryas* should be treated as subgenera of *Euphydryas* Scudder (renumbered as CXCVII), thus superseding the revision by Higgins [1978. Entomol. Gaz., 29: 109-115]. Thus parentheses should be deleted for author names associated with the species originally described as *EUPHYDRYAS*. The heading should be modified as follows:

Genus CXCVII: *Euphydryas* Scudder

1872. 4th Ann. Rept. Peabody Acad. Sci., [1871]: 48. Type-species by original designation *Papilio phaeton* Drury, [1773]. Ill. nat. Hist., 1: index et 42, pl. 21, figs. 3, 4.

= *Hypodryas* Higgins, 1978. Ent. Gazette, 29: 110-112. Type-species by original designation *Papilio maturna* Linnaeus, 1758. Syst. Nat., ed. 10, 1: 480.

= *Occidryas* Higgins, 1978. Ent. Gazette, 29: 110-112. Type-species by original designation *Melitaea anicia* Doubleday and Hewitson, [1848]. Gen. diurn. Lepid., 1: pl. 23, fig. 2 [see Note 520].

Editorial Note: It has been suggested by one author [Scott, "1978(80)". J. Res. Lepid., 17(4): 245-252] that *E. anicia* and *E. chalcedona* are conspecific based upon studies of the male genitalia. This placement has been disputed by various specialists (*in litt.*), since there is a considerable amount of field evidence to the contrary, and therefore the conservative approach is taken here. The two taxa are herein treated as separate species based upon the findings of Ferris ["1987(88)". J. Res. Lepid., 26: 109-115].

163:630 The original spelling is: *gillettii*.

163:631b The TL has been restricted by J. H. Shepard [1984. Quaestiones Entomologicae, 20: 35-44] to the vicinity of Rock Lake, Alberta (53°27'N, 118°16'W).

164:631h Emend type locality entry as follows: Sacramento [*sic*] Mtns. Sacramento was misspelled in the original description.

164:631l Add:

= ab. "venusta" Gunder, 1932. Pan-Pacif. Ent., 8: 127. TL - Cottonwood Can., above Roosevelt Camp, Roosevelt Dam, [Gila Co.], Arizona. HT in AMNH.

- 165:631t Correct spelling "Harbey" to: Harney.
- 165:632 Editorial Note: There has been a long-standing taxonomic problem relative to *chalcedona* and *E. colon* (633). Bauer in Howe [1975. Butts. N. America: 176-181], treated these taxa as separate species. Historically they have been treated by specialists as both conspecific and as separate species. Dornfeld [1980. Butts. of Oregon: 72] considered them conspecific, as did Scott ["1978(80)". J. Res. Lepid., 17(4): 245-252]. Again the conservative approach is adopted and *chalcedona* and *colon* are treated herein as conspecific. In the form that *Memoir No. 2* was published there is a conflict between entries 632d (*E. chalcedona huellemani*) and 633e (*E. colon wallacensis*). Both entities have the same type locality. This would present no problem were these butterflies clearly separable, but they are not. Ehrlich *et al.* [1975. Science, 188: 221-228] have suggested that it is possible to have two genetically distinct populations of *Euphydryas* in close geographic proximity. Field studies in Idaho by N. S. Curtis, C. D. Ferris and others have shown that *E. chalcedona* is extremely variable in facies within a given colony in that area, and there is some degree of annual variation as well. Thus it appears that the names *huellemani* and *wallacensis* apply to the same insect, with the latter having publication priority [see Note 524]. On the basis of the preceding discussion, entries 632 and 633 should be combined into the single species-group *chalcedona* with the following modifications:

- 165:632 Revise order of entries as follows:

a. c. <i>chalcedona</i> (Doubleday), etc.	632a.
b. c. <i>colon</i> (W. H. Edwards), etc.	632b.
c. c. <i>corralensis</i> T. and J. Emmel, etc.	632c.
d. c. <i>dwinellici</i> (Hy. Edwards), etc.	632d.
e. c. <i>kingstonensis</i> T. and J. Emmel, etc.	632e.
f. c. <i>klotsi</i> dos Passos, etc.	632f.
g. c. <i>macglashanii</i> (Rivers), etc.	632g.
h. c. <i>nevadensis</i> Bauer, etc.	632h.
i. c. <i>olancha</i> (W. G. Wright), etc.	632i.
j. c. <i>paradoxa</i> McDunnough, etc.	632j.
k. c. <i>perdiccas</i> (W. H. Edwards), etc.	632k.
l. c. <i>quino</i> (Behr), etc.	632l.
m. c. <i>sierra</i> (W. G. Wright), etc.	632m.
n. c. <i>sperryi</i> F. and R. Chermock, etc.	632n.
o. c. <i>wallacensis</i> Gunder, etc.	632o.

- 165:632a Insert after entry for ab. "suprafusca" as follows:

= ab. "suprafusa" dos Passos, 1964. Lepid. Soc. Mem. No. 1:
86. *Lapsus calami*.

- 165:632o This entry should now read as follows:

o. c. *wallacensis* Gunder, 1928. Pan-Pacif. Ent., 5: 50. TL - Wallace, 632o.

Shoshone Co., Idaho. HT in AMNH.

= *huellemani* dos Passos, 1964. Lepid. Soc. Mem. No. 1: 86;
change of status. TL - Wallace, Idaho. HT in LACM [see Note
524].

= ab. "huellemani" J. A. Comstock, 1926. Bull S. California
Acad. Sci., 25: 32. TL - Wallace, Idaho. HT in LACM.

- 166:633 This entire entry should be placed under 632 as noted above.

- 167:633b Change last sentence to read (now 632h as revised):
 Bauer did not designate a type specimen. For discussion [see Note 529].
 LT in Bauer collection.
- 168:634h Reference should be to Note 531, not 532.
- 169:634q Insert after 1st line as follows:
 = *nubigera* (Carpenter) *MELITAEA*, 1877. Field and For., 3: 48.
Lapsus calami.
- 169:634 Insert two additional entries as follows:
 u. e. *luestherae* Murphy and Ehrlich, 1980. J. Lepid. Soc., 34(3): 317. 634u.
 TL - Del Puerto Canyon, 22 mi. W. Patterson, Stanislaus Co., California. HT in AMNH.
- v. e. *koreti* Murphy and Ehrlich, "1983(84)". J. Res. Lepid., 22(4): 257- 634v.
 259. TL - Ridge S. Bald Mtn., White Pine Co., Nevada. HT in AMNH.
 [NB: Phenotypically *luestherae* is virtually identical to *editha*, and *koreti* to *lehmani*. The basis for naming these subspecies is habitat isolation and presumed lack of gene exchange].
- 169 Delete heading CCI and the two lines of text immediately beneath.
- 169:635a Correct entry to read: = ab. "streckeri", etc.
- 170:CCII Correct Roman numerals to: CXCVIII.
- Editorial Note: The partial revision of genus *POLYGONIA* by Scott [1984. J. Res. Lepid., 23(3): 197-210] is considered premature, and it is incomplete with respect to distribution data. Scott's taxonomic decisions are based upon slight differences in the male genitalia of the specimens examined, and upon geographic distributions. Various taxa have been lumped under *progne* (Cramer), based upon presumed allopatry, and various other taxonomic changes have been proposed. The taxon *nigrozephyrus* Scott (in the combination *P. progne nigrozephyrus*) has been erected primarily on the premise that *P. progne progne* and *P. progne nigrozephyrus* are allopatric. It is quite possible, however, that *nigrozephyrus* represents simply an occasional dark form of *zephyrus* (not *progne*) that seems to occur more frequently in females than in males. This dark phenotype ranges at least from the Jemez Mtns. in New Mexico into northern Idaho, and a single female was taken by the Editor along with normal males of *zephyrus* in 1985 on a prairie in Albany Co., Wyoming where nothing remotely resembling *progne* is found [*cf.* "limited to mountainous Colorado, Utah, southern Wyoming, and probably northern New Mexico" as stated by Scott]. Normal *progne* and *zephyrus* are sympatric and synchronic in the Pine Ridge (Sioux Co.) of Nebraska, and in Devil's Tower Nat. Mon. in Crook Co., Wyoming. Dark *zephyrus* specimens have been taken along with typical *progne* in the Pine Ridge. The conservative approach is adopted herein, and no major taxonomic changes are recommended pending further study. *P. p. nigrozephyrus* is tentatively listed, however, as a subspecies of *progne*.
- 170:636 Correct spelling to: *interrogationis*.
 Add at end of entry as follows:
 = *interrigationis* Brown *et al.*, 1957. Colorado Butts.: 91. *Lapsus calami*.
 = *tiliae* (Fabricius) *CYNTHIA*, 1938. In Bryk, J. C. Fabr. Syst. Gloss.: 90. A proposed replacement name based upon one of the larval foodplants.

170:638b Correct spelling to: *neomarsyas*.

This taxon is simply a varietal form of *satyrus* and should be treated as a junior synonym thereof and placed under 638a, renumbered as 638. All forms have been reared from a single brood (*fide* J. P. Pelham).

171:640 *P. hylas* has been treated by many contemporary authors as a subspecies of *P. faunus*. Thus revise entry 640 to 639e.

171:641 Delete as separate species and place as synonym of *faunus rusticus* (639c), *fide* J. F. Emmel.

171:643 Correct entry to read: (Grote and Robinson).

171:645 Correct spelling to: *c-argenteum*.

Based upon the editorial note above, the following changes should be made:

a. *p. progne* (Cramer), etc.

b. *p. nigrozephyrus* Scott, 1984. *J. Res. Lepid.*, 23(3): 201-202. TL - 645b.

Lump Gulch nr. Rollinsville, 8500', Gilpin Co., Colorado. HT in Univ. of Colorado Mus.

172:CCIII Correct Roman numerals to: CXXVIX.

172:646 To conform to the CODE, emend spelling to: *vaulbum*.

Correct author entry to: (Denis and Schiffermüller).

Add to entry as follows:

= *watsoni* (G. C. Hall) *AGLAIS*, 1924. *J. New York ent. Soc.*, 32: 109. TL - Sicamous, British Columbia. HT in AMNH.

172:648a Add at end of entry:

= *betulae* (Fabricius) *VANESSA*, 1938. *In* Bryk, J. C. Fabr. Syst. Gloss.: 111. A proposed replacement name based upon a larval foodplant.

173:CCIV Correct Roman numerals to: CC.

173:649b Add at end of entry:

= *melberti* (W. H. Edwards) *VANESSA*, 1872. *In* Hayden, F.V., Geol. Surv. Montana: 467. *Lapsus calami*.

173:649c Remove 649c as separate subspecies and place as junior synonym of *m. milberti* (649b). There is no appreciable difference between *milberti* and *furcillata*. See Ferris in Ferris and Brown [1980]. Butts. Rocky Mtn. States: 337].

173:CCV Correct Roman numerals to: CCI.

173:650 Insert before *huntera* entry as follows:

= *gnaphalii* (Fabricius) *CYNTHIA*, 1938. *In* Bryk, J. C. Fabr. Gloss.: 102. A proposed replacement name based upon a larval foodplant.

174:651 Insert after second line of entry as follows:

= *carduelis* Field [*nec* Seba, 1765], 1940. *Bull. Univ. Kansas*, 41(22): 342.

= ab. "elymi" Rambur, 1829. *Ann. Soc. Obs.*, 2: pl. 5, figs. 1-2. TL - France?. HT in ?? [NB: See Field, 1971. Smithsonian Contrib. Zool., 84 for a detailed discussion of the various aberrational and form names in *VANESSA*].

174:653a Insert after 4th line of entry as follows:

= *atalanta* Auct. [*nec* Linnaeus, 1758].

= *italica* Field [*nec* Stichel, 1907], 1938[1940]. *Bull. Univ. Kansas* 39(10): 264.

174:CCVI Correct Roman numerals to: CCII.

645a.

645b.

- 175:CCVII Correct Roman numerals to: CCIII.
 Correct spelling in prefatory entry to: *Euralia*.
- 175:CCVIII Correct Roman numerals to: CCIV.
- 175:656 The taxa "rubrosuffusa" and "rosa" were described as forms and not aberrations as stated.
 Add at end of entry:
 = ab. "weidenhameri" Polacek, 1925. Bull. Maryland Acad. Sci., 6(3). TL - Maryland. Loc. type unknown.
- 176:657,658 Editorial Note: Two revisionary papers have been published that affect the taxonomy of these two entries: Hafernik, J. E., Jr., 1982 [Phenetics and ecology of hybridization in buckeye butterflies (Lepidoptera: Nymphalidae), Univ. of California Pubs. in Entomology, 96: 1-109]; Turner, T. W. and J. R. Parnell, 1985 [The identification of two species of *Junonia* Hübner (Lepidoptera: Nymphalidae): *J. evarete* and *J. genoveva* in Jamaica, J. Res. Lepid., 24(2): 142-153]. Based upon these studies, entries 657 and 658 should be revised as shown.
- 176:657 *2. *genoveva* (Stoll) *PAPILIO*, [1782]. In Cramer, Uitl. Kapellen, 4: pl. 290, figs. E, F. TL - not stated, possibly Surinam. Loc. of type unknown, possibly either BM or Amsterdam.
- a. g. *zonalis* C. and R. Felder, [1864-1867]. Reise Freg. Novara, 3: 399. 657a.
 TL - "Columbia, Cuba, Puerto Rico". ST's in BM.
 = *negra* Auct. [nec C. and R. Felder, 1867].
- = *nigrosuffusa* Barnes and McDunnough, 1916. Canadian Ent., 48: 222. TL - "Palmerlee, Babaquivera [sic] Mtns., Huachuca Mtns., Arizona". ST's in USNM. Hafernik placed this taxon as a subspecies of *evarete*. T. W. Turner (*in litt.*) has suggested that *nigrosuffusa* may represent a hybrid between *coenia* and *genoveva* rather than a form or subspecies. Dark-suffused examples intermediate between the two species are regularly collected in SE Arizona and SW New Mexico. For this reason, pending further study, *nigrosuffusa* is placed as a synonym of *genoveva*. Turner has also indicated that the dark insect that has been taken on Padre Island, Texas is a dark *coenia* based upon comparative larval studies and chromosome data. *J. evarete* occurs along the Florida coast, and in Texas as a migrant from Mexico.
- 176:658 *3. *evarete* (Cramer) *PAPILIO*, [1779]. Uitl. Kapellen, 3: 18, pl. 203, 658.
 figs. C, D. TL - "Surinam". Type may be lost [BM, Amsterdam?].
 = ‡ *lavinia* (Cramer) *PAPILIO*, [1775]. Uitl. Kapellen, 1: 32, pl. 21, figs. C, D. TL - "Surinam". Loc. of type unknown [see Note 7]. Preoccupied by *Papilio lavinia* Fabricius, 1775. Syst. Ent.: 450 [see Note 565].
- 176:CCIX Correct Roman numerals to: CCV.
- 176:659 Editorial Note: Silberglied, *et al.* [1979. Psyche, 86: 219-260] revised this genus and recognized no subspecies. We do not accept this approach. Correct the omitted line as follows: TL - "Surinam". Loc. Type - not known and probably lost.
- 176:659a Correct entry as follows:
 ... Munroe, 1942. American Mus. Novitates, (1179): 2-3. TL - San Carlos 659a.
 Estate, Guantanamo, Oriente, Cuba. HT in AMNH.
- 176:660 Add: [NB: Silberglied, *et al.*, 1979, *op. cit.*, have restricted the TL to

"La Habana, Cuba"].

- 176:CCX Correct Roman numerals to: CCVI.
 177:663a Correct footnote reference from 557 to: 571.
 177:CCXI Correct Roman numerals to: CCVII.

Editorial Note: The basis for using *Basilarchia* relates to differences in larval and pupal morphology between the North American fauna and the Old World *LIMENITIS* Fabricius. On the other hand, A. P. Platt (who has been studying this group for some years) in a 1988 letter to the Committee chairman, has stated that he favors the use of *LIMENITIS* with *BASILARCHIA* as a subgenus. At one extreme, true *LIMENITIS* could be considered as a monotypic genus composed of *L. populi* (Linnaeus 1758); at the other extreme, *ADELPHA*, *BASILARCHIA*, and *LADOGA* could be considered to be subgenera under *LIMENITIS*. *BASILARCHIA* is retained here pending further study.

- 180:665 Add entry as follows:
 f. w. siennafascia (Austin and Mullins) *LIMENITIS*, (1984). J. Res. 665f.
 Lepid., 22: 225-227. TL - Chiricahua Mtns., nr. Barfoot Park, Cochise Co., Arizona. HT in AME.
 180:CCXII Correct Roman numerals to: CCVIII.
 180: A major revision of genus *EPIPHILE* has been published by Jenkins [1986. Bull. Allyn Mus., (101): 1-70]. Thus insert above Genus CCXIII: *Myscelia* Doubleday as follows:

Genus CCVIX: *Epiphile* Doubleday

1844. List Spec. Lepid. Ins. British Mus., 1: 90. Type-species by selection by Scudder, 1875. Proc. American Acad. Arts Sci., 10: 165
Temenis orea Hübner, 1823. Samml. exot. Schmett., 2: pl. 30.
 = *Epiphile* Boisduval, 1870. Consid. Lépid. Guatemala: 40. Type-species by selection by Hemming, 1943. Proc. R. ent. Soc. Lond. (B), 12: 27 *Epiphile epicaeste* Hewitson, 1857. Ill. exot. Butts., 2: 47, pl. 24, figs. 2, 4.
 1. *adrasta* Hewitson, 1861. Exot. Butts. *Epiphile* 2: [50], part [40], pl. [25], figs. 9-11. TL - "Mexico". ST's in BM.
 a. a. *adrasta* Hewitson, 1861. Exot. Butts. *Epiphile* 2: [50], part [40], 668.5a. pl. [25], figs. 9-11. TL - "Mexico". STs in BM.

- 180:CCXIII Correct Roman numerals to: CCX.

A major revision of the genus *MYSCELIA* has been published by Jenkins [1984. Bull. Allyn Mus., (87): 1-64]. Starting with the last line on p. 180 to the end of the genus entry, the text should be revised as follows:

1. *ethusa* (Doyère) *CYBDELIS*, 1840. In Cuvier, *Règne Anim.* (Disciple's edn.), Livraison 101 (Ins. 10): pl. 138, figs. 3 & 3b. TL - Unknown. ST's (?) in USNM.
 a. e. *ethusa* (Doyère) *CYBDELIS*, 1840. In Cuvier, *Règne Anim.* (Disciple's edn.), Livraison 101 (Ins. 10): pl. 138, figs. 3 & 3b. TL - Unknown. ST's (?) in USNM.
 = *ethusa* (Boisduval) *CYBDELIS*, 1840. In Cuvier, *Règne Anim.* (Disciple's edn.), Livraison 101 (Ins. 10): pl. 138, fig. 3. TL - "Mexico". "HT" male in USNM does not conform to the CODE (Art. 69) and the name is invalid. Note: This has previously been

cited by authors as "Boisduval, 1836". In his discussion of the Disciple's Edition of Cuvier's *Règne Anim.* [1976. J. Soc. Biblphy nat. Hist., 8(1): 57], Cowan has stated: "Doyère introduced about seven new names in his last livraison. All were Lepidoptera label names from Boisduval's celebrated collection, and the latter has been cited as their author. But Boisduval was never a disciple, and the responsibility for publication must rest with Doyère."

= *cyanecula* (C. and R. Felder) *FAUNIA*, [1864-1867]. Reise Freg. Novara, 3: 408, 618; pl. 53, fig. 5. TL - Mexico. Loc. of HT unknown; not in BM.

= *rogenhoferi* R. Felder, 1869. Verh. zool.-bot. Ges. Wien, (22): 472. TL - Huahuapan (= Huajuapan), Oaxaca, Mexico. ST's may be in Vienna.

*2. *cyananthe* (C. and R. Felder) *FAUNIA*, [1864-1867]. Reise Freg. Novara, 3: 408-409. TL - Mexico. HT in BM.

a. c. *skinneri* Mengel, 1894. Ent. News, 5: 96. TL - Bayemena, Sinaloa, 670a. Mexico. ST's in Reading [Pennsylvania] Public Museum and Strecker Coll. at FMNH.

181:CCXIV Correct Roman numerals to: CCXI.

182:674 Insert after 2nd line at top of page:

= *caerulea* dos Passos, 1964. Lepid. Soc. Mem. No. 1: 73. *Lapsus calami*.

182:CCXV Correct Roman numerals to: CCXII.

Add as follows:

2. *tithia* (Geyer) *SIRONIA*, 1823. In Hübner, Zutr. z. Samml. exot. Schmett., 2: 31. TL - "South Brazil". HT presumed lost. 675.1.

182:CCXVI Correct Roman numerals to: CCXIII.

Add species as noted:

3. *anna* (Guérin-Méneville) *CALLICORE*, 1844. Icon. Règne Anim.: 480. 677.1. TL - Mexico? HT presumed lost.

182:CCXVII Correct Roman numerals to: CCXIV.

182:679a It has been suggested by Masters [1970. J. Lepid. Soc., 24(3): 207] that *M. cana floridana* be sunk as a junior synonym of *M. bogatana cana*. In subsequent publications by other authors, *M. cana* (Erichson) and *M. bogatana* Felder have been treated as distinct species.

182:CCXVIII Correct Roman numerals to: CCXV.

183: Change Tribe entry as follows:

Tribe Ageroniini Doubleday

1847. Gen. diurn. Lepid., 1(8): 81.

183:CCXIX Correct Roman numerals to: CCXVI.

A major revision of the genus *HAMADRYAS* has been published by Jenkins [1983. Bull. Allyn Mus., (81): 1-146], and a subsequent paper on North American taxa [1984. J. Lepid. Soc., 38(3): 171-175]. Starting after the genus prefatory entry, the species entries should be replaced as follows:

*1. *februa* (Hübner) *AGERONIA*, 1823. Samml. exot. Schmett., 2: pl. [4]. TL - Not stated. Type probably lost [see Note 9].

a. f. *ferentina* (Godart) *NYMPHALIS*, [1824]. In Latreille, Hist Nat. 681a.

[Zool.], 9: 428 (based on figure labeled *Papilio feronia* Cramer, 1782: 140, 249, pl. 362, figs. A, B [nec Linnaeus, 1758]. TL - "East Indies" according to Cramer (1782); Surinam according to Godart [1824]. ST's - Loc. unknown.

= *gudula* Author not stated [Fruhstorfer] AGERONIA, 1914. In Seitz, Grossschmett. Erde, 5: 105d.

= *gudula* (Fruhstorfer) AGERONIA, 1916. In Seitz Grossschmett. Erde, 5: 539. TL - "Texas" (on specimen label); "W. Mexico" (in description). ST's in BM.

= *icilia* (Fruhstorfer) AGERONIA, 1916. In Seitz, Grossschmett. Erde, 5: 540. TL - Belmont, Trinidad. ST's in BM.

= *hierone* (Fruhstorfer) AGERONIA, 1916. In Seitz, Grossschmett. Erde, 5: 540. TL - Yucatán, Mexico. ST's - Loc. unknown; not in BM or ZMHU.

= *fundania* (Fruhstorfer) AGERONIA, 1916. In Seitz, Grossschmett. Erde, 5: 540. TL - San Pedro Sula, Honduras. ST's in BM.

= *fictitia* (F. M. Brown [nec Fruhstorfer]) AGERONIA, 1945. J. New York ent. Soc., 53: 41.

= f. "sodalia" (Fruhstorfer) AGERONIA, In Seitz, Grossschmett. Erde, 5: 540. TL - "E. & S. Mexico". ST's in BM.

*2. *amphichloe* (Boisduval) AGERONIA, 1870. Consid. Lépid. env. Guatemala de l'Orza. Rennes: 27. TL - Guayaquil, Ecuador. ST - BM.

= *tegyra* (Fruhstorfer) AGERONIA, 1916. In Seitz, Grossschmett. Erde, 5: 54. TL - Ecuador. ST - BM.

= *amphichloe* (Fruhstorfer [nec Boisduval, 1870]) AGERONIA, 1916. In Seitz, Grossschmett. Erde, 5: 540.

a. a. *diasia* (Fruhstorfer) AGERONIA, 1916. In Seitz, Grossschmett. 682a. Erde, 5: 541. TL - Puerto Rico and Haiti. ST's - Loc. unknown; not in BM or ZMHU.

= *antillana* (Hall) AGERONIA, 1925. The Entomologist, 58: 187. TL - Hispaniola. ST's in BM; Booth Museum.

*3. *atlantis* (Bates) AGERONIA, 1864. Ent. month. Mag., 1: 115-116. TL - Chuacús, Guatemala. HT in BM.

a. a. *lelaps* (Godman and Salvin) AGERONIA, [1883]. Biol. Centrali-Americanica, Lepid. Rhop., 1: 270. TL - Ventanas, Durango, Mexico. HT in BM.

*4. *feronia* (Linnaeus) PAPILIO, 1758. Syst. Nat. ed. 10, 1: 473. TL - "India". Loc. of type unknown (Uppsala, Linnaean Soc., London?).

a. f. *farinulenta* (Fruhstorfer) AGERONIA, 1916. In Seitz, Grossschmett. Erde, 5: 542. TL - San Pedro Sula, Honduras. ST's in BM. 684a.

= *feronia* (Godman and Salvin [partim, nec Linnaeus, 1758]) PERIDROMIA, [1883]. Biol. Centrali-Americanica, Lepid. Rhop., 1: 273.

= *guatemalena* (Godman and Salvin [partim, nec Linnaeus, 1758]) PERIDROMIA, [1883]. Biol. Centrali-Americanica, Lepid. Rhop., 1: 273.

= *nobilita* (Fruhstorfer) AGERONIA, 1916. In Seitz, Grossschmett. Erde, 5: 543. TL - Chiriquí, Panamá. ST's in BM.

- = *insularis* (Fruhstorfer) *AGERONIA*, 1916. In Seitz, Grossschmett. Erde, 5: 542. TL - Belmont, Trinidad. ST's in BM.
- = *mandragora* (Ménétriés [Fruhstorfer]) *AGERONIA*, 1916. In Seitz, Grossschmett. Erde, 5: 542. TL - "Espirito Santo". ST's in BM.
- *5. *guatemalena* (Bates) *AGERONIA*, 1864. Ent. month. Mag., 1: 115. TL - Central valleys, Guatemala. ST's in BM.
- a. g. *marmorice* (Fruhstorfer) *AGERONIA*, 1916. In Seitz, Grossschmett. Erde, 5: 542. TL - "Orizaba and Guadalajara, Mexico". ST's in BM. **684.1a.**
- *6. *iphthime* (Bates) *AGERONIA*, 1865. Ent. month. Mag., 1: 116. TL - "Columbia and Guatemala". ST's in BM.
- a. i. *joannae* Jenkins, 1983. Bull. Allyn Mus., 81: 72-73. TL - Chiltepec, 684.2a. Oaxaca, Mexico. HT in AME.
- = *iphthime* (Bates) *AGERONIA*, 1864 [*partim*]. Ent. month. Mag., 1: 116. TL - "Polochic Valley, Guatemala and Bogotá, Columbia". ST's in BM.
- = *iphimedea* (Boisduval) *AGERONIA*, 1870. Consid. Lépid. env. Guatemala de l'Orza. Rennes: 26. *Lapsus calami*.
- = *iphthime* (Bates [Fruhstorfer]) *AGERONIA*, 1916. In Seitz, Grossschmett. Erde, 5: 543.
- = *feronia* var. b. *iphthime* (Bates [Kirby]) *AGERONIA*, 1871. Syn. Cat. Lepid., 215.
- *7. *amphinome* (Linnaeus) *PAPILIO*, 1767. Syst. Nat., ed. 12, 1: 779-780. TL - "America meridionali", probably Surinam. HT may be in Uppsala or Linnaean Soc., London.
- a. a. *mexicana* (Lucas) *PERIDROMIA*, 1853. Rev. Zool. [1853]: 331. **685a.**
TL - "Mexico". Loc. of type unknown (MNHN?).
= *amphinome* (Godman and Salvin [*partim, nec Linnaeus, 1767*]) *PERIDROMIA*, [1883]. Biol. Centrali-Americana, Lepid. Rhop., 1: 271.

184:CCXX Correct Roman numerals to: CCXVII.

184:686 Correct type font to: acheronta.

184:CCXXI Correct Roman numerals to CCXVIII; add species as follows:

*2. *blomfildia* (Fabricius) *PAPILIO*, 1781. Spec. Ins., 2: 84. TL - unknown. HT presumed lost.

a. b. *datis* Fruhstorfer, 1907. Int. ent. Zschr. Guben, 1: 319. TL - "Mex- 688.1a. ico and Central America". ST's may be in BM.

185:CCXXII Correct Roman numerals to: CCXIX.

In prefatory entry, make two corrections as follows:

The lit. cit. for *Tymetes* and its type-species should be: Doyère, [1840].

In Cuvier, Règne Anim. (Disciple's edn.) Ins. 2: expl. pl. 138, pl. 139.

Correct spelling to read: *Eunargareta*.

185:689 To conform to the CODE (C. F. Cowan, *in litt.*), the first four lines of the entry must be changed as noted below. We would support a petition to the I.C.Z.N. for the suppression of *zerynthia* Hübner in favor of the well-known and long-used *coresia* (Godart).

1. *zerynthia* Hübner, [1823]. Samml. exot. Schmett., 2: pl. [51]. TL - **689.**

Not stated. Type probably lost [see note 9].

= *coresia* (Godart) *NYMPHALIS*, [1824]. Enc. Meth., 9: 359.

TL - "Brasil". HT may be in MNHN.

186:CCXXXIII Correct Roman numerals to: CCXX.

186:CCXXXIV Correct Roman numerals to: CCXXI.

187:CCXXXV Renumbered as CCXXI, add species as follows:

3. *echemus* (Doubleday and Hewitson) *CYMATOGRAMMA*, [1849]. 697.1.

Gen. diur. Lepid., 2: 316; pl. 49, fig. 4. TL - Honduras. HT in BM.

187:CCXXXV Correct Roman numerals to: CCXXII.

A complete revision of this genus has been undertaken by T. P. Friedlander ["1986(1987)"[1988]. J. Res. Lepid., 25(4): 215-338]. The portions of this work that affect *Memoir No. 2* are presented below and replace the original text.

Genus CCXXI: *Asterocampa* Röber

1916. In Seitz, Grossschmett. Erde, 5: 549. Type-species by designation of D. M. Bates, 1926. Ent. News, 37: 154, *Apatura celtis* Boisduval and Le Conte. "1833" [1835], Hist. gen. Iconogr. Lép. Amér. Sept.: 210 [see Note 608].

1. *celtis* (Boisduval and Le Conte) APATURA "1833" [1835]. Hist. gén. Iconogr. Lép. Amér. sept.: 210. TL - "Géorgie", probably northwest of Savannah, perhaps in Screven Co., Georgia. Type based upon Abbot drawing, the model for which has not been found. Regarding the date of publication, T. P. Friedlander has informed the Committee that [1835] is the best date based upon Cowan's (1969) dating of Plate LVII "*Apatura celtis*" as "1833" [1835].

= *alicia* (W. H. Edwards) APATURA, 1868. Butts. N. America, 1: [135-136]. TL - Vic. New Orleans, Louisiana. Type[s] lost; the figure in W. H. Edwards, 1868. Butts. N. America, 1: pl. Apatura I may be considered to represent types.

a. c. *celtis* (Boisduval and Le Conte) APATURA "1833" [1835]. Hist. 698a. gén. Iconogr. Lép. Amér. sept.: 210. TL - "Géorgie", probably northwest of Savannah, perhaps in Screven Co., Georgia. Type based upon Abbot drawing, the model for which has not been found.

= *alba* (W. G. Wright) APATURA, 1905. Butts. W. Coast: 58. (Change of status).

b. c. *reinthali* Friedlander "1986(1987)"[1988]. J. Res. Lepid., 25(4): 698b. 242-243. Ocoee, Florida. HT in CM.

c. c. *antonia* (W. H. Edwards) APATURA "1877" [1878]. Field and 698c. Forest, 3: 103. TL - Vic. Norse, Bosque Co., Texas; restricted by F. M. Brown, 1967. Trans. American ent. Soc., 93: 377-378. LT in CM, designated by F. M. Brown, *op. cit.*, p. 379.

= *montis* (W. H. Edwards) APATURA, 1883. Papilio, 3: 7. TL - Vic. Ft. Grant, Graham Mtns., Cochise Co., Arizona. LT in CM, designated by F. M. Brown, 1967. Trans. American ent. Soc., 93: 382.

2. *leilia* (W. H. Edwards) APATURA, 1874. Trans. American ent. Soc., 699. 5: 103. TL - "at Camp Lowell and in Sonoto [sic] Valley, Arizona". LT in CM, designated by F. M. Brown, 1967. Trans. American ent. Soc., 93: 385.

= *leila* Skinner, 1891. Ent. News, Suppl. [1891]: i-vi. *Lapsus cal-*

- ami*. Subsequently so misspelled by many authors including Miller and Brown, 1981. Lepid. Soc. Mem. No. 2: 187.
- = *cocles* (Lintner) APATURA, 1885. Papilio, 4: 141. TL - Hidalgo, Texas. LT in CM, designated by T. P. Friedlander, "1986(1987)"[1988]. J. Res. Lepid., 25(4): 255. LT in CM.
3. *clyton* (Boisduval and Le Conte) APATURA, "1833" [1835]. Hist. gén. Iconogr. Lép. Amér. sept.: 208. TL - "méridionales des États-Unis", probably Screven Co., Georgia. Type based upon Abbot drawing, the model for which has not been found [see Note 609].
- a. c. *clyton* (Boisduval and Le Conte) APATURA, "1833" [1835]. Hist. 700a. gén. Iconogr. Lép. Amér. sept.: 208. TL - "méridionales des États-Unis", probably Screven Co., Georgia. Type based upon Abbot drawing, the model for which has not been found [see Note 609].
- = *ocellata* (W. H. Edwards) APATURA, 1876. Butts. N. America, 2: [245]. TL - Coalburgh, West Virginia. LT in CM, designated by F. M. Brown, 1967. Trans. American ent. Soc., 93: 387.
- = *proserpina* (Scudder) APATURA, 1868. Proc. Boston Soc. nat. Hist., 11: 401. TL - Iowa. Type misplaced; not in MCZ.
- = *nig* (J. B. Smith) APATURA, 1903. Check List bor. America: 4. Validating excluded name "Clyton ab. nig." (H. Strecker) APATURA, 1878. Butts. Moths N. America: 145. TL - Berks Co., Pennsylvania. HT in FMNH.
- b. c. *flora* (W. H. Edwards) APATURA, 1876. Butts. N. America, 2: 700b. [247]. TL - Palatka, Florida. LT in CM, designated by F. M. Brown, 1967. Trans. American ent. Soc., 93: 389.
- c. c. *texana* (Skinner) CHLORIPPE, 1911. Trans. American ent. Soc., 700c. 37: 214. TL - Round Mt., Texas. HT in CM.
- = *subpallida* (Barnes and McDunnough) CHLORIPPE, 1913. Contr. nat. Hist. Lepid. N. America, 2: 99. TL - Babaquivera [sic] Mtns., [Pima Co.], Arizona. LT is USNM, designated by T. P. Friedlander, "1987(1987)"[1988]. J. Res. Lepid., 25(4): 263.
- d. c. *louisa* Stallings and Turner, 1947. Ent. News, 58: 38. TL - Pharr, 700d. Texas. HT in YPM.

188:CCXXVI Correct Roman numerals to: CCXXIII.

Family: SATYRIDAE Boisduval

- 188: Correct lit. cit for family name to read:
"1832" [-1834] (12 Oct., 1833). Icon. hist. Lépid. Europe, 1: 166 [see Notes 1 and 612].
- 189: Correct first part of lit. cit. in line 2 to read: Corresp.-Blatt zool.-min...
189:CCXXVII Correct Roman numerals to: CCXXIV.
189:CCXXVIII Correct Roman numerals to: CCXXV.
- 190: Correct lit. cit. for subfamily Satyrinae to read:
"1832" [-1834] (12 Oct., 1833). Icon. hist. Lépid. Europe, 1: 166 [see Notes 1 and 612].
- 190: Correct tribe entry as follows:

Tribe *Euptychiini* Reuter

1896. Acta Soc. Sci. Fennicae, 22: 367-370 (as *Euptychioid*).
 = *Euptychiini* Miller, 1968. Mem. American ent. Soc., 24: 79-95.
- 190:CCXXIX Correct Roman numerals to: CCXXVI.
- 190:715 Correct 1st line of species entry following *EUPTYCHIA* to read:
 , [1867]. Proc. zool. Soc. London 1866, (3): 499.
- 191:CCXXX Correct Roman numerals to: CCXXVII.
- 191:CCXXXI Correct Roman numerals to: CCXXVIII.
- 191:721a To conform to the CODE, emend spelling to: *areolata*.
- 192:CCXXXII Correct Roman numerals to: CCXXIX.
- 192:723b C. G. Oliver [1982. J. Lepid. Soc., 36(2): 153] has suggested that *c. cymela* and *c. viola* may be sibling species. Further study is required, as earlier noted by L. Miller [1976. Bull. Allyn Mus., (33): 4, 7].
- 192:CCXXXIII Correct Roman numerals to: CCXXX.
- 193:CCXXXIV Correct Roman numerals to: CCXXXI.
 The genus *COENONYMPHA* in North America requires considerable additional study with respect to its related species. It is clear that *haydenii* is a distinct species. The remaining taxa have been treated by some authors as subspecies of the *tullia* complex; other authors have treated *tullia* as a superspecies. Brown in compiling the Satyridae section of *Memoir No. 2* followed the general format of the 1964 dos Passos Synonymic List.
- 193:728e Add at end of entry:
 = *bottineauensis* Chermock, Simmons and Chermock, 1963. In Knudson and Post, Butts. Bottineau Co. Pub. No. 2, Dept. Agric. Ent., N. Dakota State Univ.: 23. TL - Bottineau Co., North Dakota. HT in AME.
- 193:729b Insert immediately prior to last line on page:
 = *ochracia* Chermock and Chermock, 1938. Proc. Pennsylvania Acad. Sci., 12: 49-50. *Lapsus calami*.
- 194:729f This taxon was originally described under *ampelos*, but based upon its maculation, it belongs under *ochracea*.
- 194:730e To conform to the CODE, emend spelling to: *insulana*.
- 194:731 Editorial Note: Dornfeld [1980. Butts. Oregon] has stated that *eryngii* and *ampelos* intergrade where these races meet in southwestern Oregon. On this basis, it seems wise to include *californica* and *eryngii* in the *tullia* complex. See Checklist arrangement.
- 195:CCXXXV Correct Roman numerals to: CCXXXII.
- 195:732 Editorial Note: The *Cercyonis pegala* complex requires careful study with regard to the affinities of the described taxa. T. C. Emmel proposed a revision in 1961 [J. Lepid. Soc., 23(3): 165-175]. This arrangement was followed in Howe, 1975 [Butts. N. America]. Scott [1986. Butts. N. America] recognized substantially fewer subspecies than other contemporary authors and has selected the taxon *nephela* (often treated as a form) to replace several subspecies recognized by other authors. Until this complex is sorted out, no major changes are made herein.
- 195:732b The name *maritima* should be placed as a synonym of *p. alope* (732c).
- 196:732l Some field collectors have suggested that *damei* is not a subspecies of *pegala*, but rather a hybrid between *sthenele* and *meadii*.

196:732m The taxon *blanca* is a synonym of *stephensi* (732j) and should be placed as a junior synonym of the latter. See Austin "1985(1986)" [J. Lepid. Soc., 39(2):110].

196:735b Editorial Note: The placement of *silvestris* under *oetus* has been questioned by many collectors from the western U.S. who generally associate *silverstris* as a subspecies of *sithenele*. The lectotype designated by Brown has been carefully examined by both L. D. Miller and C. D. Ferris, and compared against series of California *oetus* and *sithenele*. In their opinion, the lectotype specimen is equivocal and could be placed with either taxon. No change is made here, but it is recognized that a *sithenele* phenotype occurs in southern California to which the name *silvestris* may or may not apply. Emmel and Emmel [1973. Butts. S. California, p. 27] used the combination *C. sithenele silvestris*. Brown [1964. Trans. American ent. Soc., 90: 362] placed *silvestris* under *oetus*.

197:735 Add subspecies:

d. o. *pallescens* Emmel and Emmel, 1971. Pan-Pacif. Ent., 47: 155-157. 735d.

TL - Reese R. Valley, 4 road miles NE of Reese R. crossing on Nevada Hwy. 2, 5700', WSW of Austin, SW corner of T19N, R43E, Lander Co., Nevada. HT in T. C. Emmel Collection, University of Florida.

197:CCXXXVI Correct Roman numerals to: CCXXXIII.

Editorial Note: Scott [1986. Butts. N. America: 243-246] has made several new combinations in genus *Erebia* which are deemed unjustified in view of our present knowledge of North American and Old World fauna. Rather than duplicate comments in the taxonomic section that follows, these taxa are addressed here. The Scott combinations are shown in boldface type.

Erebia magdalena erinnyn. The taxon *erinnyn* was originally described by Staudinger as *erynnis* (= *erinna* Staudinger). In 1932, Warren [Ent. Rec., 44: 166] proposed the replacement name *erinnyn* which is the accusative case of *erynnis* and may not be used. Korshunov [1972. Entomol Rev., 51(1): 93] proposed a second replacement name *sajanensis*. The holotype illustrated by Staudinger [1894. Deut. ent. Zeit. [Iris], 7: 247-248, t. IX, f. 2] is a female. This species is reported to occur in southern Siberia in the eastern Sayan Mtns. and in the Baykal Range. Another *magdalena*-like species is *anjuika* Kurentzov [1966. In New Species in the Fauna of Siberia and Neighboring Regions, Nauka, Novosibirsk: 33-35, figs. 1-2]. This species is reported to occur in the Anyuy and Omsukchan Ranges of NE Siberia. In terms of geographic distribution, *anjuika* is probably the closest Siberian relative to *magdalena* (which has recently been found to occur widely in the Seward Peninsula of Alaska). In any case, since the type of *sajanensis* (= *erynnin*) is a female, and the type of *anjuika* is in the U.S.S.R., it is not possible to study the male genitalia of these species and their relationships to North American *magdalena mackinleyensis*. On this basis and the fact that the appropriate replacement name for *erynnis* Staudinger, 1894 is *sajanensis* Korshunov, 1972, the subspecific epithet *mackinleyensis* Gunder, 1932 must be retained. Ferris et al. [1983. Canadian Ent., 115: 837 and fig. 3] reported the occurrence of *E. sajanensis* (as *erinnyn*) in the Yukon Territory. It is not entirely clear to what species the butterflies illustrated belong. Three specimens, collected at the same time from a single locality, exist in two private collections. Most

probably they represent one of the following conditions: 1. color-reversal of *E. fasciata*; 2. aberrant or heavily maculated *E. magdalena*, which is proving to be extremely variable in the Arctic as to pattern; 3. possible hybrids between *fasciata* and *magdalena*.

Erebia theano pawloskii. This subspecies was described by Ménétriés in 1881 and ranges from Transbaykalia to NE Siberia including the Kamchatka and Amur regions. It is premature to associate it with North American fauna. Some Russian specialists treat *pawloskii* as a distinct species.

Erebia dabanensis youngi. As more than adequately demonstrated by Troubridge and Philip, ["1982(83)". J. Res. Lepid., 21(2): 107-146], *dabanensis* Erschov, 1871 and *youngi* Holland, 1900 are separate species.

Erebia kozhantshikovi lafontainei. Troubridge and Philip (*op. cit.*) clearly demonstrated that *kozhantshikovi* Sheljuzhko, 1925 occurs only in Siberia.

- 198:738 Karsholt and Nielsen, 1986 [Ent. scand., 16:443] have designated a LT for this taxon and fixed the TL as Sweden.
- 198:738a Correct date of publication to: [1849].
- 198:742a This name is preoccupied by *sophia* Acerbi, 1802 [CODE, Art. 58, item (9)], thus the name reverts to *canadensis* Warren, 1931, who correctly proposed the replacement.
- 199:746 Delete existing entry and replace as follows:
11. *lafontainei* Troubridge and Philip, "1982(83)". J. Res. Lepid., 21(2): 746. 130-133. TL - Mt. Decoeli, 1300 m, St. Elias Mtns., Yukon Territory. HT in CNC.
 12. *occulta* Roos and Kimmich, 1983. Entomol. Zeitschr., 93(6): 69-75. 746.1. TL - km 150, Dempster Hwy., Yukon Territory. HT in Kimmich, Siepe and Roos Collection.
= *phellea* Philip and Troubridge, "1982(83)". In Troubridge and Philip, J. Res. Lepid., 21(22): 118-123. TL - km 66-68 Council Rd., 9-11 km NNE of Solomon. HT in USNM. See also Philip, 1985 [J. Res. Lepid., 24(1): 81-82].
 13. *inuitica* Wyatt, 1966. Zeits. Wiener ent. Gesell., 51: 93-94. TL - Anak-746.2. *tuvuk*, Endicott Mtns., Alaska. HT in Colin Wyatt Collection in Karlsruhe Museum, West Germany. This taxon is known only from a single male holotype. The validity of this species has been disputed strongly by Scott [1986]. Butts. N. America: 246], however, the taxon appears to be valid. K. W. Philip kindly provided the Committee with copies of letters exchanged between him and the late Hans Epstein, and between Epstein and Wyatt. The type specimen was collected by a 9-year-old Eskimo boy on 28 June, 1965. It was sent by Wyatt to B. C. S. Warren for examination, who placed it in the *epiphron* group which belongs to the Old World. Based upon the documents examined, there appears to be no reason to suspect that the specimen was mislabeled as to the collection locality. The closest relative to *inuitica* seems to be *E. pharte* Hübner.
- 199:CCXXXVII Correct Roman numerals to: CCXXXIV.
- 200:CCXXXVIII Correct Roman numerals to: CCXXXV.
- 200:748 According to a recent revision of this genus by Austin [1986. Bull. Allyn Mus., (107): 1-27], *dionysus* should be placed as a junior synonym *stretchii*, and three new subspecies added as follows:

- c. r. pallidus Austin, 1986. Bull. Allyn Mus., (107): 10-13. TL - Larkin **748c**. Dry Lake Rd., 4.8 road mi. N. SR 359, Alkalai Valley 7000', R28E, T4N, S12 on Aurora NV-CA 15' quad., Mineral Co., Nevada. HT in Nevada State Museum.
- d. r. minimus Austin, 1986. Bull. Allyn Mus., (107): 14-16. TL - N. side **748d**. of city of Moose Jaw, 50°24'N, 105°30'W, Saskatchewan. HT in AME.
- e. r. neomexicanus Austin, 1986. Bull. Allyn Mus., (107): 16-18. TL - **748e**. Bluewater Can., Zuni Mtns., 7800', Valencia Co., New Mexico. HT in AME.

200:CCXXXIX Correct Roman numerals to: CCXXXVI.

- 200:749 Collecting in California during the past decade has produced series of specimens that grade from *Oeneis ivallda* into *O. chryxus stanislaus* (752f). On this basis, *ivallda* should be treated as *O. chryxus ivallda* and relocated as entry 752g. Editorial Note: dos Passos separated *ivallda* from *chryxus* based upon a chemical test developed in England by Ford [1948. Ent. News: 59: 92-96]. This test involves a temporary color change of anthoxanthin pigments located in butterfly wing scales when they are exposed to the fumes of Ammonium hydroxide (28.2%). The whitish areas on the wings of *ivallda* turned yellowish, while dos Passos reported no color change in *chryxus*. Occasionally pale specimens of *chryxus* that resemble *ivallda* are collected in Alberta and extreme northern British Columbia. C. D. Ferris has subjected specimens of *ivallda*, from various localities, and pale *chryxus* specimens to the test described. The most responsive specimens were from a clinal population taken at Carson Spur in Amador Co., California. Responses to the test by typical *ivallda* were mixed varying from virtually none to strong. Some of the pallid *chryxus* from Canada showed weak responses to this test. On the basis of these results, the validity of this method for species separation is questioned. It does appear to have some validity at the subspecies level. Slight chemical differences can produce drastic color changes in butterfly-wing pigments. Such chemical changes may be controlled by a single gene (such as sexual dimorphism in *Colias*), and also perhaps environmentally induced. Some data are summarized in Ferris, 1985 [Bull. Allyn Mus., (96): 1-51]. In the case at hand, it seems unwise in light of field data to treat *ivallda* as a separate species based upon a possibly unreliable chemical test.
- 201:752d The TL has been restricted to the vicinity of Rock Lake, Alberta (53°27'N, 118°16'W) by Shepard [1984. Quaestiones Entomologicae, 20: 35-44].
- 201:755 In a paper published in 1983 [Canadian Ent., 115: 823-840], Ferris *et al.*, recommended that the species *bore* and *taygete* be combined into a single species *bore*. Recent studies by Ferris in the western arctic have disclosed that two apparent sibling species exist, separable by behavior and phenotype, but not by recognizable characters of the male genitalia (informally reported in "1986(1987)", J. Lepid. Soc., 40(3): 131-138). Until this situation is resolved, it seems best to retain both of these taxa.
- 201:755a Correct date of *bootes* entry to: "1832" [1834].
- 202:756b Add at end of entry:
 = *acerta* T. Emmel, 1975. In Howe, Butts. N. America: 86.
Lapsus calamii.
- 202:757e HT is in AMNH.

- 202:758a Correct date of *oeno* entry to: "1832" [1834].
 202:758b Correct date of *eritiosa* entry to: "1832" [1834].
 202:759 The changes noted below for *O. polixenes* are based upon a recent paper by Troubridge and Parshall [1988. Canadian Ent., 120: 679-696].
 203:759d Comparison of series of *subhyalina* and *peartiae* in the Canadian National Collection in Ottawa has shown that these names apply to identical phenotypes. Thus *peartiae* is placed as a junior synonym of *subhyalina*, which has date priority.
 Replace former 759d with a new subspecies as follows:
 d. p. *luteus* Troubridge and Parshall, 1988. Canadian Ent., 120: 682-683. 759d.
 TL - Pink Mtn., British Columbia, Canada. HT in CNC.
 203:759e Examination of the type series in the Canadian National Collection of this subspecies has shown that it is not what many arctic collectors have called *yukonensis*. True *yukonensis* has pale gray rather diaphanous wings and occurs at high elevation. Typical low-elevation *polixenes luteus* from the Yukon Territory is normally rather dark and heavily scaled.
 203:759 Add new subspecies as follows:
 g. p. *woodi* Troubridge and Parshall, 1988. Canadian Ent., 120: 683-684. 759g.
 TL - Herschel Island, Yukon Territory, Canada. HT in CNC.
 203: At the end of the *Oeneis* entry, add two new species:
 12. *philipi* Troubridge, 1988. In Troubridge and Parshall, 1988. Canadian Ent., 120: 684-686. TL - Km 1 Dempster Highway at Klondike River, Yukon Territory, Canada. HT in CNC.
 13. *excubitor* Troubridge, Philip, Scott and Shepard, 1982. Canadian Ent., 114: 881-889. TL - Hilltop 4 km E Mt. Chambers, Ogilvie Mtns., Yukon Territory. HT in CNC. Scott [1986. Butts. N. America: 251] used the combination *O. alpina excubitor*. While *excubitor* may be the North American representative of *alpina* Kurentzov [1970. Butts. Far Eastern U.S.S.R.: 74, figs. 42, 44], this situation is far from clear. The type specimen of *alpina* came from the Omsukchan Range in Siberia. The published description for this taxon is virtually useless in establishing the true nature of *alpina*. The present location of the type is unknown.

Family DANAIDAE Duponchel

- 203:CCXL Correct Roman numerals to: CCXXXVII.
 Delete references to Notes 659 and Notes 660.
 Based upon the revision by Ackery and Vane-Wright [1984. Milkweed Butterflies: Their Caldistics and Biology], the following entries on pp. 203-204 should be deleted: *Tirumala*, *Chittira*, *Caduga*, *Melinda*, *Nasuma*, *Ravadeba*, *Phirdana*, *Bahora*, *Asthipa*, *Mangalisa*, *Badacara*, *Chlorochrospis*, *Elsa*.
 204: Correct the Note numbers as follows: 661 to 659; 662 to 660; 663 to 661; 664 to 662; 665 to 663.
 204:760 Correct Note reference to 664 from 666.
 205:760 Delete lines 3 and 4 at top of page: = *curassavicae* . . . Kiel.
 205:761 Correct Note 668 to 666.
 205:762 Correct Note references as follows: 668 to 666; 669 to 667.
 205: In text under Ituninae, correct Note numbers as follows: 670 to 668; 671 to 668, 669.

205:CCXLI Correct Roman numerals to: CCXXXVIII.

205:763a Correct Felder citation to: 2: 352.

The entry that follows below should be placed immediately following the text on page 205. All of the Ithomiidae records shown are most probably based upon mislabeled material, but the records do exist and are included here for sake of completeness.

Family ITHOMIIDAE Kirby

1894. In Allen's Naturalist's Library, Butterflies 1: x, 29.

Subfamily Ithomiinae Kirby

1894. In Allen's Naturalist's Library, Butterflies 1: x, 29.

Genus CCXXXIX: Dircenna Doubleday

1847. Gen. diurn. Lepid., 1: 119, pl. 17, fig. 2. Type-species by monotypy *Dircenna iambe* Doubleday, 1847. Gen. diurn. Lepid., 1, pl. 17, fig. 2.

1. *klugii* (Geyer) *CERATINA*, 1837. In Hübner Zutr. z. Samml. exot. 764. Schmett., 5: 5, no. 401, pl. 138, figs. 801-802. TL - "Mexico". HT - presumed lost. For Texas records, see: Kendall & McGuire, 1984. Bull. Allyn. Mus., (86): 40.

Genus CCXL: Greta Hemming

1934. Gen. Names. hol. Butts., 1: 28. Type species by designation by Scudder, through *Hymenitis* Hübner, 1816, *Papilio diaphanus* Drury, 1773, Ill. nat. Hist. 2: pl. 7, fig. 3 & Index, 13. — So placed on the Official List of generic Names by Opinion 985 [Nov., 1972. Bull. zool. Nomencl., 29(3): 117] — and *Hymenitis* Illiger, *Hymenitis* Hübner both suppressed.

*2. *polissenae* (Hewitson) *HYMENITIS*, 1863. Exot. Butts.: 3, pl. *ITHOMIA* 24, fig. 151. TL - "Quito, Ecuador". HT in BM.

a. p. *umbrana* (Haensch) *HYMENITIS*, 1909. In Seitz, Grossschmett. 765a. Erde, 5: 164, pl. 41f. TL - Costa Rica. Loc. of type unknown. For Texas records, see: Kendall & McGuire, 1984. Bull. Allyn Mus., 86: 41.

Editor's Note: *D. klugii* was first recorded from Texas in 1869 (probable date) and again in 1877. The 1877 collection of a single specimen by G. B. Sennett was reported by Lintner [1884. Papilio, 4: 140]. The most recent record is from Brownsville, Cameron Co., Texas in 1904. *G. p. umbrana* was supposedly taken in Texas in either 1869 or 1870. There are no contemporary records for either species. At one time, the former species may have been resident in the lower Rio Grande Valley. It is very doubtful that the latter species was ever resident in Texas.

CORRECTIONS TO NOTES

Citations are by Note number and not page number.

- 32 Delete this note based upon changes to main text.
- 33 Hübner died in mid-1826, so misspellings must be attributed to authors who followed him.
- 36 This text of this note is superseded by: Cowan, C. F., 1970. Annotationes Rhop-alocerologicae 1970: 53.
- 41 Change date of Kirby reference to: 1870.
- 86 Note is misnumbered 96.
- 98 Change entry to read: On this publication, see McHenry, 1962. J. Lepid. Soc., 16(2): 104.
- 122 As noted at the beginning of the **HESPERIOIDEA** section, the publications are: Trans. American ent. Soc., 106: 43-88, 1980; 113: 29-71, 1987.
- 138 Correct plate number to: pl. 31.
- 203 Revise to read: This name is very unsatisfactorily proposed. Holland [1931. Butt. Book: 321], discussed and figured afresh both *carolinianus* G. Edwards, 1898 [*in* Catesby, Nat. Hist. Uncommon Birds, 3rd ed.] and his own *floridensis* of 1898, which he equated with the ["dusky and yellow swallowtail butterfly"] figure by G. E. Edwards (1743) on plate 34 of his Nat. Hist. of Uncommon Birds. No one seems to have accepted Edwards's use of the name as proper, except Holland, and we question whether even Holland's usage of the name can be construed as a proper description.
- 211 Correct volume citation to: 33.
- 212 Art. 45(f) of the CODE means that *kahli* should be interpreted as a subspecific name, since it was not specifically designated as infrasubspecific.
- 227 Delete: "even though the latter . . ." The two names were published on the same page and *Jasoniades* has line priority.
- 234 Correct lit. cit. to read: 38: 26.
- 242 Add: The name *terlootii* is a patronym referring to Baron Terlood who collected widely in Mexico. Thus the emendation by Skinner from *terlooii* to *terlootii* is reasonable.
- 252 Delete this note as no longer applicable.
- 279 Correct this entry to read as follows:
Misidentified as: "*europome*" by Haworth, 1802. Lepid. Britain, 1: 5; as "*nastes*", "*santes*", and "*phicomene*" by Fitch, 1854. Trans. New York St. Agric. Soc., 13:378.
- 306 Insert the following sentence after "(Stoll)" in the 4th line:
Petiver died about 1718 and his work was republished later with perhaps names by an anonymous author.
- 308 In the last paragraph, change "seven species" to: but few species.
- 376 Correct "Westwood" to: Doubleday.
- 405 The I.C.Z.N. ruled on this matter in 1977, Opinion 1073. The name is **RIODINIDAE** Grote, 1895(1827).
- 410 Delete this note.
- 411 The name *Emesis caeneus* (L.) was placed as a rejected name in 1966 by I.C.Z.N. Opinion 755.
- 443 Correct to read: Guérin-Méneville published the misspelled name *Helicona* of Latreille's emended name *Heliconia* in his Icon. Règne Anim. Cuvier, 3: 472 [1844].
- 445 Add as follows: C. F. Cowan (*in litt.*) has stated that there is no evidence that p. 757 and p. [1339] were published other than simultaneously, and therefore the preferred spell-

- ing depends upon the First Reviser.
- 447 Correct "lapsi" to: *lapsus*.
- 464 Correct reference to Note 453 to read: 458.
- 469 Correct to read: *Brenthis gibsoni* and *lehmanni* were both proposed to replace *Argynnis alaskensis* Lehmann [in Seitz, 1913. Grossschmett. Erde, 5: 424], believed to be preoccupied by *Brenthis alaskensis* Holland [1900. Ent. News, 9: 383]. Since this action was prior to 1961, this homonymy must stand [CODE, Art. 59(b), (c)].
- 472 Correct "lücki" to read: "lücki" . . .
- 497 According to C. F. Cowan (*in litt.*), *Chlosyne dorothya*e meets the CODE Criteria of Availability as listed in Arts. 10 to 20 (Chapter IV).
- 500 Correct Boisduval citation to read: *in Roret, 1836. Hist. Ins., Sp. gen. Lépid.*, pl. 11 (= 7b), fig. 8.
- 505 Correct first part of entry to read: Name misspelled or emended as . . .
- 506 Add to note: Godart noted that his *tharossa* (two examples) had white annuli. Recent work by Charles Oliver (*in. litt.*) has shown that a key character is the color of the antennal clubs. In multivoltine *tharos* the color is normally black (but may vary in some western populations), while in the univoltine populations, the color is orange dorsally.
- 508 Correct first part of entry to read: Name misspelled or emended to . . .
- 517 Note reference should be to: Note 497.
- 537 Correct first part of entry to read: Name variously emended or misspelled as "phaetaena" by Hübner, [1819]. Verz. bekannt. Schmett., (2): 28; as "phaetontea" by many . . .
- 567 Add: This species has been misreported from Florida as *A. lytrea*.
- 578 Emend entry to read:
The hybrid of *weidemeyerii latifascia* x *l. lorquini* is known as . . . affinity between *weidemeyerii* and *lorquini*.
- 616 In 5th line, change "Nymphaloidea" to read: nymphaloid complex.
- 620 Replace text by the following: Godart [1824]. Enc. Meth. 9(2): 493 treated two separate taxa; No. 55 *Satyrus canthus* (Linnaeus, 1767) [= *Papilio eurydice* Linnaeus, 1763 (*in Johansson*)], from Philadelphia, giving a translation of the Johansson description, and No. 56 *Satyrus cantheus* Godart [*nom. nov.*] for *Papilio canthus* Fabricius, 1775 from North America, which he considered to be different. Thus *cantheus* Godart was a new name for *canthus* Fabricius *nec* Linnaeus, and is an available name. The name *cantheus* Godart is not a *lapsus calami* as originally stated.
- 651 Add to present text: John B. Heppner [1982. Archiv. nat. Hist., 10(2)] has dated "Esper" thoroughly. He places plate 100 as 1789 and plate 108 as [1798]. The 1805 issue was Supplement 2, with plates 117-122.
- 669 Correct number type font to bold face.

CORRECTIONS TO INDEX

Page No.

- 248 atrapraetextus misspelled as "atropraetextus". Correct spelling and position of entry.
249 Boloria. Correct note reference to N462.
252 coloro misspelled as "chloro". Correct spelling and position of entry.
Entry out of order; place "clarus" before "claudia".
256 endymion. Species reference number should be 493, not 483.
258 florenceae misspelled "florencae".
258 g-argenteum is misspelled. It should be c-argenteum. 645. Relocate under C.
260 harkenclenus should be: Harkenclenus.
261 Entries: hopfferi-hopfingeri are misplaced. Insert them after "homoperplexa".
265 martini, 163c. Correct spelling to Hesperia.
266 Mestra. Correct Roman numeral to: CCXVII (revised number CCXIV).
273 Entry correctly placed, but correct spelling of "Purginae" to: Pyrginae.
278 Add at end of U listings: utahensis (Speyeria). 573j.

NOTES

PART III

CHECKLIST OF BUTTERFLIES OF NORTH AMERICA
NORTH OF MEXICO

This list is a basic checklist only and contains no synonymies. Species numbers are shown in boldface type to the right of each entry. When a number differs from the original usage in *Memoir No. 2*, the old number is shown in parentheses. Gaps in numbering denote deleted species; see Part II for discussion. Some new genera have been added to this Supplement and are shown in ***boldface italic*** type. Date citations follow Art. 22 of the CODE in accordance with ¶ 1 of Part I.

HESPERIOIDEA

<i>Pyrrhopyge araxes arizoneae</i> Godman and Salvin, [1893].	1a.
<i>Phocides pigmalion okeechobee</i> (Worthington, 1881).	2a.
<i>Phocides palemon lilea</i> (Reakirt, [1867]).	3a.
<i>Phocides urania</i> (Westwood, [1852]).	4.
<i>Proteides mercurius mercurius</i> (Fabricius, 1787).	5a.
<i>Proteides mercurius sanantonio</i> (Lucas, 1857).	5b.
<i>Epargyreus zestos</i> (Geyer, 1832).	6.
<i>Epargyreus clarus clarus</i> (Cramer, [1775]).	7a.
<i>Epargyreus clarus huachuca</i> Dixon, 1955.	7b.
<i>Epargyreus clarus californicus</i> MacNeill, 1975.	7c.
<i>Epargyreus exadeus cruza</i> Evans, 1952.	8a.
<i>Polygonus leo histrio</i> Röber, 1925.	9a. (9).
<i>Polygonus manueli</i> Bell and W. P. Comstock, 1948.	10.
<i>Chioides catillus albofasciatus</i> (Hewitson, [1867]).	11a.
<i>Chioides zilpa zilpa</i> (Butler), 1874.	12a.
<i>Chioides zilpa namba</i> Evans, 1952.	12b.
<i>Aguna asander</i> (Hewitson, [1867]).	13.
<i>Aguna claxon</i> Evans, 1952.	14.
<i>Aguna metophis</i> (Latreille, [1824]).	15.
<i>Typhedanus undulatus</i> (Hewitson, [1867]).	16.
<i>Polythrix mexicana</i> H. A. Freeman, 1969.	17.
<i>Polythrix octomaculata</i> (Sepp, 1848).	18.
<i>Polythrix procura</i> (Plötz, 1880).	19.
<i>Zestusa dorus</i> (W. H. Edwards), 1882.	20.
<i>Codatracus alcaeus</i> (Hewitson, [1867]).	21.
<i>Codatracus melon</i> (Godman and Salvin, 1893).	22.
<i>Codatracus arizonensis</i> (Skinner, 1905).	23.
<i>Urbanus proteus</i> (Linnaeus, 1758).	24.
<i>Urbanus pronus</i> Evans, 1952.	25.
<i>Urbanus esmeraldus</i> (Butler, 1877).	26.
<i>Urbanus dorantes dorantes</i> (Stoll, [1790]).	27a.
<i>Urbanus dorantes santiago</i> (Lucas, 1862).	27b.
<i>Urbanus teleus</i> (Hübner, [1821]).	28.
<i>Urbanus tanna</i> Evans, 1952.	29.
<i>Urbanus simplicius</i> (Stoll, [1790]).	30.
<i>Urbanus procne</i> (Plötz, 1880).	31.

<i>Urbanus doryssus</i> (Swainson, 1821).	32.
<i>Astraptes fulgerator azul</i> (Reakirt, [1867]).	34a.
<i>Astraptes egregius</i> (Butler, [1870]).	35.
<i>Astraptes alardus latia</i> Evans, 1952	36a.
<i>Astraptes gilberti</i> H. A. Freeman, 1969.	37.
<i>Astraptes galesus cassius</i> Evans, 1952.	38a.
<i>Astraptes anaphus annetta</i> Evans, 1952.	39a.
<i>Autochton cellus</i> (Boisduval and Le Conte, [1837]).	40.
<i>Autochton pseudocellus</i> (Coolidge and Clemence, 1911).	41.
<i>Autochton cinctus</i> (Plötz, 1882).	41.1.
<i>Achalarus lyciades</i> (Geyer, [1832]).	42.
<i>Achalarus casica</i> (Herrich-Schäffer, 1869).	43.
<i>Achalarus albociliatus</i> (Mabille, 1877).	44.
<i>Achalarus toxaeus</i> (Plötz, 1882).	45.
<i>Achalarus jalapus</i> (Plötz, 1882).	46.
<i>Thorybes bathyllus</i> (J. E. Smith, 1797).	47.
<i>Thorybes pylades</i> (Scudder, 1870).	48.
<i>Thorybes diversus</i> Bell, 1927.	49.
<i>Thorybes mexicanus mexicanus</i> (Herrich-Schäffer, 1869).	50a.
<i>Thorybes mexicanus nevada</i> Scudder, [1871].	50b.
<i>Thorybes mexicanus dobra</i> Evans, 1952.	50c.
<i>Thorybes mexicanus blanco</i> Scott, 1981.	50d.
<i>Thorybes confusis</i> Bell, 1922.	51.
<i>Thorybes drusius</i> (W. H. Edwards, 1883).	52.
? <i>Thorybes valerianus</i> (Plötz, 1882).	53.
<i>Cabares potrillo</i> (Lucas, 1857).	54.
<i>Celaenorrhinus fritzgaertneri</i> (Bailey, 1880).	55.
<i>Celaenorrhinus stallingsi</i> H. A. Freeman, 1946.	56.
<i>Dyscophellus euribates</i> (Stoll, [1782]).	57.
<i>Spathilepia clonius</i> (Cramer, [1776]).	58.
<i>Cogia calchas</i> (Herrich-Schäffer, 1869).	59.
<i>Cogia hippalus</i> (W. H. Edwards, 1862).	60.
<i>Cogia ouis</i> (Skinner, 1894).	61.
<i>Cogia caicus caicus</i> (Herrich-Schäffer, 1869).	62a.
<i>Cogia caicus moschus</i> (W. H. Edwards, 1882).	62b.
<i>Cogia mysie</i> (Dyar, 1904).	62.1.
<i>Arteurotia tractipennis</i> <i>tractipennis</i> Butler and Druce, 1872.	62.5a.
<i>Nisoniades rubescens</i> (Möschler, 1876).	63.
<i>Pellicia angra</i> Evans, 1953.	64.
<i>Pellicia arina</i> Evans, 1953.	65.
<i>Pellicia dimidiata</i> <i>dimidiata</i> Herrich-Schäffer, 1870.	65.1a.
<i>Bolla clytius</i> (Godman and Salvin, 1897).	66.
<i>Bolla brennus</i> (Godman and Salvin, 1896).	67.
<i>Staphylus ceos</i> (W. H. Edwards, 1870).	68.
<i>Staphylus mazans</i> (Reakirt, 1866).	69.
<i>Staphylus hayhurstii</i> (W. H. Edwards, 1870).	70.
<i>Staphylus azteca</i> (Scudder, 1872).	70.1.
<i>Gorgythion begga pyralina</i> (Möschler, 1876).	71a.
<i>Sostrata bifasciata nordica</i> Evans, 1953.	72a.
<i>Carrhenes canescens</i> (R. Felder, 1869).	73.

<i>Xenophanes tryxus</i> (Stoll, [1780]).	74.
<i>Antigonus emorsus</i> (R. Felder, 1869).	74.5.
<i>Systasea pulverulenta</i> (R. Felder, 1869).	75.
<i>Systasea zampa</i> (W. H. Edwards, 1876).	76.
<i>Achlyodes thraso tamenund</i> (W. H. Edwards, 1871).	77a.
<i>Grais stigmatica</i> (Mabille), 1883.	78.
<i>Timochares ruptifasciatus</i> (Plötz, 1884).	79.
<i>Chiomara asychis georgina</i> (Reakirt, 1868).	80a.
<i>Gesta gesta invisus</i> (Butler and H. Druce, 1872).	81a.
<i>Ephyriades brunneus floridensis</i> Bell and W. P. Comstock, 1948.	82a.
<i>Erynnis icelus</i> (Scudder and Burgess, 1870).	83.
<i>Erynnis brizo brizo</i> (Boisduval and Le Conte, 1834).	84a.
<i>Erynnis brizo somnus</i> (Lintner, 1881).	84b.
<i>Erynnis brizo burgessi</i> (Skinner, 1914).	84c.
<i>Erynnis brizo lacustra</i> (W. G. Wright, 1905).	84d.
<i>Erynnis juvenalis juvenalis</i> (Fabricius, 1793).	85a.
<i>Erynnis juvenalis clitus</i> (W. H. Edwards, 1882).	85b.
<i>Erynnis telemachus</i> Burns, 1960.	86.
<i>Erynnis propertius</i> (Scudder and Burgess, 1870).	87.
<i>Erynnis meridianus</i> Bell, 1927.	88.
<i>Erynnis scuddereri</i> (Skinner, 1914).	89.
<i>Erynnis horatius</i> (Scudder and Burgess, 1870).	90.
<i>Erynnis tristis tristis</i> (Boisduval, 1852).	91a.
<i>Erynnis tristis tatus</i> (W. H. Edwards, 1882).	91b.
<i>Erynnis martialis</i> (Scudder, 1869).	92.
<i>Erynnis pacuvius pacuvius</i> (Lintner, [1876]).	93a.
<i>Erynnis pacuvius lilius</i> (Dyar, 1904).	93b.
<i>Erynnis pacuvius perniger</i> (Grinnell, 1905).	93c.
<i>Erynnis pacuvius callidus</i> (Grinnell, 1904).	93d.
<i>Erynnis zarucco</i> (Lucas, 1857).	94.
<i>Erynnis funeralis</i> (Scudder and Burgess, 1870).	95.
<i>Erynnis lucilius</i> (Scudder and Burgess), 1870.	96.
<i>Erynnis baptisiae</i> (Forbes, 1936).	97.
<i>Erynnis afranius</i> (Lintner, [1876]).	98.
<i>Erynnis persius persius</i> (Scudder, 1863).	99a.
<i>Erynnis persius borealis</i> (Cary, 1907).	99b.
<i>Erynnis persius avinoffi</i> (Holland, 1930).	99c.
<i>Erynnis persius fredericki</i> H. A. Freeman, 1943.	99d.
<i>Pyrgus centaureae freija</i> (Warren, 1924).	100a.
<i>Pyrgus centaureae wyandot</i> (W. H. Edwards, 1863).	100b.
<i>Pyrgus centaureae loki</i> Evans, 1953.	100c.
<i>Pyrgus ruralis ruralis</i> (Boisduval, 1852).	101a.
<i>Pyrgus ruralis lagunae</i> Scott, 1981.	101b.
<i>Pyrgus xanthus</i> W. H. Edwards, 1878.	102.
<i>Pyrgus scriptura</i> (Boisduval, 1852).	103.
<i>Pyrgus communis</i> (Grote, 1872).	104.
<i>Pyrgus albescens</i> Plötz, 1884.	105.
<i>Pyrgus oileus</i> (Linneaus, 1767).	106.
<i>Pyrgus philetas</i> W. H. Edwards, 1881.	107.
<i>Heliopetes domicella</i> (Erichson, 1848).	108.

<i>Heliopetes ericetorum</i> (Boisduval, 1852).	109.
<i>Heliopetes lavianus</i> (Hewitson, [1868]).	110.
<i>Heliopetes macaira</i> (Reakirt, 1866).	111.
<i>Heliopetes arsalte</i> (Linnaeus, 1758).	112.
<i>Celotes nessus</i> (W. H. Edwards, 1877).	113.
<i>Celotes limpia</i> Burns, 1974.	114.
<i>Pholisora catullus</i> (Fabricius, 1793).	115.
<i>Pholisora mejicana</i> (Reakirt, 1866).	116.
<i>Hesperopsis libya libya</i> (Scudder, 1878).	117a.
<i>Hesperopsis libya lena</i> (W. H. Edwards, 1882).	117b.
<i>Hesperopsis alpheus alpheus</i> (W. H. Edwards, 1876).	118a.
<i>Hesperopsis alpheus oricus</i> (W. H. Edwards, 1879).	118b.
<i>Hesperopsis alpheus texanus</i> Scott, 1981.	118c.
<i>Hesperopsis gracielae</i> (MacNeill, 1970).	119.
<i>Carterocephalus palaemon mandan</i> (W. H. Edwards, 1863).	120a.
<i>Piruna pirus</i> (W. H. Edwards, 1878).	121.
<i>Piruna polingii</i> (Barnes, 1900).	122.
<i>Piruna microstictus</i> (Godman, 1900).	123.
<i>Piruna haferniki</i> H. A. Freeman, 1970.	124.
<i>Synapte malitiosa pecta</i> (Herrich-Schäffer, 1865).	125.
<i>Synapte salenus</i> (Mabille, 1883).	126.
<i>Synapte syraces</i> (Godman, [1901]).	126.1.
<i>Corticea corticea</i> (Plötz, 1883).	127.
<i>Callimormus saturnus</i> (Herrich-Schäffer, 1869).	128.
<i>Vidius perigenes</i> (Godman, 1900).	129.
<i>Monca tyrtaeus</i> (Plötz, 1883).	130.
<i>Nastraea lherminier</i> (Latreille, [1824]).	131.
<i>Nastraea julia</i> (H. A. Freeman, 1945).	132.
<i>Nastraea neamatyla</i> (Skinner and R. C. Williams, 1923).	133.
<i>Cymaenes tripunctus</i> (Herrich-Schäffer, 1865).	134.
<i>Cymaenes odilia trebiius</i> (Mabille, 1891).	135a.
<i>Lerema accius</i> (J. E. Smith, 1797).	136.
<i>Lerema liris</i> Evans, 1955	137.
<i>Vettius fantasos</i> (Stoll, 1780).	137.5.
<i>Perichares philetas adela</i> (Hewitson, [1867]).	138a.
<i>Rhinthon osca</i> (Plötz, 1883).	139.
<i>Decinea perciosius</i> (Godman, 1900).	140.
<i>Decinea huasteca</i> (H. A. Freeman, 1966).	140.1.
<i>Conga chydaea</i> (Butler, 1877).	141.
<i>Ancyloxypha numitor</i> (Fabricius, 1793).	142.
<i>Ancyloxypha arene</i> (W. H. Edwards, 1871).	143.
<i>Oarisma powersheik</i> (Parker, 1870).	144.
<i>Oarisma garita</i> (Reakirt, 1866).	145.
<i>Oarisma edwardsii</i> (Barnes, 1897).	146.
<i>Copaeodes aurantiacus</i> (Hewitson), [1868].	147.
<i>Copaeodes minimus</i> (W. H. Edwards, 1870).	148.
<i>Adopaeodes prittwitzi</i> (Plötz, 1884).	149.
<i>Thymelicus lineola</i> (Ochsenheimer, 1808).	150.
<i>Hylephila phyleus phyleus</i> (Drury, [1773]).	151a.
<i>Hylephila phyleus muertovalle</i> Scott, 1981.	151b.

<i>Yvretta rhesus</i> (W. H. Edwards, 1878).	152.
<i>Yvretta carus carus</i> (W. H. Edwards, 1883).	153a.
<i>Yvretta carus subreticulata</i> (Plötz, 1883).	153b.
<i>Pseudocopaeodes eunus eunus</i> (W. H. Edwards, 1881).	154a.
<i>Pseudocopaeodes eunus wrightii</i> (W. H. Edwards, 1882).	154b.
<i>Pseudocopoaeedes eunus alinea</i> Scott, 1981.	154c.
<i>Stinga morrisoni</i> (W. H. Edwards), 1878.	155.
<i>Hesperia uncas uncas</i> W. H. Edwards, 1863.	156a.
<i>Hesperia uncas lasus</i> (W. H. Edwards, 1881).	156b.
<i>Hesperia uncas macswaini</i> MacNeill, 1964.	156c.
<i>Hesperia juba</i> (Scudder, [1871]).	157.
<i>Hesperia comma manitoba</i> (Scudder, 1874).	158a.
<i>Hesperia comma assiniboia</i> (Lyman, 1892).	158b.
<i>Hesperia comma laurentina</i> (Lyman, 1892).	158c.
<i>Hesperia comma borealis</i> Lindsey, 1942.	158d.
<i>Hesperia comma harpalus</i> (W. H. Edwards, 1881).	158e.
<i>Hesperia comma yosemite</i> Leussler, 1933.	158f.
<i>Hesperia comma leussleri</i> Lindsey, 1940.	158g.
<i>Hesperia comma tildeni</i> H. A. Freeman, [1956].	158h.
<i>Hesperia comma dodgei</i> (Bell, 1927).	158i.
<i>Hesperia comma oregonia</i> (W. H. Edwards, 1883).	158j.
<i>Hesperia comma hubbirti</i> Lindsey, 1939.	158k.
<i>Hesperia comma ochracea</i> Lindsey, 1941.	158l.
<i>Hesperia comma colorado</i> (Scudder, 1874).	158m.
<i>Hesperia comma susanae</i> L. Miller, 1962.	158n.
<i>Hesperia comma oroplate</i> Scott, 1981.	158o.
<i>Hesperia woodgatei</i> (R. C. Williams, 1914).	159.
<i>Hesperia ottoe</i> W. H. Edwards, 1866.	160.
<i>Hesperia leonardus leonardus</i> Harris, 1862.	161a.
<i>Hesperia leonardus pawnee</i> Dodge, 1874.	161b, (162).
<i>Hesperia leonardus montana</i> (Skinner, 1911).	161c, (161a).
<i>Hesperia pahaska pahaska</i> (Leussler, 1938).	163a.
<i>Hesperia pahaska williamsi</i> Lindsey, 1938.	163b.
<i>Hesperia pahaska martini</i> MacNeill, 1964.	163c.
<i>Hesperia columbia</i> (Scudder, 1872).	164.
<i>Hesperia metea metea</i> Scudder, 1864.	165a.
<i>Hesperia metea licinus</i> (W. H. Edwards, 1871).	165b.
<i>Hesperia viridis</i> (W. H. Edwards), 1883.	166.
<i>Hesperia attalus attalus</i> (W. H. Edwards, 1871).	167a.
<i>Hesperia attalus slossonae</i> (Skinner, 1890).	167b.
<i>Hesperia meskei meskei</i> (W. H. Edwards, 1877).	168a.
<i>Hesperia meskei straton</i> (W. H. Edwards, 1881).	168b.
<i>Hesperia dacotae</i> (Skinner, 1911).	169.
<i>Hesperia lindseyi</i> (Holland, 1930).	170.
<i>Hesperia sassacus sassacus</i> Harris, 1862.	171a.
<i>Hesperia sassacus manitoboides</i> (Fletcher, [1888]).	171b.
<i>Hesperia miriamae</i> MacNeill, 1959.	172.
<i>Hesperia nevada</i> (Scudder, 1874).	173.
<i>Polites peckius</i> (W. Kirby, 1837).	174.
<i>Polites sabuleti sabuleti</i> (Boisduval, 1852).	175a.

<i>Polites sabuleti tecumseh</i> (Grinnell, 1903).	175b.
<i>Polites sabuleti chusca</i> (W. H. Edwards, 1873).	175c.
<i>Polites sabuleti ministigma</i> Scott, 1981.	175d.
<i>Polites sabuleti genoa</i> (Plötz), 1883.	175e.
<i>Polites sabuleti alkaliensis</i> Austin, 1987.	175f.
<i>Polites sabuleti albamontana</i> Austin, 1987.	175g.
<i>Polites sabuleti sinemaculata</i> Austin, 1987.	175h.
<i>Polites sabuleti basinensis</i> Austin, 1988.	175i.
<i>Polites sabuleti nigrescens</i> Austin, 1987.	175j.
<i>Polites mardon</i> (W. H. Edwards, 1881).	176.
<i>Polites draco</i> (W. H. Edwards, 1871).	177.
<i>Polites baracoa</i> (Lucas, 1857).	178.
<i>Polites themistocles</i> (Latreille, [1824]).	179.
<i>Polites origenes origenes</i> (Fabricius, 1793).	180a.
<i>Polites origenes rhena</i> (W. H. Edwards, 1878).	180b.
<i>Polites mystic mystic</i> (W. H. Edwards, 1863).	181a.
<i>Polites mystic dacotah</i> (W. H. Edwards, 1871).	181b.
<i>Polites sonora sonora</i> (Scudder, 1872).	182a.
<i>Polites sonora siris</i> (W. H. Edwards, 1881).	182b.
<i>Polites sonora utahensis</i> (Skinner, 1911).	182c.
<i>Polites vibex vibex</i> (Geyer, [1832]).	183a.
<i>Polites vibex praeceps</i> (Scudder, 1872).	183b.
<i>Polites vibex brettoides</i> (W. H. Edwards, 1883).	183c.
<i>Wallengrenia otho otho</i> (J. E. Smith, 1797).	[see comment in Part II]
<i>Wallengrenia egeremet</i> (Scudder, 1864).	184.
<i>Pompeius verna</i> (W. H. Edwards), 1862.	185.
<i>Atalopedes campestris campestris</i> (Boisduval, 1852).	186.
<i>Atalopedes campestris huron</i> (W. H. Edwards, 1863).	187a.
<i>Atrytone arogos arogos</i> (Boisduval and Le Conte, [1834]).	187b.
<i>Atrytone arogos iowa</i> (Scudder, 1869).	188a.
<i>Atrytone logan logan</i> (W. H. Edwards, 1863).	188b.
<i>Atrytone logan lagus</i> (W. H. Edwards, 1881).	189a.
<i>Atrytone mazai</i> H. A. Freeman, 1969.	189b.
<i>Atrytone potosiensis</i> H. A. Freeman, 1969.	189.1.
<i>Problema byssus byssus</i> (W. H. Edwards, 1880).	189.2.
<i>Problema byssus kurnskaka</i> (Scudder, 1887).	190a.
<i>Problema bulenta</i> (Boisduval and Le Conte, [1834]).	190b.
<i>Ochlodes sylvanoides sylvanoides</i> (Boisduval, 1852).	191.
<i>Ochlodes sylvanoides pratincola</i> (Boisduval, 1852).	192a.
<i>Ochlodes sylvanoides napa</i> (W. H. Edwards, 1865).	192b.
<i>Ochlodes sylvanoides santacruzus</i> Scott, 1981.	192c.
<i>Ochlodes sylvanoides oreocoastus</i> Scott, 1981.	192d.
<i>Ochlodes sylvanoides bonnevillus</i> Scott, 1981.	192e.
<i>Ochlodes agricola agricola</i> (Boisduval, 1852).	192f.
<i>Ochlodes agricola verus</i> (W. H. Edwards, 1881).	193a.
<i>Ochlodes agricola nemorum</i> (Boisduval, 1852).	193b.
<i>Ochlodes snowi</i> (W. H. Edwards, 1877).	193c.
<i>Ochlodes yuma</i> (W. H. Edwards, 1873).	194.
<i>Poanes massasoit massasoit</i> (Scudder, 1864).	195.
<i>Poanes massasoit chermocki</i> Anderson and Simmons, 1976.	196a.
	196b.

<i>Poanes hobomok hobomok</i> (Harris, 1862).	197a.
<i>Poanes hobomok wetona</i> Scott, 1981.	197b.
<i>Poanes zabulon</i> (Boisduval and Le Conte, [1834]).	198.
<i>Poanes taxiles</i> (W. H. Edwards, 1881).	199.
<i>Poanes aaroni aaroni</i> (Skinner, 1890).	200a.
<i>Poanes aaroni howardi</i> (Skinner, 1896).	200b.
<i>Poanes yehl</i> (Skinner), 1893.	201.
<i>Poanes viator viator</i> (W. H. Edwards, 1865).	202a.
<i>Poanes viator zizaniae</i> Shapiro, 1971.	202b.
<i>Paratrytone melane melane</i> (W. H. Edwards, 1869).	203a.
<i>Paratrytone melane vitellina</i> (Herrick-Schäffer, 1869).	203b.
<i>Choranthus radians</i> (Lucas, 1856).	204.
<i>Choranthus haitensis</i> (Skinner, 1920).	205.
<i>Choranthus vitellius</i> (Fabricius, 1793)	205.1.
<i>Mellana eulogius</i> (Plötz, 1883).	206.
<i>Mellana mexicana</i> (Bell, 1942).	207.
<i>Euphyes arpa</i> (Boisduval and Le Conte, [1834]).	208.
<i>Euphyes pilatka</i> (W. H. Edwards, 1867).	209.
<i>Euphyes dion</i> (W. H. Edwards, 1879).	210.
<i>Euphyes alabamae</i> (Lindsey, 1923).	211.
<i>Euphyes dukesi</i> (Lindsey, 1923).	212.
<i>Euphyes conspicuus conspicuus</i> (W. H. Edwards, 1863).	213a.
<i>Euphyes conspicuus buchholzi</i> (Ehrlich and Gillham, 1951).	213b.
<i>Euphyes berryi</i> (Bell, 1941).	214.
<i>Euphyes macguirei</i> H. A. Freeman, 1975.	215.
<i>Euphyes bimacula bimacula</i> (Grote and Robinson, 1867).	216a, (216).
<i>Euphyes bimacula illinois</i> (Dodge, 1872).	216b, (216).
<i>Euphyes vestris vestris</i> (Boisduval, 1852).	217a.
<i>Euphyes vestris metacomet</i> (Harris, 1862).	217b.
<i>Euphyes vestris kiowah</i> (Reakirt, 1866).	217c.
<i>Euphyes vestris harbisoni</i> J. W. Brown and McGuire, 1983.	217d.
<i>Asbolis capucinus</i> (Lucas, 1857).	218.
<i>Atrytonopsis hianna hianna</i> (Scudder, 1868).	219a.
<i>Atrytonopsis hianna turneri</i> H. A. Freeman, 1948.	219b.
<i>Atrytonopsis deva</i> (W. H. Edwards, 1876).	220.
<i>Atrytonopsis lunus</i> (W. H. Edwards, 1884).	221.
<i>Atrytonopsis vierecki</i> (Skinner, 1902).	222.
<i>Atrytonopsis loammi</i> (Whitney, 1876).	223.
<i>Atrytonopsis pittacus</i> (W. H. Edwards, 1882).	224.
<i>Atrytonopsis python</i> (W. H. Edwards, 1882).	225.
<i>Atrytonopsis cestus</i> (W. H. Edwards, 1884).	226.
<i>Atrytonopsis edwardsi</i> Barnes and McDunnough, 1916.	227.
<i>Amblyscirtes simius</i> W. H. Edwards, 1881.	228.
<i>Amblyscirtes exotera</i> (Herrick-Schäffer, 1869).	229.
<i>Amblyscirtes cassus</i> W. H. Edwards, 1883.	230.
<i>Amblyscirtes aenus</i> W. H. Edwards, 1878.	231.
<i>Amblyscirtes linda</i> H. A. Freeman, 1943.	232.
<i>Amblyscirtes oslari</i> (Skinner, 1899).	233.
<i>Amblyscirtes elissa</i> Godman, [1900].	234.
<i>Amblyscirtes hegona</i> (Scudder, 1864).	235.

<i>Amblyscirtes texanae</i> Bell, 1927.	236.
<i>Amblyscirtes tolteca</i> Scudder, 1872.	237.
<i>Amblyscirtes prenda</i> Evans, 1955.	238.
<i>Amblyscirtes aesculapius</i> (Fabricius, 1793).	239.
<i>Amblyscirtes carolina</i> (Skinner, 1892).	240.
<i>Amblyscirtes reversa</i> Jones, 1926.	241.
<i>Amblyscirtes nereus</i> (W. H. Edwards, 1876).	242.
<i>Amblyscirtes nysa</i> W. H. Edwards, 1877.	243.
<i>Amblyscirtes eos</i> (W. H. Edwards, 1871).	244.
<i>Amblyscirtes vialis</i> (W. H. Edwards, 1862).	245.
<i>Amblyscirtes celia</i> Skinner, 1895.	246.
<i>Amblyscirtes belli</i> H. A. Freeman, 1941.	247.
<i>Amblyscirtes alternata</i> (Grote and Robinson, 1867).	248.
<i>Amblyscirtes phylace</i> (W. H. Edwards, 1878).	249.
<i>Amblyscirtes fimbriata</i> (Plötz, 1882).	250.
<i>Lerodea eufala</i> (W. H. Edwards, 1869).	251.
<i>Lerodea arabus</i> (W. H. Edwards, 1882).	252.
<i>Lerodea dysaules</i> Godman, [1900].	253.
<i>Oligoria maculata</i> (W. H. Edwards, 1865)	254.
<i>Calpodes ethlius</i> (Stoll, [1782]).	255.
<i>Panoquina panoquin</i> (Scudder, 1864).	256.
<i>Panoquina panoquinoides</i> (Skinner, 1891).	257.
<i>Panoquina errans</i> (Skinner, 1892).	258.
<i>Panoquina ocola</i> (W. H. Edwards, 1863).	259.
<i>Panoquina hecebola</i> (Scudder, 1872).	260.
<i>Panoquina sylvicola</i> (Herrich-Schäffer, 1865).	261.
<i>Panoquina evansi</i> (H. A. Freeman, 1946).	262.
<i>Nyctelius nyctelius</i> (Latreille, [1824]).	263.
<i>Thespieus macareus</i> (Herrich-Schäffer, 1869).	264.
<i>Agathymus neumoegeni neumoegeni</i> (W. H. Edwards, 1882).	265a, (265).
<i>Agathymus neumoegeni carlsbadensis</i> (D. Stallings and Turner, 1957).	265b, (266).
<i>Agathymus neumoegeni florenceae</i> (D. Stallings and Turner, 1957).	265c, (267).
<i>Agathymus neumoegeni judithae</i> (D. Stallings and Turner, 1957).	265d, (268).
<i>Agathymus neumoegeni diaboloensis</i> H. A. Freeman, 1962.	265e, (269).
<i>Agathymus neumeogeni mcalpinei</i> (H. A. Freeman, 1955).	265f, (270).
<i>Agathymus chisosensis</i> (H. A. Freeman), 1952.	271.
<i>Agathymus aryxna</i> (Dyar, 1905).	272.
<i>Agathymus baueri baueri</i> (D. Stallings and Turner, 1954).	273a, (273).
<i>Agathymus baueri fremani</i> (D. Stallings, Turner and J. Stallings, 1960).	273b, (274).
<i>Agathymus evansi</i> (H. A. Freeman, 1950).	275.
<i>Agathymus mariae mariae</i> (Barnes and Benjamin, 1924).	276a, (276).
<i>Agathymus mariae chinatiensis</i> H. A. Freeman, 1964.	276b, (277).
<i>Agathymus mariae lajitaensis</i> H. A. Freeman, 1964.	276c, (278).
<i>Agathymus mariae rindgei</i> H. A. Freeman, 1964.	276d, (279).
<i>Agathymus gilberti</i> H. A. Freeman, 1964.	280.
<i>Agathymus estellae valverdiensis</i> H. A. Freeman, 1966.	281a, (281).
<i>Agathymus stephensi</i> (Skinner, 1912).	282.
<i>Agathymus polingi</i> (Skinner, 1905).	283.
<i>Agathymus alliae</i> (D. Stallings and Turner, 1957).	284.
<i>Megathyimus yuccae yuccae</i> (Boisduval and Le Conte, [1834]).	285a.

<i>Megathymus yuccae buchholzi</i> H. A. Freeman, 1952.	285b.
<i>Megathymus coloradensis coloradensis</i> C. V. Riley, 1877.	286a.
<i>Megathymus coloradensis elidaensis</i> D. Stallings, Turner & J. Stallings, 1966.	286b.
<i>Megathymus coloradensis navajo</i> Skinner, 1911.	286c.
<i>Megathymus coloradensis browni</i> D. Stallings and Turner, 1960.	286d.
<i>Megathymus coloradensis stallingsi</i> H. A. Freeman, 1943.	286e.
<i>Megathymus coloradensis reinthali</i> H. A. Freeman, 1963.	286f.
<i>Megathymus coloradensis martini</i> D. Stallings and Turner, 1956.	286g.
<i>Mehathymus coloradensis maudae</i> D. Stallings, Turner and J. Stallings, 1966.	286h.
<i>Megathymus coloradensis arizonae</i> Tinkham, 1954.	286i.
<i>Megathymus coloradensis albasuffusus</i> R., D., and J. Wielgus, 1974.	286j.
<i>Megathymus coloradensis reubeni</i> D. Stallings, Turner and J. Stallings, 1963.	286k.
<i>Megathymus coloradensis winkensis</i> H. A. Freeman, 1965.	286l.
<i>Megathymus coloradensis wilsonorum</i> D. Stallings and Turner, 1958.	286m.
<i>Megathymus coloradensis louiseae</i> H. A. Freeman, 1963.	286n.
<i>Megathymus coloradensis kendalli</i> H. A. Freeman, 1963.	286o.
<i>Megathymus cofaqui</i> (Strecker, 1876).	287.
<i>Megathymus harrisi</i> H. A. Freeman, 1955.	288.
<i>Megathymus streckeri streckeri</i> (Skinner, 1876).	289a, (289).
<i>Megathymus streckeri texanus</i> Barnes and McDunnough, 1912.	289b, (290a).
<i>Megathymus streckeri leussleri</i> Holland, 1931.	289c, (290b).
<i>Megathymus ursus ursus</i> Poling, 1902.	291a.
<i>Megathymus ursus violae</i> D. Stallings and Turner, 1956.	291b.
<i>Megathymus ursus deserti</i> R. Wielgus, D. Wielgus, and J. Wielgus, 1972.	291c.
<i>Stallingsia maculosa</i> (H. A. Freeman, 1959).	292.

PAPILIONIDAE

<i>Parnassius eversmanni thor</i> Ménétriés, 1849.	293a.
<i>Parnassius clodius clodius</i> Ménétriés, 1855.	294a.
<i>Parnassius clodius strohbeeni</i> Sternitzky, 1945.	294b.
<i>Parnassius clodius sol</i> Bryk and Eisner, 1932.	294c.
<i>Parnassius clodius baldur</i> W. H. Edwards, 1877.	294d.
<i>Parnassius clodius claudianus</i> Stichel, 1907.	294e.
<i>Parnassius clodius pseudogallatinus</i> Bryk, 1913.	294f.
<i>Parnassius clodius incredibilis</i> Bryk, 1932.	294g.
<i>Parnassius clodius altaurus</i> Dyar, 1903.	294h.
<i>Parnassius clodius gallatinus</i> Stichel, 1907.	294i.
<i>Parnassius clodius menetriesi</i> Hy. Edwards, 1877.	294j.
<i>Parnassius clodius sheppardi</i> Eisner, 1966.	294k.
<i>Parnassius phoebus behrii</i> W. H. Edwards, 1870.	295a.
<i>Parnassius phoebus sternitzkyi</i> McDunnough, 1936.	295b.
<i>Parnassius phoebus olympiannus</i> Burdick, 1941.	295c.
<i>Parnassius phoebus smintheus</i> Doubleday, [1847].	295d.
<i>Parnassius phoebus montanus</i> Bryk and Eisner, 1935.	295e.
<i>Parnassius phoebus sayii</i> W. H. Edwards, 1863.	295f.
<i>Parnassius phoebus pseudorotgeri</i> Eisner, 1966.	295g.
<i>Parnassius phoebus apricatus</i> Stichel, 1906.	295h.
<i>Parnassius phoebus golovinus</i> Holland, 1930.	295i.
<i>Parnassius phoebus alaskensis</i> Eisner, 1956.	295j.

<i>Parnassius phoebus elias</i> 1934.	295k.
<i>Parnassius phoebus yukonensis</i> Eisner, 1969.	295l.
<i>Parides eurimedes mylotes</i> (Bates, 1861).	296a.
<i>Battus philenor philenor</i> (Linnaeus, 1771).	297a.
<i>Battus philenor hirsuta</i> (Skinner, 1908).	297b.
<i>Battus polydamas polydamas</i> (Linnaeus, 1758).	298a.
<i>Battus polydamas lucayus</i> Rothschild and Jordan, 1906.	298b.
<i>Battus devilliers</i> (Godart, [1824]).	299.
<i>Eurytides marcellus</i> (Cramer, [1777]).	300.
<i>Eurytides philolaus</i> (Boisduval, 1836).	301.
<i>Eurytides celadon</i> (Lucas, 1852).	302.
<i>Papilio polyxenes asterius</i> Stoll, 1775.	303a.
<i>Papilio polyxenes coloro</i> W. G. Wright, 1905.	303b, (305).
<i>Papilio joanae</i> J. R. Heitzman, [1974].	304.
<i>Papilio kahli</i> F. and R. Chermock, 1937.	306.
<i>Papilio brevicauda brevicauda</i> Saunders, 1869.	307a.
<i>Papilio brevicauda gaspeensis</i> McDunnough, 1934.	307b.
<i>Papilio brevicauda bretonensis</i> McDunnough, 1939.	307c.
<i>Papilio bairdii bairdii</i> W. H. Edwards, 1869.	308a.
<i>Papilio bairdii oregonius</i> W. H. Edwards, 1876.	308b, (309a).
<i>Papilio bairdii dodi</i> McDunnough, 1939.	308c, (309b).
<i>Papilio machaon aliaska</i> Scudder, 1869.	310a.
<i>Papilio machaon hudsonianus</i> A. H. Clark, 1932.	310b.
<i>Papilio zelicaon zelicaon</i> Lucas, 1852.	311a.
<i>Papilio zelicaon nitra</i> W. H. Edwards, 1883.	311b.
<i>Papilio indra indra</i> Reakirt, 1866.	312a.
<i>Papilio indra minori</i> Cross, 1937.	312b.
<i>Papilio indra pergamus</i> Hy. Edwards, 1875.	312c.
<i>Papilio indra kaibabensis</i> Bauer, 1955.	312d.
<i>Papilio indra fordii</i> J. A. Comstock and Martin, 1955.	312e.
<i>Papilio indra martini</i> J. and T. Emmel, 1966.	312f.
<i>Papilio indra nevadensis</i> T. and J. Emmel, 1971.	312g.
<i>Papilio indra phyllisae</i> J. Emmel, 1981.	312h.
<i>Papilio indra panamintensis</i> J. Emmel, 1981.	312i.
<i>Heraclides thoas autocles</i> (Rothschild and Jordan, 1906).	313a.
<i>Heraclides thoas oviedo</i> (Gundlach, 1866).	313b.
<i>Heraclides cresaiphontes</i> (Cramer, [1777]).	314.
<i>Heraclides aristodemus ponceanus</i> (Schaus, 1911).	315a.
<i>Heraclides andraemon bonhotei</i> (E. M. Sharpe, [1900]).	316a.
<i>Heraclides ornythion</i> (Boisduval, 1836).	317.
<i>Heraclides astyalus pallas</i> (G. R. Grey, 1852).	318a.
<i>Heraclides androgeus epidaurus</i> (Godman and Salvin, 1890).	319a.
<i>Heraclides anchisiades idaeus</i> (Fabricius, 1793).	319.1a, (327a).
<i>Pterourus glaucus glaucus</i> (Linnaeus, 1758).	320a.
<i>Pterourus glaucus canadensis</i> (Rothschild and Jordan, 1906).	320b.
<i>Pterourus glaucus maynardi</i> (Gauthier, 1984).	320c.
<i>Pterourus glaucus arcticus</i> (Skinner, 1906).	320d.
<i>Pterourus rutulus rutulus</i> (Lucas), 1852.	321a.
<i>Pterourus rutulus arizonensis</i> (W. H. Edwards, 1883).	321b, (321c).
<i>Pterourus multicaudatus</i> (W. F. Kirby, 1884).	322.

<i>Pterourus eurymedon</i> (Lucas, 1852).	323.
<i>Pterourus pilumnus</i> (Boisduval, 1836).	324.
<i>Pterourus troilus troilus</i> (Linnaeus, 1758).	325a.
<i>Pterourus troilus ilioneus</i> (J. E. Smith, 1797).	325b.
<i>Pterourus palamedes</i> (Drury, [1773]).	326.
<i>Pterourus victorinus</i> (Doubleday, 1844).	326.1.

PIERIDAE

<i>Catasticta nimbice nimbice</i> (Boisduval, 1836).	328a.
<i>Neophasia menapia menapia</i> (C. and R. Felder, 1859).	329a.
<i>Neophasia menapia tau</i> (Scudder, 1861).	329b.
<i>Neophasia menapia melanica</i> Scott, 1981.	329c.
<i>Neophasia terlootii</i> Behr, 1869.	330.
<i>Appias drusilla poeyi</i> Butler, [1872].	331a.
<i>Appias drusilla neumoegeni</i> (Skinner, 1894).	331b.
<i>Pontia beckerii</i> (W. H. Edwards, 1871).	332.
<i>Pontia sisymbrii sisymbrii</i> (Boisduval, 1852).	333a.
<i>Pontia sisymbrii flavitincta</i> (J. A. Comstock, 1924).	333b.
<i>Pontia sisymbrii elivata</i> (Barnes and Benjamin, 1926).	333c.
<i>Pontia sisymbrii nordini</i> (K. Johnson, 1977).	333d.
<i>Pontia protodice</i> (Boisduval and Le Conte, 1829).	334.
<i>Pontia occidentalis occidentalis</i> (Reakirt, 1866).	335a.
<i>Pontia occidentalis nelsoni</i> (W. H. Edwards, 1883).	335b.
<i>Pieris napi pseudobryoniae</i> Verity, 1908.	336a.
<i>Pieris napi hulda</i> W. H. Edwards, 1869.	336b.
<i>Pieris napi frigida</i> Scudder, 1861.	336c.
<i>Pieris napi oleracea</i> Harris, 1929.	336d.
<i>Pieris napi venosa</i> Scudder, 1861.	336e.
<i>Pieris napi castoria</i> Reakirt, 1866.	336f.
<i>Pieris napi marginalis</i> Scudder, 1861.	336g.
<i>Pieris napi mcdunnoughi</i> Remington, 1954.	336h.
<i>Pieris napi mogollon</i> Burdick, 1942.	336i.
<i>Pieris virginensis</i> W. H. Edwards, 1870.	337.
<i>Pieris rapae</i> (Linnaeus, 1758).	338.
<i>Ascia monuste monuste</i> (Linnaeus, 1764).	339a.
<i>Ascia monuste phileta</i> (Fabricius, 1775).	339b.
<i>Ascia monuste cleomes</i> (Boisduval and Le Conte, [1829]).	339c.
<i>Ganyra josephina josepha</i> (Salvin and Godman, 1868).	340a.
<i>Euchloe ausonides ausonides</i> (Lucas, 1852).	341a.
<i>Euchloe ausonides coloradensis</i> (Hy. Edwards, 1881).	341b.
<i>Euchloe ausonides mayi</i> F. and R. Chermock, 1940.	341c.
<i>Euchloe ausonides palaeoreios</i> K. Johnson, 1976.	341d.
<i>Euchloe creusa</i> (Doubleday, [1847]).	342.
<i>Euchloe hyantis hyantis</i> (W. H. Edwards, 1871).	343a.
<i>Euchloe hyantis andrewsi</i> Martin, 1958.	343b.
<i>Euchloe hyantis lotta</i> Beutenmüller, 1898.	343c.
<i>Euchloe olympia</i> W. H. Edwards, 1871.	344.
<i>Anthocharis cethura cethura</i> C. and R. Felder, [1865].	345a.
<i>Anthocharis cethura catalina</i> Meadows, 1937.	345b.

<i>Anthocharis pima</i> W. H. Edwards, 1888.	346.
<i>Anthocharis dammersi</i> J. A. Comstock, 1929.	347.
<i>Anthocharis sara sara</i> Lucas, 1852.	348a.
<i>Anthocharis sara inghami</i> Gunder, 1932.	348b.
<i>Anthocharis sara thoosa</i> (Scudder, 1878).	348c.
<i>Anthocharis sara julia</i> W. H. Edwards, 1872.	348d.
<i>Anthocharis sara browningi</i> Skinner, 1906.	348e.
<i>Anthocharis sara stella</i> W. H. Edwards, 1879.	348f.
<i>Anthocharis sara flora</i> W. G. Wright, 1905.	348g.
<i>Anthocharis sara alaskensis</i> Gunder, 1932.	348h.
<i>Paramidea midea midea</i> (Hübner, [1809]).	349a.
<i>Paramidea midea annickae</i> (dos Passos and Klots, 1969).	349b.
<i>Paramidea lanceolata</i> (Lucas, 1852).	350.
<i>Colias philodice philodice</i> Godart, [1819].	351a.
<i>Colias philodice vitabunda</i> Hovanitz, 1943.	351b
<i>Colias philodice eriphyle</i> W. H. Edwards, 1876.	351c.
<i>Colias eurytheme</i> Boisduval, 1852.	352.
<i>Colias occidentalis occidentalis</i> Scudder, 1862.	354a.
<i>Colias occidentalis chrysomelas</i> Hy. Edwards, 1877.	354b.
<i>Colias alexandra alexandra</i> W. H. Edwards, 1863.	355a.
<i>Colias alexandra edwardsii</i> W. H. Edwards, 1870.	355b.
<i>Colias alexandra columbiensis</i> Ferris, 1973.	355c.
<i>Colias alexandra astraea</i> W. H. Edwards, 1872.	355d.
<i>Colias alexandra christina</i> W. H. Edwards, 1863.	355e.
<i>Colias alexandra krauthii</i> Klots, 1935.	355f.
<i>Colias alexandra kluanensis</i> Ferris, 1981.	355g.
<i>Colias alexandra apache</i> Ferris, 1988	355h.
<i>Colias alexandra harfordii</i> Hy. Edwards, 1877.	355i, (353).
<i>Colias meadii meadii</i> W. H. Edwards, 1871.	356a.
<i>Colias meadii elis</i> Strecker, 1885.	356b.
<i>Colias meadii lemhiensis</i> Curtis and Ferris, 1985.	356c.
<i>Colias hecla hecla</i> Lefèvre, 1836.	357a.
<i>Colias hecla hela</i> Strecker, 1880.	357b.
<i>Colias canadensis</i> Ferris, 1982.	357.1.
<i>Colias boothii</i> Curtis, 1835.	[see comment in Part II]
<i>Colias nastes nastes</i> Boisduval, "1832" [1834].	358.
<i>Colias nastes moina</i> Strecker, 1880.	360a.
<i>Colias nastes aliaska</i> Bang-Haas, 1927.	360b.
<i>Colias nastes streckeri</i> Grum-Grischimailo, 1895.	360c.
<i>Colias scudderii</i> Reakirt, 1865.	360d.
<i>Colias gigantea gigantea</i> Strecker, 1900.	361.
<i>Colias gigantea harroweri</i> Klots, 1940.	362a.
<i>Colias pelidne pelidne</i> Boisduval and Le Conte, 1829.	362b.
<i>Colias pelidne skinneri</i> Barnes, 1897.	363a.
<i>Colias interior</i> Scudder, 1862.	363b.
<i>Colias palaeno chippewa</i> W. H. Edwards, 1872.	364.
<i>Colias palaeno baffinensis</i> Ebner and Ferris, 1977.	365a.
<i>Colias behrii</i> W. H. Edwards, 1866.	365b.
<i>Zerene eurydice</i> (Boisduval, 1855).	366.
<i>Zerene cesonia cesonia</i> (Stoll), [1790].	367.
	368a.

<i>Anteos clorinde nivifera</i> (Fruhstofer), 1907.	369a.
<i>Anteos maerula</i> (Fabricius), 1775.	370.
<i>Phoebis sennae sennae</i> (Linnaeus, 1758).	371a.
<i>Phoebis sennae eubule</i> (Linnaeus, 1767).	371b.
<i>Phoebis sennae marcellina</i> (Cramer, [1779]).	371c.
<i>Phoebis philea philea</i> (Linnaeus in Johansson, 1763).	372a.
<i>Phoebis argante argante</i> (Fabricius, 1775).	373a.
<i>Phoebis agarithe agarithe</i> (Boisduval, 1836).	374a.
<i>Phoebis agarithe maxima</i> (Neumoegen, 1891).	374b.
<i>Phoebis neocypris</i> (Hübner, [1823]).	375.
<i>Aphrissa statira jada</i> (Butler, [1870]).	376a.
<i>Aphrissa statira floridensis</i> (Neumoegen, 1891).	376b.
<i>Aphrissa orbis</i> (Poey, 1832).	376.1.
<i>Kricogonia lyside</i> (Godart, [1819]).	377.
<i>Eurema daira daira</i> (Godart, [1819]).	378a.
<i>Eurema daira palmira</i> (Poey, 1853).	378b.
<i>Eurema daira lydia</i> (C. and R. Felder, 1861).	378c.
<i>Eurema boisduvalianum</i> (C. and R. Felder, [1865]).	379.
<i>Eurema mexicanum</i> (Boisduval, 1836).	380.
<i>Eurema salome limoneum</i> (C. and R. Felder, 1861).	381a.
<i>Eurema proterpia</i> (Fabricius, 1775).	382.
<i>Eurema lisa lisa</i> (Boisduval and Le Conte, 1829).	383a.
<i>Eurema chamberlaini</i> (Butler, 1897).	384.
<i>Eurema nise nise</i> (Cramer, [1775]).	385a.
<i>Eurema nise nelphe</i> (R. Felder), 1869.	385b.
<i>Eurema messalina blakei</i> (Maynard, 1891).	386a.
<i>Eurema dina westwoodi</i> (Boisduval, 1836).	387a.
<i>Eurema dina helios</i> M. Bates, 1934.	387b.
<i>Eurema nicippe</i> (Cramer, [1779]).	388.
<i>Nathalis iole</i> Boisduval, 1836.	389.
<i>Enantia albania</i> (Bates, 1864).	390.

LYCAENIDAE

<i>Feniseca tarquinius tarquinius</i> (Fabricius, 1793).	391a.
<i>Feniseca tarquinius novascotiae</i> McDunnough, 1935.	391b.
<i>Tharsalea arota arota</i> (Boisduval, 1852).	392a.
<i>Tharsalea arota virginicensis</i> (W. H. Edwards, 1870).	392b.
<i>Tharsalea arota nubila</i> J. A. Comstock, 1926.	392c.
<i>Tharsalea arota schellbachi</i> Tilden, 1955.	392d.
<i>Lycaena phlaeas americana</i> Harris, 1862.	393a.
<i>Lycaena phlaeas feildeni</i> (M'Lachlan, 1878).	393b.
<i>Lycaena phlaeas arethusa</i> (Wolley-Dod, 1907).	393c.
<i>Lycaena phlaeas arctodon</i> Ferris, 1974.	393d.
<i>Lycaena phlaeas hypophlaeas</i> (Boisduval), 1852	393e.
<i>Lycaena cuprea cuprea</i> (W. H. Edwards, 1870).	394a.
<i>Lycaena cuprea snowi</i> (W. H. Edwards, [1881]).	394b.
<i>Lycaena cuprea henryae</i> (Cadbury, 1937).	394c.
<i>Lycaena cuprea artemisia</i> Scott, 1981.	394d.
<i>Gaeides xanthoides xanthoides</i> (Boisduval, 1852).	395a.

<i>Gaeides xanthoides dione</i> (Scudder, 1869).	395b.
<i>Gaeides editha editha</i> (Mead, 1878).	396a.
<i>Gaeides editha montana</i> (Field, 1936).	396b.
<i>Gaeides editha nevadensis</i> (Austin, 1984).	396c.
<i>Gaeides gorgon</i> (Boisduval, 1852).	397.
<i>Hyllolycaena hyllus</i> (Cramer, [1775]).	398.
<i>Chalceria rubida rubida</i> (Behr, 1866).	399a.
<i>Chalceria rubida duofacies</i> (K. Johnson and Balogh, 1977).	399b.
<i>Chalceria rubida perkinsorum</i> (K. Johnson and Balogh, 1977).	399c.
<i>Chalceria rubida longi</i> (K. Johnson and Balogh, 1977).	399d.
<i>Chalceria rubida sirius</i> (W. H. Edwards, 1871).	399e.
<i>Chalceria rubida monachensis</i> (K. Johnson and Balogh, 1977).	399f.
<i>Chalceria ferrisi</i> (K. Johnson and Balogh, 1977).	400.
<i>Chalceria heteronea heteronea</i> (Boisduval, 1852).	401a.
<i>Chalceria heteronea clara</i> (Hy. Edwards, 1880).	401b.
<i>Epidemia epixanthe epixanthe</i> (Boisduval and Le Conte, [1833]).	402a.
<i>Epidemia epixanthe phaedra</i> (G. C. Hall, 1924).	402b.
<i>Epidemia epixanthe michiganensis</i> (Rawson, 1948).	402c.
<i>Epidemia dorcas dorcas</i> (W. Kirby, 1837).	403a.
<i>Epidemia dorcas castro</i> (Reakirt, 1866).	403b.
<i>Epidemia dorcas florus</i> (W. H. Edwards, 1883).	403c.
<i>Epidemia dorcas dospassosi</i> (McDunnough, 1940).	403d.
<i>Epidemia dorcas claytoni</i> (Brower, 1940).	403e.
<i>Epidemia dorcas megaloceras</i> Ferris, 1977.	403f.
<i>Epidemia dorcas arcticus</i> Ferris, 1977.	403g.
<i>Epidemia helloides</i> (Boisduval, 1852).	404.
<i>Epidemia nivalis nivalis</i> (Boisduval, 1869).	405a.
<i>Epidemia nivalis browni</i> (dos Passos, 1938).	405b.
<i>Epidemia mariposa mariposa</i> (Reakirt, 1866).	406a.
<i>Epidemia mariposa charlottensis</i> (Holland, 1930).	406b.
<i>Epidemia mariposa penroseae</i> (Field, 1938).	406c.
<i>Hermelycaena hermes</i> (W. H. Edwards, 1870).	407.
<i>Hypaurotis crysalus crysalus</i> (W. H. Edwards, 1873).	408a.
<i>Hypaurotis crysalus citima</i> (Hy. Edwards, 1881).	408b.
<i>Habrodais grunus grunus</i> (Boisduval, 1852).	409a.
<i>Habrodais grunus lorquini</i> Field, 1938.	409b.
<i>Habrodais grunus herri</i> Field, 1938.	409c.
<i>Eumaeus atala florida</i> Röber, 1926.	410a.
<i>Eumaeus toxea</i> (Godart, [1824]).	[see comment in Part II]
<i>Atides halesus halesus</i> (Cramer, [1777]).	411.
<i>Atides halesus estesi</i> Clench, 1940.	412a.
<i>Chlorostrymon maesites</i> (Herrich-Schäffer, 1864).	412b.
<i>Chlorostrymon telea</i> (Hewitson, 1868).	413.
<i>Chlorostrymon simaethis simaethis</i> (Drury, 1770).	414.
<i>Chlorostrymon simaethis sarita</i> (Skinner, 1895).	415a.
<i>Phaeostrymon alcestis alcestis</i> (W. H. Edwards, 1871).	415b.
<i>Phaeostrymon alcestis oslari</i> (Dyar, 1904).	416a.
<i>Harkenclenus titus titus</i> (Fabricius, 1793).	416b.
<i>Harkenclenus titus mopsus</i> (Hübner, 1818).	417a.
<i>Harkenclenus titus watsoni</i> (Barnes and Benjamin, 1926).	417b.
	417c.

<i>Harkenclenus titus immaculosus</i> (W. P. Comstock, 1913).	417d.
<i>Satyrium behrii behrii</i> (W. H. Edwards, 1870).	418a.
<i>Satyrium behrii crossi</i> (Field, 1938).	418b.
<i>Satyrium behrii columbia</i> (McDunnough, 1944).	418c.
<i>Satyrium fuliginosum fuliginosum</i> (W. H. Edwards, 1861).	419a.
<i>Satyrium fuliginosum semiluna</i> Klots, 1930.	419b.
<i>Satyrium acadicum acadicum</i> (W. H. Edwards, 1862).	420a.
<i>Satyrium acadicum coolinensis</i> (Watson and W. P. Comstock, 1920).	420b.
<i>Satyrium acadicum montanensis</i> (Watson and W. P. Comstock, 1920).	420c.
<i>Satyrium acadicum watrini</i> (Dufrane, 1939).	420d.
<i>Satyrium californicum</i> (W. H. Edwards, 1862).	421.
<i>Satyrium sylvium sylvium</i> (Boisduval, 1852).	422a.
<i>Satyrium sylvium dryope</i> (W. H. Edwards, 1870).	422b.
<i>Satyrium sylvium itys</i> (W. H. Edwards, 1882).	422c.
<i>Satyrium sylvium desertorum</i> (Grinnell, 1917).	422d.
<i>Satyrium sylvium putnami</i> (Hy. Edwards, 1877).	422e.
<i>Satyrium edwardsii</i> (Grote and Robinson, 1867).	423.
<i>Satyrium calanus calanus</i> (Hübner, [1809]).	424a.
<i>Satyrium calanus falacer</i> (Godart, [1824]).	424b.
<i>Satyrium calanus godarti</i> (Field, 1938).	424c.
<i>Satyrium calanus albidus</i> Scott, 1981.	424d.
<i>Satyrium caryaevorum</i> (McDunnough, 1942).	425.
<i>Satyrium kingi</i> (Klots and Clench, 1952).	426.
<i>Satyrium liparops liparops</i> (Le Conte, 1833).	427a.
<i>Satyrium liparops strigosum</i> (Harris, 1862).	427b.
<i>Satyrium liparops fletcheri</i> (Michener and dos Passos, 1942).	427c.
<i>Satyrium liparops aliparops</i> (Michener and dos Passos, 1942).	427d.
<i>Satyrium auretorum auretorum</i> (Boisduval, 1852).	428a.
<i>Satyrium auretorum spadix</i> (Hy. Edwards, 1881).	428b.
<i>Satyrium tetra</i> (W. H. Edwards, 1870).	429.
<i>Satyrium saepium saepium</i> (Boisduval, 1852).	430a.
<i>Satyrium saepium okanaganum</i> (McDunnough, 1944).	430b.
<i>Ocaria ocrisia</i> (Hewitson, 1868).	[see comment in Part II]
<i>Ministrymon clytie</i> (W. H. Edwards, 1877).	431.
<i>Ministrymon leda</i> (W. H. Edwards), 1882.	432.
<i>Tmolus echion echiolus</i> (Draudt, 1920).	433.
<i>Tmolus azia</i> (Hewitson, 1873).	434a.
<i>Oenomaus ortygynus</i> (Cramer, [1779]).	435.
<i>Thereus zebina</i> (Hewitson, 1869).	436.
<i>Thereus spurina</i> (Hewitson, 1867).	[see comment in Part II]
<i>Thereus palegon</i> (Stoll, [1780]).	[see comment in Part II]
<i>Allosmaitia pion</i> (Godman and Salvin, 1887).	[see comment in Part II]
<i>Calycopis cecrops</i> (Fabricius, 1793).	440.
<i>Calycopis isobeon</i> (Butler and H. Druce, 1872).	441.
<i>Callophrys dumetorum dumetorum</i> (Boisduval, 1852).	442.
<i>Callophrys dumetorum perplexa</i> Barnes and Benjamin, 1923. [see comment in Part II]	443a.
<i>Callophrys dumetorum oregonensis</i> Gorelick, 1969.	443b.
<i>Callophrys comstocki</i> Henne, 1940.	443c.
<i>Callophrys lemberti</i> Tilden, 1963.	444.
<i>Callophrys apama apama</i> (W. H. Edwards, 1882).	445.
	446a.

<i>Callophrys apama homoperplexa</i> Barnes and Benjamin, 1923.	446b.
<i>Callophrys affinis affinis</i> (W. H. Edwards, 1862).	447a.
<i>Callophrys affinis washingtonia</i> Clench, 1944.	447b.
<i>Callophrys viridis</i> (W. H. Edwards, 1862). [see comment in Part II]	448.
<i>Callophrys sheridanii sheridanii</i> (W. H. Edwards, 1877).	449a.
<i>Callophrys sheridanii neoperplexa</i> Barnes and Benjamin, 1923.	449b.
<i>Callophrys sheridanii newcomeri</i> Clench, 1963.	449c.
<i>Cyanophrys miserabilis</i> (Clench, 1946).	450.
<i>Cyanophrys goodsoni</i> (Clench, 1946).	451.
<i>Mitoura spinetorum spinetorum</i> (Hewitson, 1867).	452a.
<i>Mitoura spinetorum ninus</i> (W. H. Edwards, 1871).	452b.
<i>Mitoura millerorum</i> (Clench, 1981).	452.1.
<i>Mitoura johnsoni</i> (Skinner, 1904).	453.
<i>Mitoura rosneri rosneri</i> K. Johnson, 1976.	454a.
<i>Mitoura rosneri plicataria</i> K. Johnson, 1976.	454b.
<i>Mitoura barryi barryi</i> K. Johnson, 1976.	455a.
<i>Mitoura barryi acuminata</i> K. Johnson, 1976.	455b.
<i>Mitoura byrnei</i> K. Johnson, 1976.	456.
<i>Mitoura nelsoni nelsoni</i> (Boisduval), 1869.	457a.
<i>Mitoura nelsoni muiri</i> (Hy. Edwards), 1881.	457b.
<i>Mitoura siva siva</i> (W. H. Edwards, 1874).	458a.
<i>Mitoura siva juniperaria</i> J. A. Comstock, 1925.	458b.
<i>Mitoura siva mansfieldi</i> Tilden, 1951.	458c.
<i>Mitoura siva chalcosiva</i> (Clench, 1981).	458d.
<i>Mitoura siva clenchi</i> K. Johnson, 1988.	458e.
<i>Mitoura loki</i> (Skinner, 1907).	459.
<i>Mitoura grynea grynea</i> (Hübner, [1819]).	460a.
<i>Mitoura grynea smilacis</i> (Boisduval and Le Conte, [1833]).	460b.
<i>Mitoura grynea sweadneri</i> F. H. Chermock, 1940.	460c.
<i>Mitoura grynea castalis</i> (W. H. Edwards, 1871).	460d.
<i>Mitoura hesseli</i> Rawson and Ziegler, 1950.	461.
<i>Mitoura thornei</i> J. W. Brown, [1983].	461.1.
<i>Xamia xami xami</i> (Reakirt, 1866).	462a.
<i>Xamia xami texami</i> (Clench, 1981).	462b.
<i>Sandia mcfarlandi mcfarlandi</i> P. Ehrlich and Clench, 1960.	463a.
<i>Incisalia augustinus augustinus</i> (W. Kirby, 1873).	464a.
<i>Incisalia augustinus helena</i> dos Passos, 1943.	464b.
<i>Incisalia augustinus croesioides</i> Scudder, 1876.	464c.
<i>Incisalia augustinus iroides</i> (Boisduval, 1852).	464d.
<i>Incisalia augustinus annetteae</i> dos Passos, 1943.	464e.
<i>Incisalia fotis</i> (Strecker, 1878).	465.
<i>Incisalia mossii mossii</i> (Hy. Edwards, 1881).	465.1a.
<i>Incisalia mossii schryveri</i> Cross, 1937.	465.1b.
<i>Incisalia mossii bayensis</i> (R. M. Brown, 1969).	465.1c.
<i>Incisalia mossii doudoroffi</i> dos Passos, 1940.	465.1d.
<i>Incisalia mossii windi</i> Clench, 1943.	465e.
<i>Incisalia polia polia</i> Cook and Watson, 1907.	466a.
<i>Incisalia polia obscura</i> Ferris and Fisher, 1973.	466b.
<i>Incisalia irus irus</i> (Godart, [1824]).	467a.
<i>Incisalia irus arsace</i> (Boisduval and Le Conte, [1833]).	467b.

<i>Incisalia irus hadra</i> Cook and Watson, 1909.	467c.
<i>Incisalia henrici henrici</i> (Grote and Robinson, 1867).	468a.
<i>Incisalia henrici margaretae</i> dos Passos, 1943.	468b.
<i>Incisalia henrici solata</i> Cook and Watson, 1909.	468c.
<i>Incisalia henrici turneri</i> Clench, 1943.	468d.
<i>Incisalia lanoriaensis</i> Sheppard, 1934.	469.
<i>Incisalia niphon niphon</i> (Hübner, [1823]).	470a.
<i>Incisalia niphon clarki</i> T. N. Freeman, 1938.	470b.
<i>Incisalia eryphon eryphon</i> (Boisduval, 1852).	471a.
<i>Incisalia eryphon sheltonensis</i> F. Chermock and Frechin, 1948.	471b.
<i>Arawacus jada</i> (Hewitson), [1867].	472.
<i>Fixsenia favonius</i> (J. E. Smith, 1797).	473.
<i>Fixsenia ontario ontario</i> (W. H. Edwards, 1868).	474a.
<i>Fixsenia ontario autolycus</i> (W. H. Edwards, 1868).	474b.
<i>Fixsenia ontario violae</i> (D. Stallings and Turner, 1947).	474c.
<i>Fixsenia ontario ilavia</i> (Beutenmüller, 1899).	474d.
<i>Fixsenia polingi polingi</i> (Barnes and Benjamin, 1926).	475a.
<i>Fixsenia polingi organensis</i> Ferris, 1980.	475b.
<i>Hypostrymon critola</i> (Hewitson, 1874).	476.
<i>Parrhasius m-album</i> (Boisduval and Le Conte, [1833]).	477.
<i>Strymon melinus melinus</i> Hübner, [1818].	478a.
<i>Strymon melinus humuli</i> (Harris, 1841).	478b.
<i>Strymon melinus franki</i> Field, 1938.	478c.
<i>Strymon melinus pudicus</i> (Hy. Edwards, 1876).	478d.
<i>Strymon melinus setonia</i> McDunnough, 1927.	478e.
<i>Strymon melinus atrofasciatus</i> McDunnough, 1921.	478f.
<i>Strymon avalona</i> (W. G. Wright, 1905).	479.
<i>Strymon rufofuscus</i> (Hewitson, 1877).	480.
<i>Strymon bebrycia</i> (Hewitson, 1868).	481.
<i>Strymon martialis</i> (Herrich-Schäffer, 1864).	482.
<i>Strymon yojoa</i> (Reakirt, [1866]).	483.
<i>Strymon albatus sedecia</i> (Hewitson, 1874).	484a.
<i>Strymon acis bartrami</i> (W. P. Comstock and Huntington, 1943).	485a.
<i>Strymon alea</i> (Godman and Salvin, 1887).	486.
<i>Strymon columella modesta</i> (Maynard, 1873).	487a.
<i>Strymon columella cybira</i> (Hewitson, 1874).	487b.
<i>Strymon columella istapa</i> (Reakirt, 1866).	487c.
<i>Strymon limenia</i> (Hewitson, 1868).	488.
<i>Strymon cestri</i> (Reakirt, [1866]).	489.
<i>Strymon bazochii</i> (Godart, [1824]).	[see comment in Part II]
<i>Erora laeta</i> (W. H. Edwards, 1862).	490.
<i>Erora quaderna sanfordi</i> dos Passos, 1940.	491.
<i>Electrostrymon endymion cyphara</i> (Hewitson, 1874). [see comment in Part II]	492a.
<i>Electrostrymon angelia</i> (Hewitson, 1874).	493a.
<i>Brephidium exile</i> (Boisduval, 1852).	494.
<i>Brephidium isophthalma pseudofea</i> (Morrison, 1873).	495.
<i>Leptotes cassius theonus</i> (Lucas, 1857).	496a.
<i>Leptotes cassius striata</i> W. H. Edwards, 1878).	497a.
<i>Leptotes marina</i> (Reakirt, 1868).	497b.
<i>Zizula cyna</i> (W. H. Edwards, 1881).	498.

<i>Hemiargus thomasi bethunebakeri</i> W. P. Comstock and Huntington, 1943.	500a.
<i>Hemiargus ceraunus antibubastus</i> Hübner, [1818].	501a.
<i>Hemiargus ceraunus gyas</i> (W. H. Edwards, 1871).	501b.
<i>Hemiargus ceraunus zachaenia</i> (Butler and H. Druce, 1872).	501c.
<i>Hemiargus isolata alcea</i> (W. H. Edwards, 1871).	502a.
<i>Everes comyntas comyntas</i> (Godart, [1824]).	503a.
<i>Everes comyntas texanus</i> F. Chermock, 1944.	503b.
<i>Everes amyntula amyntula</i> (Boisduval, 1852).	504a.
<i>Everes amyntula valeriae</i> Clench, 1944.	504b.
<i>Everes amyntula albrighti</i> Clench, 1944.	504c.
<i>Everes amyntula herrii</i> F. Grinnell, 1901.	504d.
<i>Celastrina argiolus ladon</i> (Cramer, [1780]).	505a.
<i>Celastrina argiolus lucia</i> (W. Kirby, 1837).	505b.
<i>Celastrina argiolus argentata</i> (Fletcher, 1903).	505c.
<i>Celastrina argiolus nigrescens</i> (Fletcher, 1903).	505d.
<i>Celastrina argiolus sidara</i> (Clench, 1944).	505e.
<i>Celastrina argiolus echo</i> (W. H. Edwards, 1864).	505f.
<i>Celastrina argiolus gozora</i> (Boisduval, 1870).	505g.
<i>Celastrina argiolus cinerea</i> (W. H. Edwards, 1883).	505h.
<i>Celastrina ebenina</i> Clench, 1972.	506.
<i>Philotes sonorensis</i> (C. and R. Felder, 1865).	507.
<i>Euphilotes battoides centralis</i> (Barnes and McDunnough, 1917).	508a.
<i>Euphilotes battoides ellisi</i> (Shields, 1975).	508b.
<i>Euphilotes battoides glaucon</i> (W. H. Edwards, 1871).	508c.
<i>Euphilotes battoides oregonensis</i> (Barnes and McDunnough, 1917).	508d.
<i>Euphilotes battoides baueri</i> (Shields, 1975).	508e.
<i>Euphilotes battoides intermedia</i> (Barnes and McDunnough, 1917).	508f.
<i>Euphilotes battoides battoides</i> (Behr, 1867).	508g.
<i>Euphilotes battoides martini</i> (Mattoni, [1955]).	508h.
<i>Euphilotes battoides bernardino</i> (Barnes and McDunnough, 1917).	508i.
<i>Euphilotes battoides comstocki</i> (Shields, 1975).	508j.
<i>Euphilotes battoides allyni</i> (Shields, 1975).	508k.
<i>Euphilotes enoptes ancilla</i> (Barnes and McDunnough, 1918).	509a.
<i>Euphilotes enoptes columbiae</i> (Mattoni, [1955]).	509b.
<i>Euphilotes enoptes enoptes</i> (Boisduval, 1852).	509c, (509d).
<i>Euphilotes enoptes bayensis</i> (Langston, [1964]).	509d, (509e).
<i>Euphilotes enoptes smithi</i> (Mattoni, [1955]).	509e, (509f).
<i>Euphilotes enoptes tildeni</i> (Langston, [1964]).	509f, (509g).
<i>Euphilotes enoptes langstoni</i> (Shields, 1975).	509g, (509h).
<i>Euphilotes enoptes dammersi</i> (J. A. Comstock and Henne, 1933).	509h, (509i).
<i>Euphilotes mojave</i> (Watson and W. P. Comstock, 1920).	510, (509c).
<i>Euphilotes rita rita</i> (Barnes and McDunnough, 1916).	511a.
<i>Euphilotes rita coloradensis</i> (Mattoni, [1966]).	511b.
<i>Euphilotes rita emmeli</i> (Shields, 1975).	511c, (511d).
<i>Euphilotes rita mattonii</i> (Shields, 1975).	511d, (511e).
<i>Euphilotes rita pallescens</i> (Tilden and Downey, 1955).	511e, (510a).
<i>Euphilotes rita elvira</i> (Mattoni, [1966]).	511f, (510b).
<i>Euphilotes spaldingi spaldingi</i> (Barnes and McDunnough, 1917).	511.1a, (511c).
<i>Euphilotes spaldingi pinjuna</i> Scott, 1981.	511.1b.
<i>Philotiella speciosa</i> (Hy. Edwards, 1876).	512.

<i>Glaucoptysche piassus piassus</i> (Boisduval, 1852).	513a.
<i>Glaucoptysche piassus sagittigera</i> (C. and R. Felder, 1865).	513b.
<i>Glaucoptysche piassus nevada</i> F. M. Brown, 1975.	513c.
<i>Glaucoptysche piassus daunia</i> (W. H. Edwards, 1871).	513d.
<i>Glaucoptysche piassus toxœuma</i> F. M. Brown, 1971.	513e.
<i>Glaucoptysche lygdamus mildredae</i> F. H. Chermock, 1944.	514a.
<i>Glaucoptysche lygdamus couperi</i> Grote, 1873.	514b.
<i>Glaucoptysche lygdamus afra</i> (W. H. Edwards, [1884]).	514c.
<i>Glaucoptysche lygdamus lygdamus</i> (Doubleday, 1841).	514d.
<i>Glaucoptysche lygdamus jacki</i> D. Stallings and Turner, 1947.	514e.
<i>Glaucoptysche lygdamus oro</i> (Scudder, 1876).	514f.
<i>Glaucoptysche lygdamus arizonensis</i> McDunnough, 1934.	514g.
<i>Glaucoptysche lygdamus australis</i> F. Grinnell, 1917.	514h.
<i>Glaucoptysche lygdamus incognita</i> Tilden, [1974].	514i.
<i>Glaucoptysche lygdamus palosverdescensis</i> E. M. Perkins and J. Emmel, 1978.	514j.
<i>Glaucoptysche lygdamus columbia</i> (Skinner, 1917).	514k.
<i>Glaucoptysche xerxes</i> (Boisduval, 1852).	515.
<i>Lycaeides idas anna</i> (W. H. Edwards), 1861.	516a.
<i>Lycaeides idas ricei</i> (Cross, 1937).	516b.
<i>Lycaeides idas lotis</i> (Lintner, [1876]).	516c.
<i>Lycaeides idas alaskensis</i> (F. H. Chermock, [1945]).	516d.
<i>Lycaeides idas scudderri</i> (W. H. Edwards, 1861).	516e.
<i>Lycaeides idas aster</i> (W. H. Edwards, 1882).	516f.
<i>Lycaeides idas empetri</i> (T. N. Freeman, 1938).	516g.
<i>Lycaeides idas ferniensis</i> (F. H. Chermock, [1945]).	516h.
<i>Lycaeides idas atrapraetextus</i> (Field, 1939).	516i.
<i>Lycaeides idas sublivens</i> Nabokov, 1949.	516j.
<i>Lycaeides idas longinus</i> Nabokov, 1949.	516k.
<i>Lycaeides idas nabokovi</i> Masters, 1972.	516l.
<i>Lycaeides melissa samuelis</i> Nabokov, 1944.	517a.
<i>Lycaeides melissa melissa</i> (W. H. Edwards, 1873).	517b.
<i>Lycaeides melissa annetta</i> (W. H. Edwards, 1882).	517c.
<i>Lycaeides melissa fridai</i> (F. H. Chermock, [1945]).	517d.
<i>Lycaeides melissa paradoxa</i> (F. H. Chermock, [1945]).	517e.
<i>Plebejus saepiolus amica</i> (W. H. Edwards, 1863).	518a.
<i>Plebejus saepiolus whitmeri</i> F. M. Brown, 1951.	518b.
<i>Plebejus saepiolus gertschi</i> dos Passos, 1938.	518c.
<i>Plebejus saepiolus insulanus</i> Blackmore, 1919.	518d.
<i>Plebejus saepiolus saepiolus</i> (Boisduval, 1852).	518e.
<i>Plebejus saepiolus hilda</i> (J. and F. Grinnell, 1907).	518f.
<i>Plebulina emigdionis</i> (F. Grinnell, 1905).	519.
<i>Icaricia icarioides pembina</i> (W. H. Edwards, 1862).	520a.
<i>Icaricia icarioides lycea</i> (W. H. Edwards, 1864).	520b.
<i>Icaricia icarioides ardea</i> (W. H. Edwards, 1871).	520c.
<i>Icaricia icarioides buchholzi</i> (dos Passos, 1938).	520d.
<i>Icaricia icarioides blackmorei</i> (Barnes and McDunnough, 1919).	520e.
<i>Icaricia icarioides montis</i> (Blackmore, 1923).	520f.
<i>Icaricia icarioides icarioides</i> (Boisduval, 1852).	520g.
<i>Icaricia icarioides pardalis</i> (Behr, 1867).	520h.
<i>Icaricia icarioides pheres</i> (Boisduval, 1852).	520i.

<i>Icaricia icarioides missionensis</i> (Hovanitz, 1937).	520j.
<i>Icaricia icarioides moroensis</i> (Sternitzky, 1930).	520k.
<i>Icaricia icarioides evius</i> (Boisduval, 1869).	520l.
<i>Icaricia icarioides helios</i> (W. H. Edwards, [1871]).	520m.
<i>Icaricia shasta minnehaha</i> (Scudder, 1874).	521a.
<i>Icaricia shasta pitkinensis</i> (Ferris, 1976).	521b.
<i>Icaricia shasta shasta</i> (W. H. Edwards, 1862).	521c.
<i>Icaricia shasta charlestonensis</i> (Austin, 1980).	521d.
<i>Icaricia acmon lutzi</i> (dos Passos, 1938).	522a.
<i>Icaricia acmon texana</i> Goodpasture, 1973.	522b.
<i>Icaricia acmon spangelatus</i> (Burdick, 1942).	522c.
<i>Icaricia acmon acmon</i> (Westwood and Hewitson, [1852]).	522d.
<i>Icaricia lupini lupini</i> (Boisduval, 1869).	523a.
<i>Icaricia lupini monticola</i> (Clemence, 1909).	523b.
<i>Icaricia lupini chlorina</i> (Skinner, 1902).	523c.
<i>Icaricia neurona</i> (Skinner, 1902).	524.
<i>Vacciniina optilete yukona</i> (Holland, 1900).	525a.
<i>Agriades franklinii</i> (Curtis, 1835).	526.
<i>Agriades rusticus rusticus</i> (W. H. Edwards, 1865).	526.1a, (526e).
<i>Agriades rusticus podarce</i> (C. and R. Felder, [1865]).	526.1b, (526f).
<i>Agriades rusticus megallo</i> (McDunnough, 1927).	526.1c, (526d).
<i>Agriades rusticus lacustris</i> (T. N. Freeman, 1939).	526.1d, (526b).
<i>Agriades rusticus bryanti</i> (Leussler, 1935).	526.1e, (526c).

RIODINIDAE

<i>Euselasia abreas</i> (W. H. Edwards, 1881).	527.
<i>Calephelis virginensis</i> (Guérin-Méneville, [1831]).	528.
<i>Calephelis borealis</i> (Grote and Robinson, 1866).	529.
<i>Calephelis nemesis australis</i> (W. H. Edwards, 1877).	530a.
<i>Calephelis nemesis nemesis</i> (W. H. Edwards, 1871).	530b.
<i>Calephelis nemesis dammersi</i> McAlpine, 1971.	530c.
<i>Calephelis nemesis californica</i> McAlpine, 1971.	530d.
<i>Calephelis perditalis</i> Barnes and McDunnough, 1918.	531.
<i>Calephelis wrighti</i> Holland, 1930.	532.
<i>Calephelis mutica</i> McAlpine, 1937.	533.
<i>Calephelis rawsoni</i> McAlpine, 1939.	534.
<i>Calephelis freemani</i> McAlpine, 1971.	535.
<i>Calephelis arizonensis</i> McAlpine, 1971.	536.
<i>Calephelis dreisbachi</i> McAlpine, 1971.	537.
<i>Caria ino melicerta</i> Schaus, 1890.	538a.
<i>Lasaia sula peninsularis</i> Clench, 1972.	539a.
<i>Melanis pixe</i> (Boisduval, [1836]).	540.
<i>Emesis zela cleis</i> (W. H. Edwards, 1870).	541a.
<i>Emesis ares</i> (W. H. Edwards, 1882).	542.
<i>Emesis emesia</i> (Hewitson, 1867).	543.
<i>Apodemia mormo mormo</i> (C. and R. Felder, 1859).	544a.
<i>Apodemia mormo langei</i> J. A. Comstock, 1938.	544b.
<i>Apodemia mormo cythera</i> (W. H. Edwards, 1873).	544c.
<i>Apodemia mormo duryi</i> (W. H. Edwards, 1882).	544d.

<i>Apodemia mormo mejicana</i> (Behr, 1865).	544e.
<i>Apodemia mormo tuolumnensis</i> Opler and Powell, 1962.	544f.
<i>Apodemia mormo virgulti</i> (Behr, 1865).	544g.
<i>Apodemia mormo deserti</i> Barnes and McDunnough, 1918.	544h.
<i>Apodemia mormo dialeuca</i> Opler and Powell, 1962	544i.
<i>Apodemia multiplaga</i> Schaus, 1902.	545.
<i>Apodemia heburni</i> Godman and Salvin, [1886].	546.
<i>Apodemia palmerii</i> (W. H. Edwards), 1870.	547.
<i>Apodemia walkeri</i> Godman and Salvin, [1886].	548.
<i>Apodemia phyciodoides</i> Barnes and Benjamin, 1924.	549.
<i>Apodemia nais</i> (W. H. Edwards, 1876).	550.
<i>Apodemia chisosensis</i> H. A. Freeman, 1964.	551.

LIBYTHEIDAE

<i>Libytheana bachmanii bachmanii</i> (Kirtland, 1852).	552a.
<i>Libytheana bachmanii larvata</i> (Strecker, [1878]).	552b.
<i>Libytheana carinenta mexicana</i> Michener, 1943.	553a.
<i>Libytheana motya</i> (Boisduval and Le Conte, [1833]).	554.

HELICONIIDAE

<i>Agraulis vanillae nigrior</i> Michener, 1942.	555a.
<i>Agraulis vanillae incarnata</i> (Riley, 1926).	555b.
<i>Dione moneta poeyi</i> Butler, 1873.	556a.
<i>Dryadula phaetusa</i> (Linnaeus, 1758).	557.
<i>Dryas iulia largo</i> Clench, 1975.	558a.
<i>Dryas iulia moderata</i> (Riley, 1926).	558b.
<i>Eueides isabella zoracoen</i> Reakirt, 1866.	559a.
<i>Heliconius charitonius tuckeri</i> W. P. Comstock and F. M. Brown, 1950.	560a.
<i>Heliconius charitonius vazquezae</i> W. P. Comstock and F. M. Brown, 1950.	560b.
<i>Heliconius erato petiveranus</i> (Doubleday, 1847).	561a.

NYMPHALIDAE

<i>Euptoieta claudia</i> (Cramer, [1775]).	562.
<i>Euptoieta hegesia hoffmanni</i> W. P. Comstock, 1944.	563a.
<i>Speyeria diana</i> (Cramer, [1775]).	564.
<i>Speyeria cybele cybele</i> (Fabricius, 1775).	565a.
<i>Speyeria cybele novascotiae</i> (McDunnough, 1935).	565b.
<i>Speyeria cybele krautwurmi</i> (Holland, 1931).	565c.
<i>Speyeria cybele pseudocarpenteri</i> (F. and R. Chermock, 1940).	565d.
<i>Speyeria cybele carpenterii</i> (W. H. Edwards, 1876).	565e.
<i>Speyeria cybele charlottii</i> (Barnes, 1897).	565f.
<i>Speyeria cybele letona</i> dos Passos and Grey, 1945.	565g.
<i>Speyeria cybele pugetensis</i> F. Chermock and Frechin, 1947.	565h.
<i>Speyeria cybele leto</i> (Behr, 1862).	565i.
<i>Speyeria aphrodite aphrodite</i> (Fabricius, 1787).	566a.
<i>Speyeria aphrodite winni</i> (Gunder, 1932).	566b.
<i>Speyeria aphrodite alcestis</i> (W. H. Edwards, 1877).	566c.

<i>Speyeria aphrodite manitoba</i> (F. and R. Chermock, 1940).	566d.
<i>Speyeria aphrodite whitehousei</i> (Gunder, 1932).	566e.
<i>Speyeria aphrodite columbia</i> (Hy. Edwards, 1877).	566f.
<i>Speyeria aphrodite ethne</i> (Hemming, 1933).	566g.
<i>Speyeria aphrodite byblis</i> (Barnes and Benjamin, 1926).	566h.
<i>Speyeria idalia</i> (Drury, [1773]).	567.
<i>Speyeria nokomis nokomis</i> (W. H. Edwards, 1862).	568a.
<i>Speyeria nokomis nitocris</i> (W. H. Edwards, 1874).	568b.
<i>Speyeria nokomis coerulescens</i> (Holland, 1900).	568c.
<i>Speyeria nokomis apacheana</i> (Skinner, 1918).	568d.
<i>Speyeria edwardsii</i> (Reakirt, 1866).	569.
<i>Speyeria coronis coronis</i> (Behr, 1864).	570a.
<i>Speyeria coronis hennei</i> (Gunder, 1934).	570b.
<i>Speyeria coronis semiramis</i> (W. H. Edwards, 1886).	570c.
<i>Speyeria coronis simaetha</i> dos Passos and Grey, 1945.	570d, (570e).
<i>Speyeria coronis snyderi</i> (Skinner, 1897).	570e, (570f).
<i>Speyeria coronis halcyone</i> (W. H. Edwards, 1869).	570f, (570g).
<i>Speyeria zerene zerene</i> (Boisduval, 1852).	571a.
<i>Speyeria zerene conchyliaetus</i> (J. A. Comstock, 1925).	571b.
<i>Speyeria zerene gloriosa</i> Moeck, 1957.	571c.
<i>Speyeria zerene sordida</i> (W. G. Wright, 1905).	[see comment in Part II]
<i>Speyeria zerene malcolmi</i> (J. A. Comstock, 1920).	571d.
<i>Speyeria zerene carolae</i> (dos Passos and Grey, 1942).	571e.
<i>Speyeria zerene hippolyta</i> (W. H. Edwards, 1879).	571f.
<i>Speyeria zerene behrensi</i> (W. H. Edwards, 1869).	571g.
<i>Speyeria zerene myrtleae</i> dos Passos and Grey, 1945.	571h.
<i>Speyeria zerene bremnerii</i> (W. H. Edwards, 1872).	[see comment in Part II]
<i>Speyeria zerene picta</i> (McDunnough, 1924).	571i.
<i>Speyeria zerene garretti</i> (Gunder, 1932).	571j.
<i>Speyeria zerene sinope</i> dos Passos and Grey, 1945.	571k.
<i>Speyeria zerene platina</i> (Skinner, 1897).	571l.
<i>Speyeria zerene gundleri</i> (J. A. Comstock, 1925).	571m.
<i>Speyeria callippe callippe</i> (Boisduval, 1852).	571n.
<i>Speyeria callippe comstocki</i> (Gunder, 1925).	572a.
<i>Speyeria callippe liliana</i> (Hy. Edwards, [1877]).	572b.
<i>Speyeria callippe semifirida</i> (McDunnough, 1924).	572c.
<i>Speyeria callippe elaine</i> dos Passos and Grey, 1945.	572d.
<i>Speyeria callippe rupestris</i> (Behr, 1863).	572e.
<i>Speyeria callippe juba</i> (Boisduval, 1869).	572f.
<i>Speyeria callippe laura</i> (W. H. Edwards, 1879).	[see comment in Part II]
<i>Speyeria callippe nevadensis</i> (W. H. Edwards, 1870).	572g.
<i>Speyeria callippe macaria</i> (W. H. Edwards, 1877).	572h.
<i>Speyeria callippe laurina</i> (W. G. Wright, 1905).	572i, (572j).
<i>Speyeria callippe harmonia</i> dos Passos and Grey, 1945.	572j, (572k).
<i>Speyeria callippe meadii</i> (W. H. Edwards, 1872).	572l.
<i>Speyeria callippe gallatini</i> (McDunnough, 1929).	572m, (572n).
<i>Speyeria callippe calgariana</i> (McDunnough, 1924).	572o.
<i>Speyeria egleis egleis</i> (Behr, 1862).	572p.
<i>Speyeria egleis tehachapina</i> (J. A. Comstock, 1920).	573a.
<i>Speyeria egleis oweni</i> (W. H. Edwards, 1892).	573b, (573e).
	573c, (573f).

<i>Speyeria egleis linda</i> (dos Passos and Grey, 1942).	573d, (573g).
<i>Speyeria egleis macdunnoughi</i> (Gunder, 1932).	573e, (573h).
<i>Speyeria egleis albrichti</i> (Gunder, 1932).	573f, (573i).
<i>Speyeria egleis utahensis</i> (Skinner, 1919).	573g, (573j).
<i>Speyeria egleis toiyabe</i> Howe, 1975.	573h, (573k).
<i>Speyeria egleis secreta</i> dos Passos and Grey, 1945.	573i, (573l).
<i>Speyeria egleis moecki</i> Hammond and Dornfeld, 1983.	573j.
<i>Speyeria adiaste adiaste</i> (W. H. Edwards, 1864).	573.1a, (573b).
<i>Speyeria adiaste atossa</i> (W. H. Edwards, 1890).	573.1b, (573d).
<i>Speyeria adiaste clemencei</i> (J. A. Comstock, 1925).	573.1c, (573c).
<i>Speyeria atlantis atlantis</i> (W. H. Edwards, 1862).	574a.
<i>Speyeria atlantis canadensis</i> (dos Passos, 1935).	574b.
<i>Speyeria atlantis hollandi</i> (F. and R. Chermock, 1940).	574c.
<i>Speyeria atlantis hesperis</i> (W. H. Edwards, 1864).	574d.
<i>Speyeria atlantis nikias</i> (Ehrmann, 1917).	574e.
<i>Speyeria atlantis dorothea</i> Moeck, 1947.	574f.
<i>Speyeria atlantis nausicaa</i> (W. H. Edwards, 1874).	574g.
<i>Speyeria atlantis schellbachii</i> Garth, 1949.	574h.
<i>Speyeria atlantis chitone</i> (W. H. Edwards, 1879).	574i.
<i>Speyeria atlantis wasatchia</i> dos Passos and Grey, 1945.	574j.
<i>Speyeria atlantis greyi</i> (Moeck, 1950).	574k.
<i>Speyeria atlantis tetonia</i> dos Passos and Grey, 1945.	574l.
<i>Speyeria atlantis viola</i> dos Passos and Grey, 1945.	574m.
<i>Speyeria atlantis dodgei</i> (Gunder, 1931).	574n.
<i>Speyeria atlantis irene</i> (Boisduval, 1869).	574o.
<i>Speyeria atlantis electa</i> (W. H. Edwards, 1878).	574p.
<i>Speyeria atlantis lurana</i> dos Passos and Grey, 1945.	574q.
<i>Speyeria atlantis hutchinsi</i> dos Passos and Grey, 1947.	574r.
<i>Speyeria atlantis beani</i> (Barnes and Benjamin, 1926).	574s.
<i>Speyeria atlantis lais</i> (W. H. Edwards, 1884).	574t.
<i>Speyeria atlantis dennisi</i> dos Passos and Grey, 1947.	574u.
<i>Speyeria atlantis ratonensis</i> Scott, 1981.	574v.
<i>Speyeria atlantis elko</i> Austin, 1983.	574w.
<i>Speyeria atlantis capitanensis</i> R. Holland, 1988.	574x.
<i>Speyeria hydaspe hydaspe</i> (Boisduval, 1869).	575a.
<i>Speyeria hydaspe viridicornis</i> (J. A. Comstock, 1925).	575b.
<i>Speyeria hydaspe purpurascens</i> (Hy. Edwards, [1877]).	575c.
<i>Speyeria hydaspe minor</i> dos Passos and Grey, 1947.	575d.
<i>Speyeria hydaspe rhodope</i> (W. H. Edwards, 1874).	575e.
<i>Speyeria hydaspe sakuratala</i> (Skinner, 1911).	575f.
<i>Speyeria hydaspe conquista</i> dos Passos and Grey, 1945.	575g.
<i>Speyeria mormonia bischoffii</i> (W. H. Edwards, 1870).	576a.
<i>Speyeria mormonia opis</i> (W. H. Edwards, 1874).	576b.
<i>Speyeria mormonia washingtonia</i> (Barnes and McDunnough, 1913).	576c, (576d).
<i>Speyeria mormonia erinna</i> (W. H. Edwards, 1883).	576d, (576e).
<i>Speyeria mormonia artonis</i> (W. H. Edwards, 1881).	576e, (576g).
<i>Speyeria mormonia mormonia</i> (Boisduval, 1869).	576f, (576h).
<i>Speyeria mormonia eury nome</i> (W. H. Edwards, 1872).	576g, (576i).
<i>Speyeria mormonia luski</i> (Barnes and McDunnough, 1913).	576h, (576j).
<i>Boloria napaea alaskensis</i> (Holland, 1900).	577a.

<i>Boloria napaea nearctica</i> Verity, 1932.	577b.
<i>Boloria napaea halli</i> Klots, 1940.	577c.
<i>Clossiana eunomia triclaris</i> (Hübner, [1821]).	578a.
<i>Clossiana eunomia alticola</i> (Barnes and McDunnough, 1913).	578b.
<i>Clossiana eunomia dawsoni</i> (Barnes and McDunnough, 1916).	578c.
<i>Clossiana eunomia laddi</i> (Klots, 1940).	578d, (578e).
<i>Clossiana eunomia ursadenitis</i> (Ferris and Groothuis, 1971).	578e, (578f).
<i>Clossiana eunomia denali</i> (Klots, 1940).	578f, (578g).
<i>Clossiana selene myrina</i> (Cramer, [1777]).	579a.
<i>Clossiana selene nebraskensis</i> (Holland, 1928).	579b.
<i>Clossiana selene sabulocollis</i> (Kohler, 1977).	579c.
<i>Clossiana selene tollandensis</i> (Barnes and Benjamin, 1925).	579d.
<i>Clossiana selene albequina</i> (Holland, 1928).	579e.
<i>Clossiana selene atrocostalis</i> (Huard, 1927).	579f.
<i>Clossiana selene terraenovae</i> (Holland, 1928).	579g.
<i>Clossiana bellona bellona</i> (Fabricius, 1775).	580a.
<i>Clossiana bellona toddi</i> (Holland, 1928).	580b.
<i>Clossiana bellona jenistae</i> (D. Stallings and Turner, 1946).	580c.
<i>Clossiana frigga saga</i> (Staudinger, 1861).	581a.
<i>Clossiana frigga gibsoni</i> (Barnes and McDunnough, 1926).	581b.
<i>Clossiana frigga sagata</i> (Barnes and McDunnough, 1923).	581c.
<i>Clossiana improba improba</i> (Butler, 1877).	582a.
<i>Clossiana improba harryi</i> Ferris, 1984.	582b.
<i>Clossiana acrocema</i> (Gall and Sperling, 1980).	582.1.
<i>Clossiana kriemhild</i> (Strecker, [1878-1879]).	583.
<i>Clossiana epithore epithore</i> (W. H. Edwards, [1864]).	584a.
<i>Clossiana epithore chermocki</i> (E. and S. Perkins, 1966).	584b.
<i>Clossiana epithore uslui</i> Koçak, 1984.	584c.
<i>Clossiana epithore sierra</i> (E. Perkins, 1973).	584d.
<i>Clossiana polaris polaris</i> (Boisduval, [1828]).	585a.
<i>Clossiana polaris stellata</i> (Masters, 1972).	585b.
<i>Clossiana freija freija</i> (Thunberg, 1791).	586a.
<i>Clossiana freija tarquinius</i> (Curtis, 1835).	586b.
<i>Clossiana freija natazhati</i> (Gibson, 1920).	[see comment in Part II]
<i>Clossiana freija browni</i> (Higgins, 1953).	586d, (586e).
<i>Clossiana alberta</i> (W. H. Edwards, 1890).	587.
<i>Clossiana astarte astarte</i> (Doubleday and Hewitson, [1847]).	588a.
<i>Clossiana astarte distincta</i> (Gibson, 1920).	588b.
<i>Clossiana astarte tschukotkensis</i> Wyatt, 1961.	588c.
<i>Clossiana titania boisduvalii</i> (Duponchel, [1832]).	589a.
<i>Clossiana titania rainieri</i> (Barnes and McDunnough, 1913).	589b.
<i>Clossiana titania grandis</i> (Barnes and McDunnough, 1916).	589c.
<i>Clossiana titania montinus</i> (Scudder, 1863).	589d.
<i>Clossiana titania ingens</i> (Barnes and McDunnough, 1918).	589e.
<i>Clossiana titania helena</i> (W. H. Edwards, 1871).	589f.
<i>Clossiana chariclea arctica</i> (Zetterstedt, [1839]).	590a.
<i>Clossiana chariclea butleri</i> (W. H. Edwards, 1883).	590b.
<i>Poladryas minuta minuta</i> (W. H. Edwards, 1861).	591a, (591).
<i>Poladryas minuta nympha</i> (W. H. Edwards, 1884).	591b, (592b).
<i>Poladryas arachne arachne</i> (W. H. Edwards, 1869).	592a.

<i>Poladryas arachne monache</i> (J. A. Comstock, 1918).	592b, (592c).
<i>Thessalia theona thekla</i> (W. H. Edwards, 1870).	593a.
<i>Thessalia theona bolli</i> (W. H. Edwards, 1877).	593b.
<i>Thessalia chinatiensis</i> (Tinkham, 1944).	594.
<i>Thessalia cyneas cyneas</i> (Godman and Salvin, 1878).	595a, (595).
<i>Thessalia fulvia fulvia</i> (W. H. Edwards, 1879).	596a, (596).
<i>Thessalia fulvia coronado</i> M. Smith and Brock, 1988.	596b.
<i>Thessalia fulvia pariaensis</i> M. Smith and Brock, 1988.	596c.
<i>Thessalia leanira leanira</i> (C. and R. Felder, 1860).	597a.
<i>Thessalia leanira wrighti</i> (W. H. Edwards, 1886).	597b.
<i>Thessalia leanira cerrita</i> (W. G. Wright, 1905).	597c.
<i>Thessalia leanira alma</i> (Strecker, 1878).	597d.
<i>Thessalia leanira oregonensis</i> (Bauer, 1975).	597e.
<i>Chlosyne californica</i> (W. G. Wright, 1905).	598.
<i>Chlosyne lacinia crocale</i> (W. H. Edwards, 1874).	599a.
<i>Chlosyne lacinia adjutrix</i> Scudder, 1875.	599b.
<i>Chlosyne definita</i> (E. M. Aaron, 1884).	600.
<i>Chlosyne endeis</i> (Godman and Salvin, 1894).	601.
<i>Chlosyne erodyle</i> (H. W. Bates, 1864).	602.
<i>Chlosyne janais</i> (Drury, [1782]).	603.
<i>Chlosyne rosita browni</i> Bauer, 1960.	604a.
<i>Chlosyne ehrenbergerii</i> (Geyer, 1833).	604.1.
<i>Chlosyne melitaeoides</i> (C. and R. Felder, 1867).	604.2.
<i>Charidryas gorgone gorgone</i> (Hübner, 1810).	605a.
<i>Charidryas gorgone carlota</i> (Reakirt, 1866).	605b.
<i>Charidryas nycteis nycteis</i> (Doubleday and Hewitson, [1847]).	606a.
<i>Charidryas nycteis drusius</i> (W. H. Edwards, 1884).	606b.
<i>Charidryas nycteis reversa</i> (F. and R. Chermock, 1940).	606c.
<i>Charidryas harrisii harrisii</i> (Scudder, 1864).	607a.
<i>Charidryas harrisii liggettii</i> (Avinoff, 1930).	607b.
<i>Charidryas harrisii hanhami</i> (Fletcher, 1904).	607c.
<i>Charidryas palla palla</i> (Boisduval, 1852).	608a.
<i>Charidryas palla whitneyi</i> (Behr, 1863). [see comment in Part II]	608b.
<i>Charidryas palla vallismortis</i> (J. W. Johnson, 1938).	608c.
<i>Charidryas palla calydon</i> (Holland, 1931).	608d.
<i>Charidryas palla flavula</i> (Barnes and McDunnough, 1918).	608e.
<i>Charidryas palla sterope</i> (W. H. Edwards, 1870).	608f.
<i>Charidryas acastus</i> (W. H. Edwards, 1874).	609.
<i>Charidryas neuvoegeni neuvoegeni</i> (Skinner, 1895).	610a.
<i>Charidryas neuvoegeni sabina</i> (W. G. Wright, 1905).	610b.
<i>Charidryas gabbii</i> (Behr, 1863).	611.
<i>Charidryas damoetas damoetas</i> (Skinner, 1902). [see comment in Part II]	612a.
<i>Charidryas damoetas malcolmi</i> (J. A. Comstock), 1926.	612b.
<i>Charidryas hoffmanni hoffmanni</i> (Behr, 1863).	613a.
<i>Charidryas hoffmanni segregata</i> (Barnes and McDunnough, 1918).	613b.
<i>Charidryas hoffmanni manchada</i> (Bauer, 1959[1960]).	613c.
<i>Microtia elva</i> H. W. Bates, 1864.	614.
<i>Dymasia dymas dymas</i> (W. H. Edwards, 1877).	615a.
<i>Dymasia dymas chara</i> (W. H. Edwards, 1883).	615b.
<i>Dymasia dymas imperialis</i> (Bauer, [1959]).	615c.

<i>Texola elada ulrica</i> (W. H. Edwards, 1877).	616a.
<i>Texola elada perse</i> (W. H. Edwards, 1882).	616b.
<i>Anthanassa texana texana</i> (W. H. Edwards, 1863).	617a.
<i>Anthanassa texana seminole</i> (Skinner, 1911).	617b.
<i>Anthanassa frisia</i> (Poey, 1832).	618.
<i>Anthanassa pitolyca</i> (H. W. Bates, 1864).	619.
<i>Anthanassa tulcis</i> (H. W. Bates, 1864).	620.
<i>Phyciodes vesta</i> (W. H. Edwards, 1869).	621.
<i>Phyciodes phaon</i> (W. H. Edwards, 1864).	622.
<i>Phyciodes tharos arcticus</i> dos Passos, 1935.	623a.
<i>Phyciodes tharos tharos</i> (Drury, [1773]).	623b.
<i>Phyciodes tharos distinctus</i> Bauer, 1975.	623c.
<i>Phyciodes tharos pascoensis</i> W. G. Wright, 1905. [see comment in Part II]	623d.
<i>Phyciodes batesii</i> (Reakirt, 1865).	624.
<i>Phyciodes pratensis pratensis</i> (Behr, 1863).	625a.
<i>Phyciodes pratensis montanus</i> (Behr, 1863).	625b.
<i>Phyciodes pratensis camillus</i> W. H. Edwards, 1871.	625c.
<i>Phyciodes pictus pictus</i> (W. H. Edwards, 1865).	626a.
<i>Phyciodes orseis orseis</i> W. H. Edwards, 1871.	627a.
<i>Phyciodes orseis herlani</i> Bauer, 1975.	627b.
<i>Phyciodes pallidus pallidus</i> (W. H. Edwards, 1864).	628a.
<i>Phyciodes pallidus barnesi</i> Skinner, 1897.	628b.
<i>Phyciodes mylitta mylitta</i> (W. H. Edwards, 1861).	629a.
<i>Phyciodes mylitta callina</i> (Boisduval, 1869).	629b.
<i>Phyciodes mylitta thebais</i> Godman and Salvin, 1878.	629c.
<i>Tegosa anieta luka</i> Higgins, 1981.	629.5a.
<i>Euphydryas gillettii</i> (Barnes, 1921).	630.
<i>Euphydryas anicia alena</i> Barnes and Benjamin, 1926.	631a.
<i>Euphydryas anicia anicia</i> (Doubleday and Hewitson, [1848]).	631b.
<i>Euphydryas anicia bakeri</i> D. Stallings and Turner, 1945.	631c.
<i>Euphydryas anicia bernadetta</i> Leussler, 1920.	631d.
<i>Euphydryas anicia capella</i> (Barnes, 1897).	631e.
<i>Euphydryas anicia carmentis</i> Barnes and Benjamin, 1926.	631f.
<i>Euphydryas anicia chuskae</i> (Ferris and R. Holland, 1980).	631g.
<i>Euphydryas anicia claudcrofti</i> (Ferris and R. Holland, 1980).	631h.
<i>Euphydryas anicia effi</i> D. Stallings and Turner, 1945.	631i.
<i>Euphydryas anicia eurytion</i> (Mead, 1875).	631j.
<i>Euphydryas anicia helvia</i> (Scudder, 1869).	631k.
<i>Euphydryas anicia hermosa</i> (W. G. Wright, 1905).	631l.
<i>Euphydryas anicia hopfingeri</i> Gunder, 1931.	631m.
<i>Euphydryas anicia howlandi</i> D. Stallings and Turner, [1947].	631n.
<i>Euphydryas anicia irelandi</i> Gunder, 1929.	631o.
<i>Euphydryas anicia macyi</i> Fender and Jewett, 1953.	631p.
<i>Euphydryas anicia magdalena</i> Barnes and McDunnough, 1918.	631q.
<i>Euphydryas anicia maria</i> (Skinner, 1899).	631r.
<i>Euphydryas anicia morandi</i> Gunder, 1928.	631s.
<i>Euphydryas anicia veaziae</i> Fender and Jewett, 1953.	631t.
<i>Euphydryas anicia wheeleri</i> (Hy. Edwards, 1881).	631u.
<i>Euphydryas anicia windi</i> Gunder, 1932.	631v.
<i>Euphydryas chalcedona chalcedona</i> (Doubleday, [1847]).	632a.

<i>Euphydryas chalcedona colon</i> (W. H. Edwards, 1881).	632b, (633a).
<i>Euphydryas chalcedona corralensis</i> T. and J. Emmel, [1973].	632c, (632b).
<i>Euphydryas chalcedona dwinellei</i> (Hy. Edwards, 1881).	632d, (632c).
<i>Euphydryas chalcedona kingstonensis</i> T. and J. Emmel, [1973].	632e.
<i>Euphydryas chalcedona klotzi</i> dos Passos, 1938.	632f.
<i>Euphydryas chalcedona macglashanii</i> (Rivers, 1888).	632g.
<i>Euphydryas chalcedona nevadensis</i> Bauer, 1975.	632h, (633b).
<i>Euphydryas chalcedona olancha</i> (W. G. Wright, 1905).	632i, (632h).
<i>Euphydryas chalcedona paradoxa</i> McDunnough, 1927.	632j, (633c).
<i>Euphydryas chalcedona perdiccas</i> (W. H. Edwards, 1881).	632k, (633d).
<i>Euphydryas chalcedona quino</i> (Behr, 1863).	632l, (632i).
<i>Euphydryas chalcedona sierra</i> (W. G. Wright, 1905).	632m, (632j).
<i>Euphydryas chalcedona sperryi</i> F. and R. Chermock, 1945.	632n, (632k).
<i>Euphydryas chalcedona wallacensis</i> Gunder, 1928.	632o, (633e).
<i>Euphydryas editha alebarki</i> Ferris, [1971].	634a.
<i>Euphydryas editha augusta</i> (W. H. Edwards, 1890).	634b.
<i>Euphydryas editha aurilacus</i> Gunder, 1928.	634c.
<i>Euphydryas editha baroni</i> (W. H. Edwards, 1879).	634d.
<i>Euphydryas editha bayensis</i> Sternitzky, 1937.	634e.
<i>Euphydryas editha beani</i> (Skinner, 1897).	634f.
<i>Euphydryas editha colonia</i> (W. G. Wright, 1905).	634g.
<i>Euphydryas editha editha</i> (Boisduval, 1852).	634h.
<i>Euphydryas editha edithana</i> (Strand, 1914).	634i.
<i>Euphydryas editha fridayi</i> Gunder, 1931.	634j.
<i>Euphydryas editha gunninsonensis</i> F. M. Brown, [1971].	634k.
<i>Euphydryas editha hutchinsi</i> McDunnoughi, 1928.	634l.
<i>Euphydryas editha insularis</i> T. and J. Emmel, [1975].	634m.
<i>Euphydryas editha lawrencei</i> Gunder, 1931.	634n.
<i>Euphydryas editha lehmani</i> Gunder, 1929.	634o.
<i>Euphydryas editha monoensis</i> Gunder, 1928.	634p.
<i>Euphydryas editha nubigena</i> (Behr, 1863).	634q.
<i>Euphydryas editha rubicunda</i> (Hy. Edwards, 1881).	634r.
<i>Euphydryas editha taylori</i> (W. H. Edwards, 1888).	634s.
<i>Euphydryas editha wrighti</i> Gunder, 1929.	634t.
<i>Euphydryas editha luestherae</i> Murphy and P. Ehrlich, 1980.	634u.
<i>Euphydryas editha koreti</i> Murphy and P. Ehrlich, (1984).	634v.
<i>Euphydryas phaeton phaeton</i> (Drury, [1773]).	635a.
<i>Euphydryas phaeton ozarkae</i> Masters, 1968.	635b.
<i>Polygonia interrogationis</i> (Fabricius, 1798).	636.
<i>Polygonia comma</i> (Harris, 1842).	637.
<i>Polygonia satyrus</i> (W. H. Edwards, 1869).	638.
<i>Polygonia faunus faunus</i> (W. H. Edwards, 1862).	639a.
<i>Polygonia faunus smythi</i> A. H. Clark, 1937.	639b.
<i>Polygonia faunus rusticus</i> (W. H. Edwards, 1874).	639c.
<i>Polygonia faunus arcticus</i> Leussler, 1935.	639d.
<i>Polygonia faunus hylas</i> (W. H. Edwards, 1872).	639e, (640).
<i>Polygonia zephyrus</i> (W. H. Edwards, 1870).	642.
<i>Polygonia gracilis</i> (Grote and Robinson, 1867).	643.
<i>Polygonia oreas oreas</i> (W. H. Edwards, 1869).	644a.
<i>Polygonia oreas silenus</i> (W. H. Edwards, 1870).	644b.

<i>Polygonia progne progne</i> (Cramer, [1776]).	645a.
<i>Polygonia progne nigrozephyrus</i> Scott, 1984.	[see comment in Part II] 645b.
<i>Nymphalis vaualbum j-album</i> (Boisduval and Le Conte, [1883]).	646.
<i>Nymphalis californica californica</i> (Boisduval, 1852).	647a.
<i>Nymphalis californica herri</i> Field, 1936.	647b.
<i>Nymphalis antiopa antiopa</i> (Linnaeus, 1758).	648a.
<i>Nymphalis antiopa hyperborea</i> (Seitz, 1914).	648b.
<i>Aglais milberti viola</i> dos Passos, 1938.	649a.
<i>Aglais milberti milberti</i> (Godart, [1819]).	649b.
<i>Vanessa virginensis</i> (Drury, 1773).	650.
<i>Vanessa cardui</i> (Linnaeus, 1758).	651.
<i>Vanessa annabella</i> (Field, 1971).	652.
<i>Vanessa atalanta rubria</i> (Fruhstorfer, 1909).	653a.
<i>Hypanartia lethe</i> (Fabricius, 1793).	654.
<i>Hypolimnas misippus</i> (Linnaeus, 1764).	655.
<i>Junonia coenia</i> Hübner, [1822].	656.
<i>Junonia genoveva zonalis</i> C. and R. Felder, [1864-1867].	657a.
<i>Junonia evarete</i> (Cramer, [1779]).	658.
<i>Anartia jatrophae guantanamo</i> Munroe, 1942.	659a.
<i>Anartia jatrophae luteipicta</i> Fruhstorfer, 1907.	659b.
<i>Anartia chrysopoelea</i> Hübner, [1824].	660.
<i>Anartia fatima</i> (Fabricius, 1793).	661.
<i>Siproeta stelenes biplagiata</i> (Fruhstorfer, 1907).	662a.
<i>Basilarchia arthemis arthemis</i> (Drury, [1773]).	663a.
<i>Basilarchia arthemis rubrofasciata</i> Barnes and McDunnough, 1916.	663b.
<i>Basilarchia arthemis astyanax</i> (Fabricius, 1775).	663c.
<i>Basilarchia arthemis arizonensis</i> (W. H. Edwards, 1882).	663d.
<i>Basilarchia archippus archippus</i> (Cramer, [1776]).	664a.
<i>Basilarchia archippus floridensis</i> (Strecker, 1878).	664b.
<i>Basilarchia archippus watsoni</i> dos Passos, 1938.	664c.
<i>Basilarchia archippus obsoleta</i> (W. H. Edwards, 1882).	664d.
<i>Basilarchia archippus lahontani</i> (Herlan, 1971).	664e.
<i>Basilarchia weidemeyerii oberfoelli</i> (F. M. Brown, 1960).	665a.
<i>Basilarchia weidemeyerii weidemeyerii</i> W. H. Edwards, 1861.	665b.
<i>Basilarchia weidemeyerii latifascia</i> (E. M. and S. F. Perkins, 1967).	665c.
<i>Basilarchia weidemeyerii nevadae</i> Barnes and Benjamin, 1924.	665d.
<i>Basilarchia weidemeyerii angustifascia</i> Barnes and Benjamin, 1912.	665e.
<i>Basilarchia weidemeyerii siennafascia</i> (Austin and Mullins, 1984).	665f.
<i>Basilarchia lorquini burrisoni</i> (Maynard, 1891).	666a.
<i>Basilarchia lorquini lorquini</i> (Boisduval, 1852).	666b.
<i>Adelpha fessonia</i> (Hewitson, 1847).	667.
<i>Adelpha bredowii eulalia</i> (Doubleday and Hewitson, [1848]).	668a.
<i>Adelpha bredowii californica</i> (Butler, [1865]).	668b.
<i>Epiphile adrasta adrasta</i> Hewitson, 1876.	668.5a.
<i>Myscelia ethusa ethusa</i> (Doyère, 1840).	669a.
<i>Myscelia cyananthe skinneri</i> Mengel, 1894.	670a, (670, 672).
<i>Eunica monima</i> (Stoll, [1782]).	673.
<i>Eunica tatila tatilista</i> Kaye, 1926.	674a.
<i>Dynamine dyonis</i> Geyer, [1837].	675.
<i>Dynamine titlia</i> (Geyer, 1823).	675.1.

<i>Diaethria clymena</i> (Cramer, [1776]).	676.
<i>Diaethria asteria</i> (Godman and Salvin, 1894).	677.
<i>Diaethria anna</i> (Guérin-Méneville, 1844).	677.1.
<i>Mestra amymone</i> (Ménétriés, 1857).	678.
<i>Mestra cana</i> (Erichson, 1848).	679.
<i>Biblis hyperia aganisa</i> Boisduval, 1836.	680a.
<i>Hamadryas februa ferentina</i> (Godart, [1824]).	681a.
<i>Hamadryas amphichloe diasia</i> (Fruhstorfer, 1916).	682a.
<i>Hamadryas atlantis lelaps</i> (Godman and Salvin, [1883]).	683a.
<i>Hamadryas feronia farinulenta</i> (Fruhstorfer, 1916).	684a, (682a).
<i>Hamadryas guatemalena marmarice</i> (H. W. Fruhstorfer, 1916).	684.1a.
<i>Hamadryas iphthime joannae</i> Jenkins, 1983.	684.2a.
<i>Hamadryas amphinome mexicana</i> (Lucas, 1853).	685a.
<i>Historis odius</i> (Fabricius, 1775).	686.
<i>Historis acheronta cadmus</i> (Cramer, [1775]).	687a.
<i>Smyrna karwinskii</i> Geyer, [1883].	688.
<i>Smyrna blomfildia datis</i> Fruhstorfer, 1781.	688.1a.
<i>Marpesia zerynthia</i> Hübner, [1823].	689.
<i>Marpesia chiron</i> (Fabricius, 1775).	690.
<i>Marpesia petreus</i> (Cramer, [1776]).	691.
<i>Marpesia eleuchea</i> Hübner, [1818].	692.
<i>Anaea aidea</i> (Guérin-Méneville, [1844]).	693.
<i>Anaea floridalis</i> F. Johnson and W. P. Comstock, 1941.	694.
<i>Anaea andria</i> Scudder, 1875.	695.
<i>Memphis glycerium</i> (Doubleday, [1850]).	696.
<i>Memphis pithyusa</i> (R. Felder, 1869).	697.
<i>Memphis echemus</i> (Doubleday and Hewitson, [1849]).	697.1.
<i>Asterocampa celitis celitis</i> (Boisduval and Le Conte, [1837]).	698a, (698).
<i>Asterocampa celitis reinthali</i> Friedlander, "1986(1987)"[1988].	698b.
<i>Asterocampa celitis antonia</i> (W. H. Edwards, [1878]).	698c, (699).
<i>Asterocampa leilia</i> (W. H. Edwards, 1874).	699, (702a).
<i>Asterocampa clyton clyton</i> (Boisduval and Le Conte, [1837]).	700a, (704).
<i>Asterocampa clyton flora</i> (W. H. Edwards, 1876).	700b, (705).
<i>Asteroicampa clyton texana</i> (Skinner, 1911).	700c, (706).
<i>Asterocampa clyton louisa</i> Stallings and Turner, 1947.	700d, (707).
<i>Doxocopa pavon</i> (Latreille, [1809]).	708.
<i>Doxocopa laure</i> (Drury, [1773]).	709.

SATYRIDAE

<i>Enodia portlandia portlandia</i> (Fabricius, 1781).	710a.
<i>Enodia portlandia floralae</i> (J. R. Heitzman and dos Passos, 1974).	710b.
<i>Enodia portlandia missarkae</i> (J. R. Heitzman and dos Passos, 1974).	710c.
<i>Enodia anthedon</i> A. H. Clark, 1936.	711.
<i>Enodia creola</i> (Skinner, 1897).	712.
<i>Satyrodes eurydice eurydice</i> (Linnaeus in Johansson, 1763).	713a.
<i>Satyrodes eurydice fumosa</i> Leussler, 1916.	713b.
<i>Satyrodes appalachia appalachia</i> (R. L. Chermock, 1947).	714a.
<i>Satyrodes appalachia leeuwi</i> (Gatrell and Arbogast, 1974).	714b.
<i>Cyllopsis pyramon nabokovi</i> L. Miller, 1974.	715a.

<i>Cyllopsis henshawi</i> (W. H. Edwards, 1876).	716.
<i>Cyllopsis pertepida dorothea</i> (Nabokov, 1942).	717a.
<i>Cyllopsis pertepida maniola</i> (Nabokov, 1942).	717b.
<i>Cyllopsis pertepida avicula</i> (Nabokov, 1942).	717c.
<i>Cyllopsis gemma gemma</i> (Hübner, [1808]).	718a.
<i>Cyllopsis gemma freemani</i> (D. Stallings and Turner, 1946).	718b.
<i>Hermeuptychia hermes</i> (Fabricius, 1775).	719.
<i>Hermeuptychia sosybius</i> (Fabricius 1793).	720.
<i>Neonympha areolata areolata</i> (J. E. Smith, 1797).	721a.
<i>Neonympha areolata septentrionalis</i> Davis, 1924.	721b.
<i>Neonympha mitchellii</i> French, 1889.	722.
<i>Megisto cymela cymela</i> (Cramer, [1777]).	723a.
<i>Megisto cymela viola</i> (Maynard, 1891).	723b.
<i>Megisto rubricata rubricata</i> (W. H. Edwards, 1871).	724a.
<i>Megisto rubricata smithorum</i> (Wind, 1946).	724b.
<i>Megisto rubricata cheneyorum</i> (R. Chermock, [1949]).	724c.
<i>Paramacera allyni</i> L. Miller, 1972.	725.
<i>Coenonympha haydenii</i> (W. H. Edwards, 1872).	726.

Coenonympha tullia Complex

<i>kodiak</i> W. H. Edwards, 1869.	727a.
<i>yukonensis</i> Holland, 1900.	727b.
<i>mixturrata</i> Alpheraky, 1897.	727c.
<i>mcisaaci</i> dos Passos, 1936.	727d, (728a).
<i>nipisiquit</i> McDunnough, 1939.	727e, (728b).
<i>heinemani</i> F. M. Brown, [1959].	727f, (728c).
<i>inornata</i> W. H. Edwards, 1861.	727g, (728d).
<i>benjamini</i> McDunnough, 1928.	727h, (728e).
<i>mackenziei</i> Davenport, 1936.	727i, (729a).
<i>ochracea</i> W. H. Edwards, 1861.	727j, (729b).
<i>brenda</i> W. H. Edwards, 1869.	727k, (729c).
<i>subfusca</i> Barnes and Benjamin, 1926.	727l, (729d).
<i>furcae</i> Barnes and Benjamin, 1926.	727m, (729e).
<i>mono</i> Burdick, 1942.	727n, (729f).
<i>elko</i> W. H. Edwards, 1881.	727o, (730a).
<i>columbiana</i> McDunnough, 1928.	727p, (730b).
<i>eunomia</i> Dornfeld, 1967.	727q, (730c).
<i>ampelos</i> W. H. Edwards, 1871.	727r, (730d).
<i>insulana</i> McDunnough, 1928.	727s, (730e).
<i>california</i> Westwood, [1851].	727t, (731a).
<i>eryngii</i> Hy. Edwards, 1877.	727u, (731b).

<i>Cercyonis pegala abbotti</i> F. M. Brown, 1969.	732a.
<i>Cercyonis pegala pegala</i> (Fabricius, 1775).	732b.
<i>Cercyonis pegala alope</i> (Fabricius, 1793).	732c.
<i>Cercyonis pegala nephele</i> (W. Kirby, 1837).	732d.
<i>Cercyonis pegala olympus</i> (W. H. Edwards, 1880).	732e.
<i>Cercyonis pegala texana</i> (W. H. Edwards, 1880).	732f.
<i>Cercyonis pegala ino</i> Hall, 1924.	732g.

<i>Cercyonis pegala boopis</i> (Behr, 1864).	732h.
<i>Cercyonis pegala ariane</i> (Boisduval, 1852).	732i.
<i>Cercyonis pegala stephensi</i> (W. G. Wright, 1905).	732j.
<i>Cercyonis pegala wheeleri</i> (W. H. Edwards, 1873).	732k.
<i>Cercyonis pegala damei</i> Barnes and Benjamin, 1926.	732l.
<i>Cercyonis meadii meadii</i> (W. H. Edwards, 1872).	733a.
<i>Cercyonis meadii alamosa</i> T. and J. Emmel, 1969.	733b.
<i>Cercyonis meadii melania</i> (Wind, 1946).	733c.
<i>Cercyonis meadii mexicana</i> R. Chermock, 1948.	733d.
<i>Cercyonis sthenele masoni</i> Cross, 1937.	734a.
<i>Cercyonis sthenele paulus</i> (W. H. Edwards, 1879).	734b.
<i>Cercyonis sthenele sthenele</i> (Boisduval, 1852).	734c.
<i>Cercyonis oetus charon</i> (W. H. Edwards, 1872).	735a.
<i>Cercyonis oetus silvestris</i> (W. H. Edwards, 1861). [see comment in Part II]	735b.
<i>Cercyonis oetus oetus</i> (Boisduval, 1869).	735c.
<i>Cercyonis oetus pallescens</i> Emmel and Emmel, 1971.	735d.
<i>Erebia vidleri</i> Elwes, 1898.	736.
<i>Erebia rossii ornata</i> Leussler, 1935.	737a.
<i>Erebia rossii rossii</i> (Curtis, 1835).	737b.
<i>Erebia rossii gabrieli</i> dos Passos, 1949.	737c.
<i>Erebia rossii kuskoquima</i> Holland, 1931.	737d.
<i>Erebia disa mancinus</i> Doubleday and Hewitson, [1849].	738a.
<i>Erebia disa steckeri</i> Holland, 1930.	738b.
<i>Erebia disa subarctica</i> McDunnough, 1937.	738c.
<i>Erebia magdalena magdalena</i> Strecker, 1880.	739a.
<i>Erebia magdalena mackinleyensis</i> Gunder, 1932.	739b.
<i>Erebia fasciata fasciata</i> Butler, 1868.	740a.
<i>Erebia fasciata avinoffi</i> Holland, 1930.	740b.
<i>Erebia fasciata semo</i> Grum-Grshmaïlo, 1899.	740c.
<i>Erebia discoidalis discoidalis</i> (W. Kirby, 1837).	741a.
<i>Erebia discoidalis mcdunnoughi</i> dos Passos, 1940.	741b.
<i>Erebia theano canadensis</i> Warren, 1931.	742a.
<i>Erebia theano alaskensis</i> Holland, 1900.	742b.
<i>Erebia theano ethela</i> W. H. Edwards, 1891.	742c.
<i>Erebia theano demmia</i> Warren, 1936.	742d.
<i>Erebia youngi herscheli</i> Leussler, 1935.	743a.
<i>Erebia youngi youngi</i> Holland, 1900.	743b.
<i>Erebia youngi rileyi</i> dos Passos, 1947.	743c.
<i>Erebia epipsodea rhodia</i> W. H. Edwards, 1871.	744a.
<i>Erebia epipsodea freemani</i> P. Ehrlich, 1954.	744b.
<i>Erebia epipsodea epipsodea</i> Butler, 1868.	744c.
<i>Erebia epipsodea hopfingeri</i> P. Ehrlich, 1954.	744d.
<i>Erebia epipsodea remingtoni</i> P. Ehrlich, 1952.	744e.
<i>Erebia callias</i> W. H. Edwards, 1871.	745.
<i>Erebia lafontainei</i> Troubridge and Philip, "1982(83)".	746.
<i>Erebia occulta</i> Roos and Kimmich, 1983.	746.1.
<i>Erebia inuitica</i> Wyatt, 1966. [see comment in Part II]	746.2.
<i>Gyrocheilus patrobas tritonaria</i> (W. H. Edwards, 1874).	747a.
<i>Neominois ridingsii ridingsii</i> (W. H. Edwards, 1865).	748a.
<i>Neominois ridingsii stretchii</i> (W. H. Edwards, 1870).	748b.

<i>Neominois ridingsii pallidus</i> Austin, 1986.	748c.
<i>Neominois ridingsii minimus</i> Austin, 1986.	748d.
<i>Neominois ridingsii neomexicanus</i> Austin, 1986.	748e.
<i>Oeneis nevadensis gigas</i> Butler, 1868.	750a.
<i>Oeneis nevadensis nevadensis</i> (C. and R. Felder, [1866]).	750b.
<i>Oeneis nevadensis iduna</i> (W. H. Edwards, 1874).	750c.
<i>Oeneis macounii</i> (W. H. Edwards, 1885).	751.
<i>Oeneis chryxus strigulosa</i> McDunnough, 1934.	752a.
<i>Oeneis chryxus calais</i> (Scudder), 1865.	752b.
<i>Oeneis chryxus caryi</i> Dyar, 1904.	752c.
<i>Oeneis chryxus chryxus</i> (Doubleday and Hewitson, [1849]).	752d.
<i>Oeneis chryxus valerata</i> Burdick, [1958].	752e.
<i>Oeneis chryxus stanislaus</i> Hovanitz, 1937.	752f.
<i>Oeneis chryxus ivallda</i> (Mead, 1878).	752g, (749).
<i>Oeneis uhleri varuna</i> (W. H. Edwards), 1882.	753a.
<i>Oeneis uhleri nahanni</i> Dyar, 1904.	753b.
<i>Oeneis uhleri cairnesi</i> Gibson, 1920.	753c.
<i>Oeneis uhleri uhleri</i> (Reakirt, 1866).	753d.
<i>Oeneis uhleri reinthali</i> F. M. Brown, 1953.	753e.
<i>Oeneis alberta alberta</i> Elwes, 1893.	754a.
<i>Oeneis alberta oslari</i> Skinner, 1911.	754b.
<i>Oeneis alberta capulinensis</i> F. M. Brown, 1970.	754c.
<i>Oeneis alberta daura</i> (Strecker, 1894).	754d.
<i>Oeneis taygete taygete</i> Geyer, [1830].	755a.
<i>Oeneis taygete gaspeensis</i> dos Passos, 1949.	755b.
<i>Oeneis taygete fordi</i> dos Passos, 1949.	755c.
<i>Oeneis taygete edwardsi</i> dos Passos, 1949.	755d.
<i>Oeneis bore hanburyi</i> Watkins, 1928.	756a.
<i>Oeneis bore mckinleyensis</i> dos Passos, 1949.	756b.
<i>Oeneis jutta terraenovae</i> dos Passos, 1935.	757a.
<i>Oeneis jutta ascerta</i> Masters and Sorenson, 1968.	757b.
<i>Oeneis jutta ridingiana</i> F. and R. Chermock, 1940.	757c.
<i>Oeneis jutta harperi</i> F. H. Chermock, 1969.	757d.
<i>Oeneis jutta leussleri</i> Bryant, 1935.	757e.
<i>Oeneis jutta alaskensis</i> Holland, 1900.	757f.
<i>Oeneis jutta chermocki</i> Wyatt, 1965.	757g.
<i>Oeneis jutta reducta</i> McDunnough, 1929.	757h.
<i>Oeneis melissa melissa</i> (Fabricius, 1775).	758a.
<i>Oeneis melissa semidea</i> (Say, 1828).	758b.
<i>Oeneis melissa semplei</i> Holland, 1931.	758c.
<i>Oeneis melissa assimilis</i> Butler, 1868.	758d.
<i>Oeneis melissa gibsoni</i> Holland, 1931.	758e.
<i>Oeneis melissa beanii</i> Elwes, 1893.	758f.
<i>Oeneis melissa lucilla</i> Barnes and McDunnough, 1918.	758g.
<i>Oeneis polixenes polixenes</i> (Fabricius, 1775).	759a.
<i>Oeneis polixenes katahdin</i> (Newcomb, 1901).	759b.
<i>Oeneis polixenes subhyalina</i> (Curtis, 1835).	759c.
<i>Oeneis polixenes luteus</i> Troubridge and Parshall, 1988.	759d.
<i>Oeneis polixenes yukonensis</i> Gibson, 1920.	759e.
<i>Oeneis polixenes brucei</i> (W. H. Edwards, 1891).	759f.

<i>Oeneis polixenes woodi</i> Troubridge and Parshall, 1988.	759g.
<i>Oeneis philipi</i> Troubridge, 1988.	759.1.
<i>Oeneis excubitor</i> Troubridge, Philip, Scott and Shepard, 1982.	759.2.

DANAIDAE

<i>Danaus plexippus</i> (Linnaeus, 1758).	760.
<i>Danaus gilippus berenice</i> (Cramer, [1775]).	761a.
<i>Danaus gilippus strigosus</i> (H. W. Bates, 1864).	761b.
<i>Danaus eresimus montezuma</i> Talbot, 1943.	762a.
<i>Danaus eresimus tethys</i> Forbes, 1943.	762b.
<i>Lycorea cleobaea demeter</i> C. and R. Felder, [1865].	763a.
<i>Lycorea cleobaea atergatis</i> Doubleday and Hewitson, [1847].	763b

ITHOMIIDAE

<i>Dircenna klugii</i> (Geyer), 1837.	764.
<i>Greta polissena umbrana</i> (Haensch, 1909).	765a.

INDEX NOTE

As stated in the Introduction on p. vii, no Index is provided for this Supplement to Memoir No. 2. The Checklist entries are keyed to the corresponding entry numbers in this Supplement and the original Memoir No. 2. In this sense, the Checklist serves as a basic Index. The Index to Memoir No. 2 should be used to locate taxa. A list of names added in Memoir No. 3 and not included in Memoir No. 2 begins on page 101. These names are keyed to Part II based upon either species number or genus number. A name set in () is a *nomen nudum*. The List of Added Names does not include misspellings.

Entries in Part Two of this Supplement are arranged by page and species numbers keyed exactly to the original publication. Thus to locate a particular item, the reader should follow the page/species numbers that appear along the left margin of each page in this Supplement. Instructions for using Memoir No. 3 appear in the Introduction on p. vii.

NOTES and ADDITIONS

After the final typesetting of *Memoir No. 3* was completed, the following information was received:

Based upon the revision by R. K. Robbins [Evolution, comparative morphology, and identification of the eumaeinae butterfly genus *Rekoa* Kaye (Lycaenidae: Theclinae), Smithsonian Contr. Zool., in press], on p. 104 of *Memoir No. 2*, *Thereus* Hübner becomes *Rekoa* Kaye, 1904. *Trans. ent. Soc. Lond.* 1904: 198. What have been recorded as *spurina* and *zebina* (misidentified) from our region are both *Rekoa marius* Lucas, of which *spurina* is a junior synonym. *T. zebina* = *R. zebina* is distinct species not found in our area. These changes affect entries 437 (which is deleted), 438 (which becomes *R. marius* Lucas, 1857), and 439 (which becomes *R. palegon*).

130:CLXXIV (new number CLXXIV), add species as follows:

4. *tenedia* C. and R. Felder, 1861. *Wien. ent. Monats.*, 5(4): 99. TL - ?? 543.1.
HT - probably in BM. This species was recorded from southern Texas in 1987, and reported by A. B. Swengel in the Winter 1988 issue of *Wings*, pp. 10-11, published by the Xerces Society. Add also to Checklist.

A			
<i>abebalus</i>	137.5	<i>coloro</i>	303b
<i>acrocnema</i>	582.1	<i>commodus</i>	205.1
<i>adrasta</i>	668.5a	<i>corinna</i>	65.1a
<i>aglaia</i>	569	<i>coronado</i>	596b
<i>alba</i>	698a	<i>curassavica</i>	184
<i>albamontana</i>	175g	D	
<i>albaria</i>	390	<i>datis</i>	688.1a
<i>albidus</i>	424d	<i>diasia</i>	682a
<i>albimedia</i>	74.5	<i>didia</i>	65.1a
<i>alinea</i>	154c	<i>dimidiata</i>	65.1a
<i>alkaliensis</i>	175f	<i>Dircenna</i>	CCXXXIX
<i>amanda</i>	192c	<i>draudti</i>	614
<i>amphichloe</i>	682a	E	
<i>anethi</i>	325a	<i>echemus</i>	697.1
<i>angelika</i>	336	<i>ehrenbergerii</i>	604.1
<i>angelina</i>	345a	"ehrmanni"	320a
<i>anieta</i>	629.5a	<i>ekisi</i>	336
<i>anna</i>	677.1	<i>elissa</i>	234
<i>annonae</i>	300	<i>elko</i>	574w
<i>Antigonus</i>	XXIXA	<i>emorsus</i>	74.5
<i>antillana</i>	682a	<i>Epiphile</i>	CCVIX
<i>apache</i>	355h	<i>eucherus</i>	137.5
<i>argiolus</i>	505	<i>Eucolias</i>	CVIII
<i>artemisia</i>	394d	<i>excubitor</i>	759.2
<i>Arteurotia</i>	XXIA	F	
<i>atlantis</i>	683	<i>fantasos</i>	137.5
<i>azteca</i>	70.1	<i>ferentina</i>	681a
B		<i>fictitia</i>	681a
<i>basinensis</i>	175i	<i>Fixsenia</i>	CXLVII
<i>betulae</i>	648a	<i>flaveola</i>	187a
<i>bilinea</i>	65.1a	"flavolineata"	167
<i>blanco</i>	50d	<i>frechini</i>	310a
<i>blomfildia</i>	688.1	<i>fundania</i>	681a
<i>bobae</i>	65.1a	G	
<i>bogatana</i>	697a	<i>gnaphalii</i>	650
<i>bonnevillus</i>	192f	<i>gracehus</i>	303a
<i>boreale</i>	424b	<i>Greta</i>	CCXL
<i>bottineauensis</i>	728e	<i>guatemalena</i>	684
(<i>browni</i>)	336	<i>guatemalena</i>	684.1
C		<i>gudula</i>	681a
<i>caenus</i>	528	<i>guppyi</i>	336
<i>californicus</i>	7c	H	
<i>canadensis</i>	357.1	<i>harbesoni</i>	217d
<i>capitanensis</i>	574x	"harperi"	624
<i>cervantes</i>	99a	<i>harryi</i>	582b
<i>chalcosiva</i>	458d	<i>heathii</i>	424b
<i>charlestonensis</i>	521d	<i>helios</i>	520m
<i>cinctus</i>	41.1	"herse"	623d
<i>clenchi</i>	458e	<i>hierone</i>	681a
		"hopfingeri"	608b

horni	614	nevadensis	396c
huasteca	140.1	nigrescens	175j
huebneri	205.1	"nigrescens"	623d
(hyatti)	336	"nigrocapitata"	167
hypsipyle	65.1a	nigrozephyrus	645b
I		nivonicus	65.1a
icilia	681a	nobilita	684a
idas	516	O	
insularis	205.1	occulta	746.1
insularis	684a	orbis	376.1
inuitica	746.2	oreocoastus	192e
iphitime	684.2	oregonensis	597e
italica	653a	organensis	475b
J		oroplatia	158o
joannae	684.2a	P	
juvenis	85a	pales	577b
K		pallescens	735d
kluanensis	355g	pallida	175i
klugii	764	pallidus	748c
koreti	634v	panamintensis	312i
(kwakwapoochesi)	310a	pandonius	327a
L		(paradoxa)	448
lafontainei	746	Paramidea	CVII
lagunae	101b	pariaensis	596c
leilia	699	phellea	746.1
lelaps	683a	philipi	759.1
lemhiensis	356c	phyllisae	312h
luestherae	634u	pinjuna	511.1b
luka	629.5a	pinkensis	239a
luteus	759d	plautus	470a
(luxuriosus)	314	plinius	25
M		polissena	765
mackenziei	120a	portensis	205.1
maculata	18	potosiensis	189.2
mandragora	684a	(prestoni)	310a
"marginata"	167	pronus	25
mariamne	562	pulchella	623b
marmorice	684.1a	R	
maynardi	320c	ratonensis	574v
mazai	189.1	reicheli	336
meckyae	336	reinthali	698b
melanica	329c	ribbei	62.5a
melitaeoides	604.2	rogenhoferi	669a
millerorum	452.1	rotundatus	41.1
minimus	748d	"rubrosuffusa"	167
ministigma	175d	rusticus	526.1
muertovalle	151b	S	
N		sangredecresto	589f
negra	657a	santacruzus	192d
neomexicanus	748e	serpentariae	297a

<i>(shapiroi)</i>	336	<i>(tremblayi)</i>	336
<i>sienna fascia</i>	665f	<i>tschukotkensis</i>	588c
<i>sinemaculata</i>	175h	<i>turneri</i>	179
<i>singularis</i>	148	U	
" <i>sodalia</i> "	681a	<i>umbrana</i>	765a
" <i>sternitzkyi</i> "	398	<i>usuli</i>	584c
<i>strophius</i>	440	V	
<i>syraces</i>	126.1	" <i>venusta</i> "	631l
T		<i>Vettius</i>	LIA
<i>Tegosa</i>	CXCVI	<i>victorinus</i>	326.1
<i>tegyra</i>	682	<i>vitellius</i>	205.1
<i>texami</i>	462b	W	
<i>texanus</i>	118c	" <i>weidenhameri</i> "	656
<i>thornei</i>	461.1	<i>wetona</i>	197b
<i>tiliae</i>	636	<i>woodi</i>	759
<i>tithia</i>	675.1	Y	
<i>tractipennis</i>	62.5a	<i>(yukonensis)</i>	355g

